

Pharmacologic Therapy in Acute Renal Failure (ARF)

الترطب

Monitor for Hyperkalemia



Diagnostic Tests

المراقبة

Vitals & Urine Output



Reduce Fluid Overload

تفيم وظلف الكلى

Renal Function Tests



Fluid Out



حافظ على كليتك، تجنب الفشل الكلوي الحاد

Protect Your Kidneys, Prevent ARF

تعويض السوائل

Blood & Fluids



Renal Blood Flow

علاج انخفاض الضغط

Treat Hypotension



العناية القسطرة

Prevent Infections



Nutritional Therapy

- The patient is weighed daily.
- Dietary proteins are limited to about 1 g/kg during the oliguric phase to minimize protein breakdown.
- Caloric requirements are met with high-carbohydrate meals.
- Foods and fluids containing potassium or phosphorus are restricted.

Nursing Management

Monitoring Fluid and Electrolyte Balance

- The nurse monitors the patient's serum electrolyte levels and physical indicators of these complications during all phases of the disorder
- The patient's cardiac function and musculoskeletal status are monitored closely for signs of hyperkalemia.
- The nurse monitors fluid status by paying careful attention to fluid intake.
- Accurate daily weights, as well as intake and output records, are essential.
- Indicators of deteriorating fluid and electrolyte status are reported immediately to the physician

Reducing Metabolic Rate

- Bed rest may be indicated to reduce exertion and the metabolic rate during the most acute stage of the disorder.
- Fever and infection, both of which increase the metabolic rate and catabolism, are prevented or treated promptly.

Promoting Pulmonary Function

- Attention is given to pulmonary function, and the patient is assisted to turn, cough, and take deep breaths frequently.

Preventing Infection

- Asepsis is essential with invasive lines and catheters to minimize the risk of infection and increased metabolism.

Providing skin care

- Massaging bony prominences, turning the patient frequently, and bathing the patient with cool water are often comforting and prevent skin breakdown.

Providing Support

- The patient and family need assistance, explanation, and support during this time.
- The purpose and rationale of the treatments are explained to the patient