

IDP Talk

A Formalisation of Sturm's Theorem  
in Isabelle/HOL

Manuel Eberl  
eberlm@in.tum.de

## Motivation

We have: a polynomial with real coefficients

## Motivation

**We have:** a polynomial with real coefficients

**We want:** the number of real roots in a specific interval

## Motivation

We have: a polynomial with  $\overbrace{\text{real coefficients}}^{\text{not really}}$

We want: the number of real roots in a specific interval

## Motivation

We have: a polynomial with  $\overbrace{\text{real coefficients}}^{\text{not really}}$

We want: the number of real roots in a specific interval

For “real” computations: restricted to appropriate subset of  $\mathbb{R}$ , such as  $\mathbb{Q}$ .

## Motivation

The solution: Sturm's Theorem

Provides a method for counting real roots algorithmically.

$\Rightarrow$  Let's formalise it in Isabelle/HOL

## Notation

Sign changes:  $\sigma(P_0, \dots, P_{n-1}; x)$  denotes denotes the number of sign changes in the sequence  $P_0(x), \dots, P_{n-1}(x)$

For the functionally inclined:

$$\sigma(ps; x) = (\text{length} \circ \text{remdups\_adj} \circ \text{filter } (\neq 0) \circ \text{map } (\lambda p. p(x))) \text{ ps} - 1$$

## Sturm's Theorem

Sturm's Theorem: Let  $P$  be a real polynomial and  $P_0, \dots, P_{n-1}$  a Sturm sequence of  $P$ . Then

$$\sigma(P_0, \dots, P_{n-1}; a) - \sigma(P_0, \dots, P_{n-1}; b)$$

is the number of real roots of  $P$  in the interval  $(a; b]$ .



## Sturm sequence

But what is a Sturm sequence of  $P$ ?

A (reasonably) general definition:

- $P_0 = P$

## Sturm sequence

But what is a Sturm sequence of  $P$ ?

A (reasonably) general definition:

- $P_0 = P$
- $P_0$  and  $P_1$  have no common roots

## Sturm sequence

But what is a Sturm sequence of  $P$ ?

A (reasonably) general definition:

- $P_0 = P$
- $P_0$  and  $P_1$  have no common roots
- $P_{n-1}$  (last element) does not change its sign

## Sturm sequence

But what is a Sturm sequence of  $P$ ?

A (reasonably) general definition:

- $P_0 = P$
- $P_0$  and  $P_1$  have no common roots
- $P_{n-1}$  (last element) does not change its sign
- if  $x_0$  is root of  $P_0$ :  $P_0 P_1(x) < 0$  in some left-NH and  $P_0 P_1(x) > 0$  in some right-NH of  $x_0$

## Sturm sequence

But what is a Sturm sequence of  $P$ ?

A (reasonably) general definition:

- $P_0 = P$
- $P_0$  and  $P_1$  have no common roots
- $P_{n-1}$  (last element) does not change its sign
- if  $x_0$  is root of  $P_0$ :  $P_0P_1(x) < 0$  in some left-NH and  $P_0P_1(x) > 0$  in some right-NH of  $x_0$
- if  $x_0$  is root of another  $P_i$ :  $P_{i-1}P_{i+1}(x_0) < 0$

## Assessment

The good news:

formalisation of real analysis, polynomials, algebra  
already exists

## Assessment

### The good news:

formalisation of real analysis, polynomials, algebra  
already exists

### The bad news:

no formalisation of limits of polynomials, very little on  
divisibility

## Assessment

### The good news:

formalisation of real analysis, polynomials, algebra  
already exists

### The bad news:

no formalisation of limits of polynomials, very little on  
divisibility

### The ugly news:

textbook proofs of Sturm's theorem are extremely  
informal proof sketches at best



## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$   
 $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$   
 $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$   
 $\Rightarrow$  signs  $[1, \_, -1]$  or  $[-1, \_, 1]$  in the entire NH, i.e. one sign change

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$ 
  - $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$
  - $\Rightarrow$  signs  $[1, \_, -1]$  or  $[-1, \_, 1]$  in the entire NH, i.e. one sign change
  - $\Rightarrow$  total number of sign changes not influenced

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$ 
  - $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$
  - $\Rightarrow$  signs  $[1, \_, -1]$  or  $[-1, \_, 1]$  in the entire NH, i.e. one sign change
  - $\Rightarrow$  total number of sign changes not influenced
- if  $P_0$  has  $x_0$  as root,  $P_0P_1(x_0) < 0$  in left-NH of  $x_0$ ,  $> 0$  in right-NH

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$ 
  - $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$
  - $\Rightarrow$  signs  $[1, \_, -1]$  or  $[-1, \_, 1]$  in the entire NH, i.e. one sign change
  - $\Rightarrow$  total number of sign changes not influenced
- if  $P_0$  has  $x_0$  as root,  $P_0P_1(x_0) < 0$  in left-NH of  $x_0$ ,  $> 0$  in right-NH
  - $\Rightarrow$  signs are different left of  $x_0$  and the same right of  $x_0$

## Proving Sturm's Theorem

Assume we already have a Sturm chain. Why does it count roots?  
Follow  $x \mapsto \sigma(P_0, \dots, P_{n-1}; x)$  passing over  $\mathbb{R}$ . Obviously, it can only change at  $x_0$  if one of the  $P_i$  has a root at  $x_0$

- if  $P_i \neq P_0$  has  $x_0$  as a root,  $P_{i-1}P_{i+1}(x_0) < 0$ 
  - $\Rightarrow$  signs of  $P_{i-1}, P_{i+1}$  are  $\neq 0$ , opposite, constant in NH of  $x_0$
  - $\Rightarrow$  signs  $[1, \_, -1]$  or  $[-1, \_, 1]$  in the entire NH, i.e. one sign change
  - $\Rightarrow$  total number of sign changes not influenced
- if  $P_0$  has  $x_0$  as root,  $P_0P_1(x_0) < 0$  in left-NH of  $x_0$ ,  $> 0$  in right-NH
  - $\Rightarrow$  signs are different left of  $x_0$  and the same right of  $x_0$
  - $\Rightarrow$  total number of sign changes decreases by one



## Proving Sturm's Theorem

Formal proof: a lot of induction on the sequences and number of roots  
 $\implies$  messy and not terribly interesting, I'll spare you the details

## Proving Sturm's Theorem

We now know that Sturm sequences can count roots.  
But how do we construct one?

## Construction Sturm sequences

Canonical construction for  $P$  with no multiple roots (i.e.  $\gcd(P, P') = 1$ ):

$$P_i = \begin{cases} P & \text{for } i = 0 \\ P' & \text{for } i = 1 \\ -(P_{i-2} \bmod P_{i-1}) & \text{otherwise} \end{cases}$$

## Construction Sturm sequences

Why does it work? Nonobvious parts:

## Construction Sturm sequences

Why does it work? Nonobvious parts:

If  $x_0$  is root of  $P_0 = P$ :  $PP'(x_0) < 0$  in left-NH and  $PP'(x_0) > 0$  in right-NH

- pick neighbourhood without roots of  $P_0$  and  $P_1$  (except for  $x_0$ ),  
apply mean value theorem

## Construction Sturm sequences

Why does it work? Nonobvious parts:

If  $x_0$  is root of  $P_0 = P$ :  $PP'(x_0) < 0$  in left-NH and  $PP'(x_0) > 0$  in right-NH

- pick neighbourhood without roots of  $P_0$  and  $P_1$  (except for  $x_0$ ),  
apply mean value theorem

If  $x_0$  is root of another  $P_i$ :  $P_{i-1}P_{i+1}(x_0) < 0$  in some NH of  $x_0$

- by construction,  $P_{i-1} = Q \cdot P_i - P_{i+1}$  for some  $Q \in \mathbb{R}[X]$   
 $\implies P_{i-1}(x_0) = -P_{i+1}(x_0)$   
also:  $P_{i-1}(x_0) \neq 0$  since  $\gcd(P_{i-1}, P_i) = \gcd(P_0, P_1) = 1$

## Construction Sturm sequences

This construction assumed no multiple roots.  
What do we do if there are multiple roots?

## Construction Sturm sequences

In case of multiple roots: Let  $D := \gcd(P, P')$ . Then:



## Construction Sturm sequences

In case of multiple roots: Let  $D := \gcd(P, P')$ . Then:

The obvious way:

- compute canonical Sturm chain of  $P/D$   
("divide out" multiple roots)

## Construction Sturm sequences

In case of multiple roots: Let  $D := \gcd(P, P')$ . Then:

The obvious way:

- compute canonical Sturm chain of  $P/D$   
("divide out" multiple roots)

The clever way:

- we can compute the canonical Sturm chain of  $P$   
and divide by  $D$  afterwards

## Construction Sturm sequences

In case of multiple roots: Let  $D := \gcd(P, P')$ . Then:

The obvious way:

- compute canonical Sturm chain of  $P/D$   
("divide out" multiple roots)

The clever way:

- we can compute the canonical Sturm chain of  $P$   
and divide by  $D$  afterwards
- but: if  $D(x) \neq 0$ , dividing by  $D$  does not change the  
number of sign changes at  $x$

## Construction Sturm sequences

In case of multiple roots: Let  $D := \gcd(P, P')$ . Then:

The obvious way:

- compute canonical Sturm chain of  $P/D$   
("divide out" multiple roots)

The clever way:

- we can compute the canonical Sturm chain of  $P$  and divide by  $D$  afterwards
- but: if  $D(x) \neq 0$ , dividing by  $D$  does not change the number of sign changes at  $x$
- $\implies$  unless the interval bounds are multiple roots, we can use the canonical construction without changes

## Making a decision procedure

`count_roots_between p a b`: picks the most efficient Sturm chain construction and:

$$\text{count\_roots\_between } p \ a \ b \ = \ |\{x. a < x \wedge x \leq b \wedge p(x) = 0\}|$$

## Making a decision procedure

count\_roots\_between p a b: picks the most efficient Sturm chain construction and:

$$\text{count\_roots\_between } p \text{ a b} = |\{x. a < x \wedge x \leq b \wedge p(x) = 0\}|$$

Some fluff:

- case distinctions allow arbitrary combination of  $\leq$  and  $<$  in bounds
- “limit signs” allow infinite bounds

In summary: we can count roots in any open/halfopen/closed, bounded/unbounded real interval

## Making a decision procedure

Some more fluff:

- and/or: count  $x$  with

$$P(x) = 0 \wedge Q(x) = 0 \quad \text{or} \quad P(x) = 0 \vee Q(x) = 0$$

## Making a decision procedure

Some more fluff:

- and/or: count  $x$  with

$$P(x) = 0 \wedge Q(x) = 0 \quad \text{or} \quad P(x) = 0 \vee Q(x) = 0$$

- $\forall$ -inequalities:

$$\forall x. P(x) \neq Q(x) \wedge R(x) \neq S(x) \vee T(x) \neq U(x)$$



## Making a decision procedure

Some more fluff:

- and/or: count x with

$$P(x) = 0 \wedge Q(x) = 0 \quad \text{or} \quad P(x) = 0 \vee Q(x) = 0$$

- $\forall$ -inequalities:

$$\forall x. P(x) \neq Q(x) \wedge R(x) \neq S(x) \vee T(x) \neq U(x)$$

- $\forall$  with  $<$  and  $>$ :

$$\forall x. P(x) < Q(x) \wedge R(x) > S(x) \vee T(x) \neq U(x)$$

## Making a decision procedure

### Examples:

lemma "card {x::real.  $(x-1)^2 * (x+1) = 0$ } = 2" **by** sturm

lemma "card {x::real.  $-0.010831 < x \wedge x < 0.010831 \wedge$   
poly [0, -17/2097152, -49/16777216, 1/6, 1/24, 1/120:] x = 0} = 3" **by** sturm

lemma "card {x::real.  $x^3 + x = 2 * x^2 \wedge x^3 - 6 * x^2 + 11 * x = 6$ } = 1" **by** sturm

lemma "(x::real)<sup>2</sup> + 1 > 0" **by** sturm

## Size of the formalisation

3725 LOC in total, 185 of that ML, the rest Isabelle

