


# GENETICS

JANUARY 2022 · VOLUME 220 · GROUP 5



# RNA-Seq Analysis in Bladder Cancer Patients

Emiliano Navarro-Garre<sup>1, 2, †, \*</sup>, Pol Bonet<sup>1, 2, †</sup>, Alberto Carrasco<sup>1, 2, †</sup>, Aina Comas<sup>1, 2, †</sup> and Sofía Redondo<sup>1, 2, †</sup>

<sup>1</sup>Departament de Genètica i de Microbiologia, Facultat de Biociències, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain.

<sup>2</sup>Group 5, Current Topics in Bioinformatics Subject.

<sup>†</sup>These authors contributed equally to this work.

\*Corresponding author: Emiliano Navarro-Garre, Departament de Genètica i de Microbiologia, Facultat de Biociències, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain. Mail: [emiliano.navarro@autonoma.cat](mailto:emiliano.navarro@autonoma.cat)

## Abstract

RNA-seq is a recent approach to carry out expression profiling using high-throughput sequencing technologies, being the preferred option to simultaneously measure the expression of tens of thousands of genes for multiple samples. In this study, we walk through a gene-level RNA-seq differential expression analysis using Bioconductor packages to find genes over- or under-expressed in bladder cancer patients, one of the types of cancer most affected by tobacco use, in an early and advanced tumoral stages, finding a certain number of genes that may be associated with the uncontrolled development of bladder tissue.

**Keywords:** RNA-seq; Bladder cancer; Statistics

## Introduction

Bladder cancer is any of several types of cancer arising from the tissues of the urinary bladder, where the main symptoms are blood in the urine, painful urination and lower back pain and appear when epithelial cells that line the bladder become malignant. Most bladder cancers are diagnosed at an early stage, when the cancer is highly treatable. But even early-stage bladder cancers can come back after successful treatment, due to different risk factors such as smoking, family history, recurrent urinary tract infections, exposure to certain chemicals, or having certain mutation in the genes that are linked to bladder cancer (Board 2002).

In this study we have performed an **RNA-seq analysis**, in order to **detect differentially expressed genes in bladder cancer patients in an early and advanced tumoral stage**, i.e. find genes over- or under-expressed in these cancer patients. RNA-seq is a recent approach to carry out expression profiling using high-throughput sequencing technologies, being the preferred option to simultaneously measure the expression of tens of thousands of genes for multiple samples (Coronado and Carrón 2021).

The aim of this study is to **compare the transcriptomic profile of patients in an early stage with ones in an advanced stage** of this cancer to find the differential expressed genes.

## Methods

All the information about methods and steps has been extracted from the practice script (Coronado and Carrón 2021).

## Packages and tools used


To carry out this study, different **Bioconductor** (Morgan 2021) packages have been used, as it has packages that support high-throughput sequencing data analysis, including RNA-seq. The

packages that were used in this study include core packages maintained by the Bioconductor core team for importing and processing raw sequencing data and loading gene annotations:

- **SummarizedExperiment**: contains one or more assays, each represented by a matrix-like object of numeric or other mode. The rows typically represent genomic ranges of interest and the columns represent samples (Morgan *et al.* 2021).
- **DESeq2**: estimate variance-mean dependence in count data from high-throughput sequencing assays and test for differential expression based on a model using the negative binomial distribution (Love *et al.* 2014).
- **org.Hs.eg.db**: genome wide annotation for Human, primarily based on mapping using Entrez Gene identifiers (Carlson 2021).
- **biomaRt**: provides an interface to a growing collection of databases implementing the BioMart software suite, such as Ensembl (Durinck *et al.* 2009).
- **edgeR**: differential expression analysis of RNA-seq expression profiles with biological replication. Implements a range of statistical methodology based on the negative binomial distributions, including empirical Bayes estimation, exact tests, generalized linear models and quasi-likelihood tests (Robinson *et al.* 2010).
- **tweedEseq**: differential expression analysis of RNA-seq using the Poisson-Tweedie family of distributions (Esnaola *et al.* 2013).
- **GOSTats**: a variety of basic manipulation tools for graphs, hypothesis testing and other simple calculations (Falcon and Gentleman 2007).
- **tweedEseqCountData**: RNA-seq count data employed to illustrate the use of the Poisson-Tweedie family of distributions with the tweedEseq package (Gonzalez and Esnaola










**Figure 2** Gene ontology enrichment analysis of differently expressed genes in late bladder cancer.

### Data availability

The data set used and the code to perform the analyses described in this article are presented below:

-  [Data.](#)
-  [RMarkdown file.](#)
-  [HTML file with the steps of the analysis.](#)

### Literature cited

- Board PATE. 2002. Bladder cancer treatment (pdq®): Patient version. PDQ Cancer Information Summaries. .
- Carlson M. 2021. *org.Hs.eg.db: Genome wide annotation for Human*. R package version 3.14.0.
- Coronado M, Carrón N. 2021. Practice 4: Finding differentially expressed genes in cancer. *Current Topics in Bioinformatics*. .
- Durinck S, Spellman PT, Birney E, Huber W. 2009. Mapping identifiers for the integration of genomic datasets with the r/bioconductor package biomart. *Nature protocols*. 4:1184–1191.
- Esnaola M, Puig P, Gonzalez D, Castelo R, Gonzalez JR. 2013. A flexible count data model to fit the wide diversity of expression profiles arising from extensively replicated rna-seq experiments. *BMC Bioinformatics*. 14:254.
- Falcon S, Gentleman R. 2007. Using GStats to test gene lists for GO term association. *Bioinformatics*. 23:257–8.
- Gentleman R. 2021. *annotate: Annotation for microarrays*. R package version 1.72.0.
- Gonzalez JR, Esnaola M. 2021. *tweeDEseqCountData: RNA-seq count data employed in the vignette of the tweeDEseq package*. R package version 1.32.0.
- Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome Biology*. 15:550.
- Mills BR. 2021. *MetBrewer: Color Palettes Inspired by Works at the Metropolitan Museum of Art*. R package version 0.1.0.
- Morgan M. 2021. *BiocManager: Access the Bioconductor Project Package Repository*. R package version 1.30.16.
- Morgan M, Obenchain V, Hester J, Pagès H. 2021. *SummarizedExperiment: SummarizedExperiment container*. R package version 1.24.0.
- Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. 2008. Mapping and quantifying mammalian transcriptomes by rna-seq. *Nature Methods* 2008 5:7. 5:621–628.
- Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 26:139–140.
- Robinson MD, Oshlack A. 2010. A scaling normalization method for differential expression analysis of rna-seq data. *Genome Biology*. 11:1–9.
- Slowikowski K. 2021. *ggrepel: Automatically Position Non-Overlapping Text Labels with 'ggplot2'*. R package version 0.9.1.
- Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Appendix  
Supplementary Figures




Figure 3 Number of patients in each tumoral stage.




Figure 4 Representation of the 3 different methods to normalize counts.




Figure 5 Differently expressed genes in early vs late bladder cancer




Figure 6 Most differentially expressed genes.