# Duals of Hilbert Spaces; Riesz Representation Theorem; Operators in Hilbert Spaces


| | |
|---|---|
| Ahmed AlOtaibi | 202153590 |
| Abdulaziz AlMutairi | 202168150 |
| Mohammed AlGhudiyan | 202266040 |

King Fahd University of Petroleum and Minerals (KFUPM)

Department of Mathematics and Statistics


MATH 535: Functional Analysis

Dr. Mohammed AlGharabli

March 30, 2026

# Contents


0.1 Riesz Representation Theorem . . . . . 2
0.2 The Adjoint . . . . . 2
0.3 The Adjoint . . . . . 5
0.4 Normal Operators . . . . . 6
0.5 Self-adjoint operators . . . . . 7
0.6 Unitary operators . . . . . 8
0.7 . . . . . 8

- The objectives are:

1-Study the adjoint of an operator
2-Study (Self-odjonit, hormal, unitary operabors)
3-Study the Spectrum of an operater.

## 0.1 Riesz Representation Theorem

**Theorem 1** (Riesz Representation Theorem)**.** *Let $H$ be a Hilbert space. For any bounded linear functional $\varphi \in H^*$, there exists a unique element $y \in H$ such that*

$$\varphi(x) = \langle y, x \rangle$$

*for all $x \in H$ and $\|\varphi\| = \|y\|$*

**Corollary 1.** *Every Hilbert space is reflexive.*

**Corollary 2** (The weak compactness property)**.** *Every bounded sequence in a Hilbert space $H$ containing a weakly convergent subsequence.*

*Proof.* Let $\{ x_n \}$ be a bounded sequence in $H$ with bound $M$, i.e. $\|x_n\| \leq M$, and let $X = \text{span} \{ x_m : m \in \mathbb{N} \}$ . Then the closed unit ball $B[0]_X$ is closed and bounded. By Theorem 2.7 from the course notes it follows that $X$ is finite dimensional. Hence, $X$ is closed. Let $y_n^1 = \langle x_1, x_n \rangle$ be a sequence of real numbers. Since $y_n^1$, by the Bolzano–Weierstrass theorem $y_n^1$ has a convergent subsequence, denote it by $y_{n_k}^1 = \langle x_1, x_{n_k} \rangle$. Similarly, for $y_n^2 = \langle x_2, x_n \rangle$ we obtain its convergent subsequence $y_{n_k}^2 = \langle x_2, x_{n_k} \rangle$.
NOT DONE □

**Remark** The above Corollary is used to prove that
NOT DONE

**Corollary 3.** *Let $T : H \to H^*$ be a bounded linear operator on a Hilbert space $H$ into its dual space $H^*$. Assume that there exists $\alpha > 0$ such that:*

$$(T(x), x) \geqslant \alpha \|x\|^2 \quad \forall x \in H$$

*where $(\cdot, \cdot)$ denotes the duality pairing between $H$ and $H^*$, then $T(x) = f, x \in H$, has a unique solution for all $f \in H^*$.*

## 0.2 The Adjoint

**Definition 1** (Adjoint Operator)**.** *Let $H$ and $G$ be Hilbert spaces and let $T : H \longrightarrow G$ be a bounded linear operator. An operator $T^* : G \to H$ such that*

$$\langle T(x), y \rangle_G = \langle x, T^*(y) \rangle_H \tag{1}$$

*for all $x \in H, y \in G$, is called the adjoint of $T$.*

**Theorem 2.** *For any bounded linear operator $T$ on a Hilbert space $H$, the adjoint operator $T^*$ exists and is unique.*

*Proof.* **Existence of the Adjoint:** We would like to show that for all $y \in G$, there is $w \in H$ such that

$$\langle Tx, y\rangle = \langle x, w\rangle$$

for all $x \in H$. Let

$$\begin{aligned} f_y : H &\longrightarrow \mathbb{F} \\ x &\longmapsto \langle Tx, y\rangle \end{aligned}$$

This map is linear, since it is the composition of two linear maps, namely $T$ and $\langle \cdot, y\rangle$. Also, this map is bounded since

$$\begin{aligned} |f_y(x)| = |\langle Tx, y\rangle| &\leq \|y\|\|Tx\| \\ &\leq \|y\|\|T\|\|x\| \end{aligned}$$

So, $f_y \in H^*$. By Reisz Representation Theorem, for some $w \in H$, we can write $f_y$ as

$$\langle Tx, y\rangle = f_y(x) = \langle x, w\rangle$$

**Uniqueness of the Adjoint:** Let $S : H \to G$ and $S' : H \to G$ be adjoints of $T$, i.e. for all $x, y \in G$

$$\begin{aligned} \langle Tx, y\rangle &= \langle x, Sy\rangle \\ \langle Tx, y\rangle &= \langle x, S'y\rangle \end{aligned}$$

We then have

$$\langle x, (S - S')y\rangle = \langle x, Sy\rangle - \langle x, S'y\rangle = \langle Tx, y\rangle - \langle Tx, y\rangle = 0$$

This implies that $(S - S')y = 0$, and hence, $S - S' = 0 \implies S = S'$. □

The defining relation of the adjoint (1) shows that the action of $A$ on $H_0$ is mimicked by the action of the adjoint $A^*$ on $H_1$; this mimicry is visible only through the applicable inner products, so the adjoint itself depends on these inner products. Roughly, if $A$ has some effect, $A^*$ preserves the geometry of that effect while acting with a reversed domain and codomain.

**Theorem 3.** *Let $H, G$ be Hilbert spaces and let $T : H \to G$ and $S : H \to G$ be linear operator, then*

1. $(T^*)^* = T$
2. $(T + S)^* = T^* + S^*$
3. $(\alpha T)^* = \bar{\alpha} T^*$
4. $(TS)^* = S^* T^*$
5. $T^*(\alpha u + \beta v) = \bar{\alpha} T^*(u) + \bar{\beta} T^*(v)$

*Proof.* 1. We have

$$\langle (T^*)^* x, y\rangle = \overline{\langle y, (T^*)^* x\rangle} = \overline{\langle T^* y, x\rangle} = \langle x, T^* y\rangle = \langle Tx, y\rangle$$

Therefore, $(T^*)^* = T$.

2. We have
$$\begin{aligned}\langle x,(T+S)^*y\rangle = \langle (T+S)x,y\rangle &= \langle Tx,y\rangle + \langle Sx,y\rangle \\ &= \langle x,T^*y\rangle + \langle x,S^*y\rangle \\ &= \langle x,(T^*+S^*)y\rangle\end{aligned}$$
Therefore, $(T+S)^* = T^* + S^*$.

3. We have
$$\begin{aligned}\langle x,(\alpha T)^*y\rangle = \langle \alpha Tx,y\rangle = \alpha\langle Tx,y\rangle = \alpha\langle x,T^*y\rangle &= \alpha\overline{\langle T^*y,x\rangle} \\ &= \overline{\bar{\alpha}\langle T^*y,x\rangle} \\ &= \overline{\langle \bar{\alpha}T^*y,x\rangle} \\ &= \langle x,\bar{\alpha}T^*y\rangle\end{aligned}$$
Therefore, $(\alpha T)^* = \bar{\alpha}T^*$.

4. We have
$$\langle x,(TS)^*y\rangle = \langle TSx,y\rangle = \langle Sx,T^*y\rangle = \langle x,S^*T^*y\rangle$$
Therefore, $(TS)^* = S^*T^*$.

5. We have
$$\begin{aligned}\langle x,T^*(\alpha u+\beta v)\rangle = \langle T(x),\alpha u+\beta v\rangle &= \overline{\langle \alpha u+\beta v,T(x)\rangle} \\ &= \bar{\alpha}\overline{\langle u,T(x)\rangle} + \bar{\beta}\overline{\langle v,T(x)\rangle} \\ &= \bar{\alpha}\langle T(x),u\rangle + \bar{\beta}\langle T(x),v\rangle \\ &= \bar{\alpha}\langle x,T^*(u)\rangle + \bar{\beta}\langle x,T^*(v)\rangle \\ &= \langle x,\bar{\alpha}T^*(u)+\bar{\beta}T^*(v)\rangle\end{aligned}$$
Therefore, $T^*(\alpha u+\beta v) = \bar{\alpha}T^*u + \bar{\beta}T^*v$.

□

**Example 1.** *Let $I$ be the identity operator on a Hilbert space $H$. then $I^* = I$*

**Example 2.** $T_k : C[0,1] \to C[0,1]$
$$g(t) \longmapsto k(t)g(t)$$
*for any $k \in C[0,1]$.*
*then $(T_k)^* = T_{\bar{k}}$*

**Example 3.** *Let $T_1 : H \to H$ be given by $T_1(x) = \alpha x$ for some scalar $\alpha$. For any $x$ and $y$ in $H$,*
$$\langle T_1(x),y\rangle = \langle \alpha x,y\rangle = \alpha\langle x,y\rangle = \langle x,\bar{\alpha}y\rangle$$
*where the first equality follows from the definition of $T_1$, the second from the linearity in the first component of the inner product, and the third from the conjugate linearity in the second component of the inner product. Comparing with (1), the adjoint of $T_1$ is $T_1^*(y) = \bar{\alpha}y$. Hence, the adjoint of multiplication by a scalar is multiplication by the conjugate of the scalar.*

From the above example it follows that if $H$ is a real Hilbert space, then $T_1 = T_1^*$.

**Example 4.** *Let $T : \mathbb{R}^n \to \mathbb{R}^m$ be a linear operator represented by the matrix $A \in \mathbb{R}^{m\times n}$, i.e. $T(x) = Ax$. The matrix of the adjoint of $T$ is the transpose matrix $A^T \in \mathbb{R}^{n\times m}$, since for any $x \in \mathbb{R}^m$ and $y \in \mathbb{R}^n$:*

$$\langle x, T(y)\rangle_{\mathbb{R}^m} = \langle x, Ay\rangle_{\mathbb{R}^m} = x \cdot (Ay) = x^t(Ay) = (x^tA)y = (A^tx)^ty = (A^tx) \cdot y.$$

In component notation, this identity is expressed by swapping the order of summation:

$$\sum_{i=1}^{m} x_i \left(\sum_{j=1}^{n} a_{ij}y_j\right) = \sum_{j=1}^{n} \left(\sum_{i=1}^{m} a_{ij}x_i\right) y_j.$$

For a linear map on complex spaces $T : \mathbb{C}^n \to \mathbb{C}^m$, the matrix of the adjoint with respect to the standard inner product is the Hermitian conjugate matrix, $A^* = \overline{A^T}$, satisfying $\langle x, Ay\rangle = \langle A^*x, y\rangle$.

**Example 5.** *Suppose that $S$ and $T$ are the right and left shift operators, respectively, on the sequence space $l^2(\mathbb{R})$, defined by*

$$S(x_1, x_2, x_3, \dots) = (0, x_1, x_2, x_3, \dots), \quad T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots).$$

*Then $S^* = T$, since*

$$\langle T(x), y\rangle = x_1(0) + x_2y_1 + x_3y_2 + x_4y_3 + \cdots = \langle x, S(y)\rangle.$$

*Also, $T^* = S$, since*

$$\langle S(x), y\rangle = (0)y_1 + x_1y_2 + x_2y_3 + x_3y_4 + \cdots = \langle x, T(y)\rangle.$$

**Theorem 4.** *Suppose that $A : \mathcal{H} \to \mathcal{H}$ is a bounded linear operator on a Hilbert space $\mathcal{H}$ with closed range. Then the equation $Ax = y$ has a solution for $x$ if and only if $y$ is orthogonal to $\ker A^*$.*

**Example 6.** *Consider the multiplication operator $M : L^2([0, 1]) \to L^2([0, 1])$ defined by*

$$Mf(x) = xf(x).$$

*Then $M^* = M$, and $M$ is one-to-one, so every $g \in L^2([0, 1])$ is orthogonal to $\ker M^*$; but the range of $M$ is a proper dense subspace of $L^2([0, 1])$, so $Mf = g$ is not solvable for every $g \in L^2([0, 1])$.*

## 0.3 The Adjoint

**Theorem 5.** *Let $H, G$ be Hilbert spaces, and $T : H \to G$. then*
*ii)* $\|T^*\| = \|T\|$
*iii)* $\|T^*T\| = \|T\|^2$
*iv)* $\ker T = (\operatorname{Im} T^*)^\perp$
*v)* $\ker T^* = (\operatorname{Im} T)^\perp$
*vi)* $\ker T^* = \{0\} \Leftrightarrow \operatorname{Im} T$ *dense in* $G$.

**Theorem 6.** *Let $H$ be a Hilbert space and $T : H \to H$ then $T$ is invertible $\Longleftrightarrow \ker T^* = \{0\}$ and $\exists\alpha > 0$ such that $\|T(x)\| \geqslant \alpha\|x\| \quad \forall x \in H$.*

**Example 7.** *Let $T : l^2 \to l^2$ where*

$$T\left(x_1, x_2 \ldots\right) = (0, x_1, x_2 \ldots)$$

*As shown in Example 5, we have $T^*(x_1, x_2, \ldots) = (x_2, x_3, \ldots)$. So,*

$$\ker T^* = \{u \in \mathbb{F} : (u, 0, 0, \ldots)\} \neq \{(0, 0, 0, \ldots)\}$$

*By Theorem 6, $T$ is not invertible.*

## 0.4 Normal Operators

***Definition 2***. Let $H$ be a Hilbert space and $T$ a lin. operaton $T : H \to H$. then if $TT^* = T^*T, T$ is said to be normal of eratu.

**Example 8.** *Consider*

$$\begin{aligned} T : \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ x &\longmapsto \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} x \end{aligned}$$

*We see that*

$$T^*(x) = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}^T x = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

*Since*

$$\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$$

*we have $T^*T = TT^*$.*

**Example 9.** *If $T$ is the right shift operator defined in Example 5, then $T$ is not normal. This can be shown by finding $T^*T$ and $TT^*$. We have*

$$T^*T(x_1, x_2, \ldots) = T^*(0, x_1, x_2, \ldots) = (x_1, x_2, \ldots)$$

*and*

$$TT^*(x_1, x_2, \ldots) = T(x_2, x_3, \ldots) = (0, x_2, x_3, \ldots)$$

*which are not equal.*

***Example 5***. If $T$ is normal, then $T$ - $\lambda I$ is also normal.

**Lemma 1.** *Let $T : H \to H$ be a normal operator, then*

1. $\|T(x)\| = \|T^*(x)\|\,; \quad \forall x \in H$
2. *If $\|T(x)\| \geqslant \alpha\|x\| \quad \forall x \in H$ then $\ker T^* = \{0\}$. ($\alpha > 0$ ????)*

*Proof.* 1. We have

$$\|Tx\| = \langle Tx, Tx\rangle = \langle x, T^*Tx\rangle = \langle x, TT^*x\rangle = \langle T^*x, T^*x\rangle = \|T^*x\|$$

2. Let $u \in \ker T^*$, i.e. $T^*(u) = 0$. By 1, we have

$$0 = \|T^*u\| = \|Tu\| \geq \alpha\|u\| \geq 0$$

Hence, $\|u\| = 0 \implies u = 0$

□

By the lemma above and theorem 6, the next theorem immediately follows.

**Theorem 7.** *Let $H$ be a Hilbert space and $T : H \to H$ is a normal operator, then $T$ is invertible $\iff \exists \alpha >$ such thar $\|T(x)\| \geqslant \alpha\|x\| \forall x \in H$.*

Hence it is easier to determine whether a normal operator invertible compared with any operator.

## 0.5 Self-adjoint operators

An interesting subset of the normal operators is the self-adjoint operators.

**Definition 2.** *$T : H \to H$ is called a self-adjoint if $T = T^*$*

***Theorem 5***.Let $H$ be Hlibert space & $T_1$ and $T_2$ be self-adjont oterator, then $1 - \alpha T_1 + \beta T_2$ is a self-adjon, $\alpha\beta \in \mathbb{R}$ 2 - The set of self-odjun operuter is a clseel subset of $B(H)$.

**Example 10.** *For any $T \in B(H)$, we observe that $TT^*$ and $T^*T$ are self-adjoint, since*

$$(T^*T)^* = T^*(T^*)^* = T^*T$$

*and*

$$(TT^*)^* = (T^*)^*T^* = TT^*$$

**Proposition 1.** *Any operator $T \in B(H)$, $T = R + iS$ for $R$ and $S$ self-adjoint*

*Proof.* Let $R$ and $S$ be the following

$$R = \frac{T + T^*}{2} \quad S = \frac{T - T^*}{2i}$$

then we have

$$R + iS = \left(\frac{T + T^*}{2}\right) + i\left(\frac{T - T^*}{2i}\right) = \frac{T + T^*}{2} + \frac{T - T^*}{2} = \frac{2T}{2} = T$$

and

$$R^* = \left(\frac{T + T^*}{2}\right)^* = \frac{(T + T^*)^*}{2} = \frac{T^* + (T^*)^*}{2} = \frac{T^* + T}{2} = \frac{T + T^*}{2} = R$$

$$S^* = \left(\frac{T - T^*}{2i}\right)^* = -\frac{(T - T^*)^*}{2i} = -\frac{T^* - (T^*)^*}{2} = -\frac{T^* - T}{2} = \frac{T - T^*}{2} = S$$

□

## 0.6 Unitary operators

**Definition 3.** *An operator $T \in B(H)$ called unitary, if $TT^* = T^*T = I$.*

**Example 11.** *It can be see that the identity operator, $I$, is unitary, since $I^*I = II = I = II = II^*$.*

***Lemma***.If $S, T \in B(H)$ and
$\langle Sz, z\rangle = \langle Tz, 3\rangle \quad \forall z \in H$
$\Rightarrow \quad S = T$.

***Theorem 6***.Let $T \in B(H)$, then $T$ is unitary if $T$ is isometry.

**Theorem 8.** *Let $T, S$ be unitary operators, then*

1. *$T^*$ is unitary and $\|T\| = \|T^*\| = 1$*
2. *$TS$ is unitary*
3. *$T^{-1}$ is unitary.*

**Remark 1.** *By Theorem 8, Example 11 and the associativity of function composition, we observe that the set of all unitary operators forms a group under composition. Moreover, if $T : \mathbb{C}^n \to \mathbb{C}^n$, this group is called the **unitary group**, denoted $U(n)$. There is a special subgroup of $\mathrm{U}(n)$, which is the **special unitary group**, denoted $\mathrm{SU}(n)$, which is the set of all unitary operators from $\mathbb{C}^n$ to $\mathbb{C}^n$ such that its determinant is $1$. This forms a subgroup since if we let $A, B \in \mathrm{SU}(n)$, we have*

$$\det AB^{-1} = \frac{\det A}{\det B} = \frac{1}{1} = 1$$

*showing that $AB^{-1} \in \mathrm{SU}(n)$.*

## 0.7