Duals of Hilbert Spaces; Riesz Representation Theorem; Operators in Hilbert Spaces

Ahmed AlOtaibi	202153590
Abdulaziz AlMutairi	202168150
Mohammed AlGhudiyan	202266040

King Fahd University of Petroleum and Minerals (KFUPM)

Department of Mathematics and Statistics

MATH 535: Functional Analysis

Dr. Mohammed AlGharabli

March 28, 2026

Contents

0.1 Riesz Representation Theorem 2
0.2 The Adjoint 2
0.3 The Adjoint 4
0.4 Normal Operators 6
0.5 Self-adjoint operators 7
0.6 Unitary operators 7
0.7 7

- The objectives are:

1-Study the adjoint of an operator
2-Study (Self-odjonit, hormal, unitary operabors)
3-Study the Spectrum of an operater.

0.1 Riesz Representation Theorem

Theorem 1 (Riesz Representation Theorem)**.** *Let H be a Hilbert space. For any bounded linear functional $\varphi \in H^*$, there exists a unique element $y \in H$ such that*

$$\varphi(x) = \langle y, x \rangle \quad \text{for all } x \in H \text{ and } \|\varphi\| = \|y\|$$

Corollary 1 (The weak compactness property)**.** *Every bounded sequence in a Hilbert space H containing a weakly convergent subsequence.*

Proof. Let $\{ x_n \}$ be a bounded sequence in H with bound M, i.e. $\|x_n\| \leq M$, and let $X = \text{span} \{ x_m : m \in \mathbb{N} \}$. Then the closed unit ball $B[0]_X$ is closed and bounded. By Theorem 2.7 from the course notes it follows that X is finite dimensional. Hence, X is closed. Let $y_n^1 = \langle x_1, x_n \rangle$ be a sequence of real numbers. Since y_n^1, by the Bolzano–Weierstrass theorem y_n^1 has a convergent subsequence, denote it by $y_{n_k}^1 = \langle x_1, x_{n_k} \rangle$. Similarly, for $y_n^2 = \langle x_2, x_n \rangle$ we obtain its convergent subsequence $y_{n_k}^2 = \langle x_2, x_{n_k} \rangle$.
NOT DONE □

Remark The above Corollary is used to prove that
NOT DONE

Corollary 2. *Let $T : H \to H^*$ be a bounded linear operator on a Hilbert space H. Assume that T is strongly monotone, i.e.*

$$\langle T(x), x \rangle \geqslant \alpha \|x\|^2 \quad \forall x \in H \quad (\alpha > 0)$$

then $T(x) = f, x \in H$, has a unique solution for all $f \in H^*$.

0.2 The Adjoint

Definition 1 (Adjoint Operator)**.** *Let H and G be Hilbert spaces and let $T : H \longrightarrow G$ be a bounded linear operator. An operator $T^* : G \to H$ such that*

$$\langle Tx, y \rangle_G = \langle x, T^* y \rangle_H \tag{1}$$

for all $x \in H, y \in G$, is called the adjoint of T.

Proposition 1. *For any bounded linear operator T on a Hilbert space H, the adjoint operator T^* exists and is unique.*

The defining relation of the adjoint (1) shows that the action of A on H_0 is mimicked by the action of the adjoint A^* on H_1; this mimicry is visible only through the applicable inner products, so the adjoint itself depends on these inner products. Roughly, if A has some effect, A^* preserves the geometry of that effect while acting with a reversed domain and codomain.

Example 1 (Adjoint Operators)**.** *Let $T_1 : H \to H$ be given by $T_1(x) = \alpha x$ for some scalar α. For any x and y in H,*

$$\langle T_1(x), y\rangle = \langle \alpha x, y\rangle = \alpha\langle x, y\rangle = \langle x, \bar{\alpha} y\rangle$$

where the first equality follows from the definition of T_1, the second from the linearity in the first component of the inner product, and the third from the conjugate linearity in the second component of the inner product. Comparing with (1), the adjoint of T_1 is $T_1^*(y) = \bar{\alpha} y$. Hence, the adjoint of multiplication by a scalar is multiplication by the conjugate of the scalar.

From the above example it follows that if H is a real Hilbert space, then $T_1 = T_1^*$.

Example 2. *The matrix of the adjoint of a linear map on $\mathbb{R}^n$ with matrix A is A^T since*

$$x \cdot (Ay) = (A^T x) \cdot y.$$

In component notation, we have

$$\sum_{i=1}^{n} x_i \left(\sum_{j=1}^{n} a_{ij} y_j \right) = \sum_{j=1}^{n} \left(\sum_{i=1}^{n} a_{ij} x_i \right) y_j.$$

The matrix of the adjoint of a linear map on $\mathbb{C}^n$ with complex matrix A is the Hermitian conjugate matrix, $A^* = \overline{A^T}$.

Example 3. *Suppose that S and T are the right and left shift operators, respectively, on the sequence space $l^2(\mathbb{R})$, defined by*

$$S(x_1, x_2, x_3, \dots) = (0, x_1, x_2, x_3, \dots), \quad T(x_1, x_2, x_3, \dots) = (x_2, x_3, x_4, \dots).$$

Then $S^* = T$, since

$$\langle T(x), y\rangle = x_1(0) + x_2 y_1 + x_3 y_2 + x_4 y_3 + \cdots = \langle x, S(y)\rangle.$$

Also, $T^* = S$, since

$$\langle S(x), y\rangle = (0) y_1 + x_1 y_2 + x_2 y_3 + x_3 y_4 + \cdots = \langle x, T(y)\rangle.$$

Example 4 (Local Averaging and its Adjoint)**.** *The operator A given by*

$$(Ax)_n = \int_{n-1/2}^{n+1/2} x(t)dt, \quad n \in \mathbb{Z},$$

takes local averages of the function $x(t)$ to yield a sequence $(Ax)_n$. A is a bounded linear operator from $\mathcal{L}^2(\mathbb{R})$ to $l^2(\mathbb{Z})$. For the adjoint:

$$\langle Ax, y\rangle_{l^2} = \sum_{n\in\mathbb{Z}} (Ax)_n y_n^* = \sum_{n\in\mathbb{Z}} \left(\int_{n-1/2}^{n+1/2} x(t)dt \right) y_n^* = \int_{-\infty}^{\infty} x(t) \left(\sum_{n\in\mathbb{Z}} y_n 1_{[n-1/2, n+1/2)}(t) \right)^* dt$$

$$= \langle x, A^* y \rangle_{\mathcal{L}^2}.$$

The adjoint of local averaging forms a piecewise-constant function:

$$(A^* y)(t) = \sum_{n \in \mathbb{Z}} y_n 1_{[n-1/2, n+1/2)}(t).$$

Theorem 2. *Suppose that $A : \mathcal{H} \to \mathcal{H}$ is a bounded linear operator on a Hilbert space $\mathcal{H}$ with closed range. Then the equation $Ax = y$ has a solution for x if and only if y is orthogonal to* $\ker A^*$.

Example 5. *Consider the multiplication operator $M : L^2([0,1]) \to L^2([0,1])$ defined by*

$$Mf(x) = xf(x).$$

Then $M^* = M$, and M is one-to-one, so every $g \in L^2([0,1])$ is orthogonal to $\ker M^*$*; but the range of M is a proper dense subspace of $L^2([0,1])$, so $Mf = g$ is not solvable for every $g \in L^2([0,1])$.*

0.3 The Adjoint

Definition 2. *Let K and G be Hilbert spaces and let $T : H \longrightarrow G$ be a linear operator. An operator $T^* : G \to H$ such that*

$$\langle Tx, y \rangle_G = \langle x, T^* y \rangle_H \qquad \text{for all } x \in \mathcal{H}, y \in G$$

*is called **the adjoint** of T.*

Theorem 3. *For any linear operator $T : H \to G$. the adjoint T^* exists and is unique.*

Proof. **Existence of the Adjoint:** We would like to show that for all $y \in G$, there is $w \in H$ such that

$$\langle Tx, y \rangle = \langle x, w \rangle$$

for all $x \in H$. Let

$$\begin{aligned} f_y : H &\longrightarrow \mathbb{F} \\ x &\longmapsto \langle Tx, y \rangle \end{aligned}$$

This map is linear, since it is the composition of two linear maps, namely T and $\langle \cdot, y \rangle$. Also, this map is bounded since

$$\begin{aligned} |f_y(x)| = |\langle Tx, y \rangle| &\leq \|y\| \|Tx\| \\ &\leq \|y\| \|T\| \|x\| \end{aligned}$$

So, $f_y \in H^*$. By Reisz Representation Theorem, for some $w \in H$, we can write f_y as

$$\langle Tx, y \rangle = f_y(x) = \langle x, w \rangle$$

Uniqueness of the Adjoint: Let $S : H \to G$ and $S' : H \to G$ be adjoints of T, i.e. for all $x, y \in G$

$$\begin{aligned} \langle Tx, y \rangle &= \langle x, Sy \rangle \\ \langle Tx, y \rangle &= \langle x, S'y \rangle \end{aligned}$$

We then have

$$\langle x, (S - S')y\rangle = \langle x, Sy\rangle - \langle x, S'y\rangle = \langle Tx, y\rangle - \langle Tx, y\rangle = 0$$

This implies that $(S - S')y = 0$, and hence, $S - S' = 0 \implies S = S'$. $\square$

Example 6. *Let I be the identity operator on a Hilbert space H. then $I^* = I$*

Example 7. $T_k : C[0,1] \to C[0,1]$

$$g(t) \longmapsto k(t)g(t)$$

for any $k \in C[0,1]$.
then $(T_k)^* = T_{\bar{k}}$

Theorem 4. *Let H, G be Hilbert spaces and let $T : H \to G$ and $S : H \to G$ be linear operator, then*

1. $(T^*)^* = T$
2. $(T + S)^* = T^* + S^*$
3. $(\alpha T)^* = \bar{\alpha} T^*$
4. $(TS)^* = S^* T^*$
5. $T^*(\alpha u + \beta v) = \bar{\alpha} T^*(u) + \bar{\beta} T^*(v)$

Proof.

1. We have

$$\langle (T^*)^* x, y\rangle = \overline{\langle y, (T^*)^* x\rangle} = \overline{\langle T^* y, x\rangle} = \langle x, T^* y\rangle = \langle Tx, y\rangle$$

Therefore, $(T^*)^* = T$.

2. We have

$$\begin{aligned}\langle x, (T+S)^* y\rangle = \langle (T+S)x, y\rangle &= \langle Tx, y\rangle + \langle Sx, y\rangle \\ &= \langle x, T^* y\rangle + \langle x, S^* y\rangle \\ &= \langle x, (T^* + S^*)y\rangle\end{aligned}$$

Therefore, $(T + S)^* = T^* + S^*$.

3. We have

$$\begin{aligned}\langle x, (\alpha T)^* y\rangle = \langle \alpha Tx, y\rangle = \alpha\langle Tx, y\rangle = \alpha\langle x, T^* y\rangle &= \alpha\overline{\langle T^* y, x\rangle} \\ &= \overline{\bar{\alpha}\langle T^* y, x\rangle} \\ &= \overline{\langle \bar{\alpha} T^* y, x\rangle} \\ &= \langle x, \bar{\alpha} T^* y\rangle\end{aligned}$$

Therefore, $(\alpha T)^* = \bar{\alpha} T^*$.

4. We have

$$\langle x, (TS)^* y\rangle = \langle TSx, y\rangle = \langle Sx, T^* y\rangle = \langle x, S^* T^* y\rangle$$

Therefore, $(TS)^* = S^* T^*$.

5. We have

$$\begin{aligned}\langle x, T^*(\alpha u+\beta v)\rangle = \langle T(x), \alpha u+\beta v\rangle &= \overline{\langle \alpha u+\beta v, T(x)\rangle} \\ &= \bar{\alpha}\overline{\langle u, T(x)\rangle} + \bar{\beta}\overline{\langle v, T(x)\rangle} \\ &= \bar{\alpha}\langle T(x), u\rangle + \bar{\beta}\langle T(x), v\rangle \\ &= \bar{\alpha}\langle x, T^*(u)\rangle + \bar{\beta}\langle x, T^*(v)\rangle \\ &= \langle x, \bar{\alpha}T^*(u) + \bar{\beta}T^*(v)\rangle\end{aligned}$$

Therefore, $T^*(\alpha u+\beta v) = \bar{\alpha}T^*u + \bar{\beta}T^*v$.

□

Theorem 5. *Let H, G be Hilbert spaces, and $T : H \to G$. then*
ii) $\|T^*\| = \|T\|$
iii) $\|T^*T\| = \|T\|^2$
iv) $\ker T = (\operatorname{Im} T^*)^\perp$
v) $\ker T^* = (\operatorname{Im} T)^\perp$
vi) $\ker T^* = \{0\} \Leftrightarrow \operatorname{Im} T$ *dense in* G.

Theorem 6. *Let H be a Hilbert space and $T : H \to H$ then T is invertible* $\Longleftrightarrow \ker T^* = \{0\}$ *and* $\exists \alpha > 0$ *such that* $\|T(x)\| \geqslant \alpha\|x\| \quad \forall x \in H$.

Theorem 7. *Let $T : l^2 \to l^2$ Where*
$T(x_1, x_2 \ldots) = (0_1, x_1, x_2 \ldots)$
T is not invertible.

0.4 Normal Operators

Definition 2. Let H be a Hilbert space and T a lin. operaton $T : H \to H$. then if $TT^* = T^*T, T$ is said to be normal of eratu.

Example 8. *Consider*

$$\begin{aligned} T : \mathbb{R}^2 &\longrightarrow \mathbb{R}^2 \\ x &\longmapsto \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} x \end{aligned}$$

We see that

$$T^*(x) = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}^T x = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} x$$

Since

$$\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$$

we have $T^*T = TT^*$.

Example 4.T is Exp3 is not normal.

Example 5. If T is normal, then T - λI is also normal.

Lemma.Let $T : H \to H$ be a normal operator, then
a) $\|T(x)\| = \|T^*(x)\|\,; \quad \forall x \in H$
b) If $\|T(x)\| \geqslant \alpha\|x\| \quad \forall x \in H$ then $\ker T^* = \{0\}$.

Theorem 4. Let H be a Hilbert space and $T : H \to H$ is a normal operator, then
T is invertible $\Longleftrightarrow \exists \alpha >$ such thar $\quad \|T(x)\| \geqslant \alpha\|x\| \forall x \in H$.
Hence it is easier to determine whether a normal operator invertible compared with any operator.

0.5 Self-adjoint operators

An interesting subset of the normal operators is the self-adjoint operators.
Definition 3. $T : H \to H$ is called a self-adjoint if $T = T^*$
Theorem 5.Let H be Hlibert space & T_1 and T_2 be self-adjont oterator, then $1 - \alpha T_1 + \beta T_2$ is a self-adjon, $\alpha\beta \in \mathbb{R}$ 2 - The set of self-odjun operuter is a clseel subset of $B(H)$.
Example 6. For any $T \in B(H)$, show TT^* and T^*T are self-adjoint.

Example 7. Any operator $T \in B(H)$, $T = R + iS$ for $R\&S$ self-adjoint

0.6 Unitary operators

Definition 4. An operator $T \in B(H)$ called unitary, if $\Pi T^* = T^*T = I$.
Lemma.If $S, T \in B(H)$ and
$\langle Sz, z\rangle = \langle Tz, 3\rangle \quad \forall z \in H$
$\Rightarrow \quad S = T$.
Theorem 6.Let $T \in B(H)$, then T is unitary if T is isometry.
Theorem 7. let T, S be unitary opearturs, then
a) T^* is unitary & $\|T\| = \|T^*\| = 1$
b) TS is unitary
c) T^{-1} is unitary.

0.7