

The Spectrum of an Operator

Abdulaziz AlMutairi

April 27, 2026

Outline

Introduction

Outline

Introduction

Spectrum

Outline

Introduction

Spectrum

Examples of Spectra of Operators

Outline

Introduction

Spectrum

Examples of Spectra of Operators

Definition

Let T be a linear operator, $\lambda \in \mathbb{C}$ is called an eigenvalue of T , if there exists $x \in H$ with $x \neq 0$ such that

$$Tx = \lambda x$$

Theorem

Let H be a Hilbert space and $T : H \rightarrow H$ then T is invertible $\iff$ $\ker T^ = \{0\}$ and $\exists \alpha > 0$ such that $\|T(x)\| \geq \alpha \|x\| \quad \forall x \in H$.*

Theorem

Let H be a Hilbert space and $T : H \rightarrow H$ then T is invertible $\iff \ker T^ = \{0\}$ and $\exists \alpha > 0$ such that $\|T(x)\| \geq \alpha \|x\| \quad \forall x \in H$.*

Corollary

Let H be a Hilbert space and $T : H \rightarrow H$ then T is invertible $\iff \exists \beta > 0$ such that $\|T^(x)\| \geq \beta \|x\|$ and $\exists \alpha > 0$ such that $\|T(x)\| \geq \alpha \|x\| \quad \forall x \in H$.*

Proof.

If T is invertible, then so is T^* . So, by Theorem 1, $\exists \beta > 0$ such that $\|T^*(x)\| \geq \beta\|x\|$ and $\exists \alpha > 0$ such that $\|T(x)\| \geq \alpha\|x\|$ for all $x \in H$. Conversely, if $\exists \beta > 0$ such that $\|T^*(x)\| \geq \beta\|x\|$, then if $T^*u = 0$ for some $u \in H$, then

$$\begin{aligned} 0 &= \|T^*u\| \geq \beta\|u\| \\ &\implies u = 0 \end{aligned}$$

So, $\ker T^* = \{0\}$. By Theorem 1, T is invertible. □

Outline

Introduction

Spectrum

Examples of Spectra of Operators

Definition

Let T be a linear operator. The resolvent set of T , denoted by $\rho(T)$, is $\rho(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is invertible}\}$. The spectrum of T , denoted by $\sigma(T)$ is $\sigma(T) = \mathbb{C} \setminus \rho(T)$, or equivalently, $\sigma(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible}\}$

Remark

Notice that, if $T \in B(H)$, by Theorem 1, we can rewrite the resolvent set of T as

$$\rho(T) = \{\lambda \in \mathbb{C} : \ker(T - \lambda I)^* = \{0\} \\ \text{and } \|(T - \lambda I)x\| \geq \alpha\|x\| \text{ for some } \alpha > 0\}$$

Alternatively, we can rewrite the resolvent set of T as

$$\rho(T) = \{\lambda \in \mathbb{C} : \|(T - \lambda I)^*x\| \geq \beta\|x\| \\ \text{and } \|(T - \lambda I)x\| \geq \alpha\|x\| \text{ for some } \beta, \alpha > 0\}$$

by Corollary 1

Theorem

If λ is an eigenvalue of T , then $\lambda \in \sigma(T)$

Proof.

Suppose that $\lambda \notin \sigma(T)$, then $T - \lambda I$ is invertible. So, we have $(T - \lambda I)u = 0 \implies u = (T - \lambda I)^{-1}(T - \lambda I)u = (T - \lambda I)^{-1}(0) = 0$. So, if there is $u \in \mathcal{H}$ such that $Tu = \lambda u$, then $u = 0$. So, λ is not an eigenvalue of T . □

Theorem

Let H be a finite dimensional Hilbert space, and let $T : H \rightarrow H$ be linear. If $\lambda \in \sigma(T)$, then λ is an eigenvalue of T .

Remark

For finite dimensional Hilbert space, the spectrum of T consist only the eigenvalues of T . However, in infinite dimensional Hilbert space, we may have no eigenvalues. For instance, consider the shift operator. Here, it has no eigenvalues, since if we consider $(T - \lambda I)x = 0$, we see that

$$(-\lambda x_1, x_1 - \lambda x_2, x_2 - \lambda x_3, \dots) = (0, 0, 0, \dots) \implies x = (0, 0, 0, \dots)$$

So, T does not have an eigenvalue. However, we showed that T is not invertible. So, $T - 0I$ is not invertible and $0 \in \sigma(T)$. So, the spectrum of an operator does not only consist of eigenvalues.

Proof.

Suppose that λ is not an eigenvalue of T , then there is nonzero $u \in H$ such that $Tu = \lambda u$. So, there is no nonzero $u \in H$ such that $(T - \lambda I)u = 0$. So, $\ker(T - \lambda I) = \{0\}$. By rank-nullity theorem, we have

$$\begin{aligned}\dim H &= \dim \ker(T - \lambda I) + \dim \operatorname{Im}(T - \lambda I) \\ &= 0 + \dim \operatorname{Im}(T - \lambda I) \\ &= \dim \operatorname{Im}(T - \lambda I)\end{aligned}$$

So, $\operatorname{Im}(T - \lambda I) = H$. This means that $(T - \lambda I)$ is surjective, and therefore, invertible. So, $\lambda \in \rho(T)$, and therefore, $\lambda \notin \sigma(T)$. $\square$

Proposition

Let $\mathcal{H}$ be a Hilbert space. Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be linear operator. The following properties hold:

1. $\sigma(I) = \{1\}$
2. $\sigma(T^*) = \{\bar{\lambda} : \lambda \in \sigma(T)\}$
3. If T is invertible, then $\sigma(T^{-1}) = \{\mu^{-1} : \mu \in \sigma(T)\}$
4. If T is self-adjoint, then $\sigma(T) \subseteq \mathbb{R}$.
5. If $U : \mathcal{H} \rightarrow \mathcal{H}$ is invertible, then $\sigma(UTU^{-1}) = \sigma(T)$

Lemma

If $\mathcal{H}$ is a complex Hilbert space and $U \in B(\mathcal{H})$ is unitary then

$$\sigma(U) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\}$$

Theorem

Let $\mathcal{H}$ be a complex Hilbert space and let $T \in B(\mathcal{H})$. If $|\lambda| > \|T\|$ then $\lambda \notin \sigma(T)$.

Corollary

If $T \in B(H)$, then $\sigma(T)$ is bounded.

Proof.

If $\lambda \in \sigma(T)$, then, by the contrapositive of Theorem 4, $|\lambda| \leq \|T\|$.
So, $\lambda \in \overline{B(0, \|T\|)}$. Hence, $\sigma(T) \subseteq \overline{B(0, \|T\|)}$, i.e. $\sigma(T)$ is bounded. □

Theorem

If $T \in B(H)$, then $\sigma(T) \subseteq \mathbb{C}$ is a closed set.

Corollary

If $T \in B(H)$, then $\sigma(T)$ is compact in $\mathbb{C}$

Outline

Introduction

Spectrum

Examples of Spectra of Operators

We would like to find the spectrum of the unilateral shift, $T : \ell^2 \rightarrow \ell^2$, defined by

$$T(x_1, x_2, \dots) = (0, x_1, x_2, \dots)$$

Since T has been shown to have no eigenvalue, we will consider the eigenvalues of T^* instead and use the properties of the spectrum to have a lower bound for the spectrum.

Suppose that $|\lambda| < 1$, then consider the sequence

Suppose that $|\lambda| < 1$, then consider the sequence

$$x = (1, \lambda, \lambda^2, \dots)$$

Suppose that $|\lambda| < 1$, then consider the sequence

$$x = (1, \lambda, \lambda^2, \dots)$$

Claim: $x \in \ell^2$

Suppose that $|\lambda| < 1$, then consider the sequence

$$x = (1, \lambda, \lambda^2, \dots)$$

Claim: $x \in \ell^2$

We have

$$\sum_{n=0}^{\infty} |x_n|^2 = \sum_{n=0}^{\infty} (|\lambda|^n)^2 = \sum_{n=0}^{\infty} (|\lambda|^2)^n = \frac{1}{1 - |\lambda|^2} < \infty$$

Now, observe that

$$\begin{aligned} T_X &= T(1, \lambda, \lambda^2, \dots) \\ &= (\lambda, \lambda^2, \lambda^3, \dots) = \lambda(1, \lambda, \lambda^2, \dots) = \lambda x \end{aligned}$$

Now, observe that

$$\begin{aligned}Tx &= T(1, \lambda, \lambda^2, \dots) \\ &= (\lambda, \lambda^2, \lambda^3, \dots) = \lambda(1, \lambda, \lambda^2, \dots) = \lambda x\end{aligned}$$

So, every $\lambda \in \mathbb{C}$ such that $|\lambda| < 1$ is an eigenvalue of T^* .

Now, observe that

$$\begin{aligned}Tx &= T(1, \lambda, \lambda^2, \dots) \\ &= (\lambda, \lambda^2, \lambda^3, \dots) = \lambda(1, \lambda, \lambda^2, \dots) = \lambda x\end{aligned}$$

So, every $\lambda \in \mathbb{C}$ such that $|\lambda| < 1$ is an eigenvalue of T^* . Now, we have

$$\{\lambda : |\lambda| < 1\} = \{\bar{\lambda} : |\lambda| < 1\} \subseteq \sigma(T)$$

So,

$$\{\lambda \in \mathbb{C} : |\lambda| < 1\} \subseteq \sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}$$

So,

$$\{\lambda \in \mathbb{C} : |\lambda| < 1\} \subseteq \sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}$$

or

$$B(0, 1) \subseteq \sigma(T) \subseteq \overline{B(0, 1)}$$

So,

$$\{\lambda \in \mathbb{C} : |\lambda| < 1\} \subseteq \sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}$$

or

$$B(0,1) \subseteq \sigma(T) \subseteq \overline{B(0,1)}$$

Since $\sigma(T)$ is closed, we have

$$\sigma(T) = \overline{B(0,1)} = \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}$$