

Plants And Soil Microorganisms: Removal of Formaldehyde, Xylene, and Ammonia from the Indoor Environment

B. C. Wolverton and John D. Wolverton
Wolverton Environmental Services
514 Pine Grove Road
Picayune, Mississippi 39466

Abstract

Interior plants in potting soil and potting soil without plants were evaluated for their ability to remove formaldehyde, xylene, and ammonia from sealed chambers. Interior paneling sections made of particle board were also used as a continuous out-gassing source of formaldehyde. Over thirty interior plants were tested and *Nephrolepis exaltata* "Bostoniensis" (Boston fern), *Chrysanthemum morifolium* (pot mum), and *Phoenix roebelenii* (dwarf date palm) were the most effective in removing formaldehyde with 1863 µg, 1450 µg, and 1385 µg removed per hour, respectively. The dwarf date palm was the most effective plant in removing xylene with 610 µg removed per hour. *Rhapis excelsa* (lady palm) was one of the most effective plants in removing ammonia with 7,356 µg removed per hour. The Boston fern was also the most effective plant in continuously removing out-gassing formaldehyde from particle board in a sealed chamber. This data indicates that house plants such as ferns, pot mums, and palms may be a cost effective means of improving the indoor air quality in tightly constructed facilities ranging from mobile homes to high-rise office facilities. Microorganisms maintained by the plants in the rhizosphere and plant leaves both appear to be important in removing indoor air polluting organic chemicals and improve with exposure time.

Introduction.

Most materials found in modern buildings,

including mobile homes, emit hundreds of volatile organic chemicals (VOCs), such as formaldehyde. Indoor air polluting chemicals come from sources such as paints, adhesives, carpeting, upholstery, furniture, paneling, plastic, vinyl, copying machines, computers, cleaning agents, and hundreds of other products found inside offices, hospitals, homes, and other buildings. Although the indoor air pollution levels are generally highest in new and renovated buildings, out-gassing from furniture, paneling and other materials may continue at trace levels for years. Some personnel exposed to VOCs in new buildings and mobile homes may not demonstrate acute reactions immediately, but become sensitized as a result of exposure. However, they may later demonstrate acute reactions when again exposed to trace levels of these chemicals.

In August, 1989, the U.S. Environmental Protection Agency (EPA) submitted a report to Congress on the quality of indoor air found in ten public access buildings, including hospitals and office buildings. This report stated that more than 900 organic chemicals have been identified in newly constructed buildings and that some chemical levels were one hundred times greater than normal levels. This report also stated that sufficient evidence exists to conclude that indoor air pollution may pose serious acute and chronic health risks (EPA, 1989).

Before the 1973-74 energy crisis, building ventilation standards called for approximately 0.42 cubic meters per minute (15 cfm) of outside air for each building

occupant. As a result of the 1973 oil embargo, national energy conservation measures called for a reduction in the amount of outside air provided for ventilation to approximately 0.14 cubic meters per minute (5 cfm) per occupant. This is believed to be a contributing factor in the deteriorating indoor air quality and the development of "sick building syndrome," which had become widespread by 1984. The building industry is presently in a dilemma over how to maintain tightly sealed, energy-efficient buildings without sacrificing indoor air quality.

The first evidence that office and house plants, along with the microorganisms associated with their roots, were capable of removing indoor air polluting chemicals was demonstrated by Wolverton et al., 1984 and 1985. The ability of soil microorganisms to degrade toxic organic chemicals has been known for many years (Davies and Evans, 1964; Evans et al., 1965; Tabak et al., 1981; LaPat-Polaske et al., 1984; Saber et al., 1985; Nelson et al., 1986). The ability of plants to exude substances from their roots which stimulate the growth of some microbes while inhibiting the growth of others on and around their roots, an area called the rhizosphere, is a well-known phenomenon (Rovira, 1959, 1963, 1970; Katznelson, 1970; Rovira and Davey, 1974). Wolverton and Harrison (1973) demonstrated the importance of this phenomenon with aquatic plants in removing toxic insecticides from water. When water contaminated with a highly toxic insecticide was exposed to soil

containing *Nymphaea odorata* (water lily) and *Juncus repens* (rush), the insecticide was detoxified rapidly in the water containing the water lily but remained toxic in the water containing the rush. This paper presents the results of additional studies demonstrating the ability of house and office plants and their associated microorganisms to remove formaldehyde, xylene, and ammonia from sealed chambers.

Materials and Methods.

Clear, cubical, plastic chambers with a volume of approximately 310 liters were used to maintain the plants in a sealed environment during test periods. The chamber tops were removable and fitted with a gasket, bolts and wing nuts to provide an airtight seal. Plant grow lights, equipped with timers to control light and dark cycles, were mounted outside the chambers to prevent heat build-up inside the chambers. All experiments of 24 hours or greater were conducted using 12 hour dark:12 hour light cycles. Light levels of 1150 ± 50 lux were maintained during the light cycle. Small electrical fans and temperature and relative humidity measuring devices were installed inside each chamber. Portholes fitted with airtight septa were used to seal the fans' electrical cords and remove air samples. The air inside the chambers was contaminated with formaldehyde by pumping air into the chamber via a gas scrubbing apparatus filled with 250 mL of a 37% formaldehyde solution. Ammonia was introduced via the same gas scrubbing apparatus filled with 250 mL of a commercial ammonia cleaning solution. The chambers were contaminated with xylene by placing one drop of xylene

on a piece of paper towel and placing the paper towel inside each chamber. The small air circulating fan was operated for approximately thirty minutes before the first samples were taken for xylene. Before taking formaldehyde and ammonia samples, the circulation fans were operated for approximately five minutes.



To evaluate the ability of plants to continuously remove out-gassing formaldehyde from paneling, furniture and other products manufactured with particle board, small sections ($30.5 \times 25.4 \times 0.6$ cm) of interior paneling obtained from a local building material supplier were placed inside two chambers. The two chambers were identical, except that one chamber contained a plant with the paneling section and the other chamber contained only a beaker of water with the paneling section. Sampling was conducted using a Sensidyne-Gastec air sampling pump and gas detector tubes specific for formaldehyde, xylene, and ammonia. The formaldehyde, xylene, and ammonia tubes had lower detection levels of 0.05 ppm, 1.0 ppm and 1.0 ppm, respectively. Soil samples were analyzed for bacteria by means of the pour plate technique to determine the number of "colony forming units" per gram of soil (cfu/g). Plate Count Agar (PCA) was used as the microbiological media and incubated at 26°C . Gram-positive and gram-negative bacteria were determined by using standard gram stain methods. Baccto soil from the Michigan Peat Company was used for the comparison studies with plants and soil controls. Potting soil controls were sterilized by using steam pressure sterilization at 6.8 kg at 121°C for 20 minutes. Sand used to cover the potting soil surface with and without

plants was also sterilized. All tests were repeated three or more times, including leak and chamber sorption controls.

Plants listed in Tables 1, 2, and 3 were standard nursery stock and tested over a five-month period. Data represents mean averages of results obtained during the five-month test period. Plant heights in the 35.6 cm pots were 82-90 cm. Plant heights in the 25.4 cm pots and in the 20.3 cm pots were 45-66 cm, while the heights of plants in the 15.2 cm pots were 36-46 cm.

Results and Discussion.

The most effective house plant tested for removing formaldehyde from sealed chambers was *Nephrolepis exaltata* "Bostoniensis" (Boston fern). This plant removed 1863 μg of formaldehyde per hour from a sealed chamber (Table 1). The U.S. Environmental Protection Agency (EPA) found mean indoor concentrations of formaldehyde and xylene in newly constructed office buildings to be 0.173 μg and 0.022 μg per liter of air, respectively (EPA, 1988). At this contamination level, the air in a 9.3 m² office with a 2.4 m ceiling would contain 3916 μg of formaldehyde and 493 μg of xylene. At a removal rate of 1863 $\mu\text{g/hr}$, two Boston ferns should be capable of removing the formaldehyde from the air in this office. Based on data from Table 1, approximately three *Dracaena deremensis* (Janet Craigs) would be required to remove the same level of formaldehyde from office areas. With a xylene contamination level of 493 $\mu\text{g/office}$, two Boston ferns or three Janet Craigs should also remove xylene from the same office area (Table 2). The numbers and varieties of other plants required to remove formaldehyde, xylene, and ammonia from



contaminated air can be determined from Tables 1, 2, and 3.

Although ammonia has not been identified by EPA as a serious indoor air pollution problem, it is sometimes used to treat particle board and interior paneling to reduce formaldehyde levels. Therefore, it could in the future present indoor air pollution problems.

To determine the formaldehyde removal rates of potting soil without plants and the importance of bacteria in removing formaldehyde, sterilized and unsterilized soils were compared. As shown in Table 4, sterilized soil and unsterilized soil covered with sterilized sand did not remove detectable levels of formaldehyde from sealed chambers; whereas, exposed unsterilized soil that had been in pots for several months removed 188 µg/hr of formaldehyde. Fresh potting soil without plants and the same type potting soil with *Ficus benjamina*, *Spathiphyllum* sp., *Sansevieria* sp., and *Kalanchoe* sp. were evaluated for their ability to remove formaldehyde from sealed chambers seven days after being prepared and again five months later. The numbers and types of bacteria present in the soil were also determined at seven days and five months as shown in Table 5. It is interesting to observe that different plants grown in the same potting soil demonstrated significantly different formaldehyde removal rates.

Fresh potting soil without plants removed 46 µg/hr of formaldehyde from sealed chambers (Table 5). After seven days, *Spathiphyllum* sp. and *Kalanchoe* sp. planted in the same potting soil removed 300 µg/hr and 142 µg/hr, respectively. The potting soil which was maintained under the same conditions of watering and fertilizing as the *Spathiphyllum* sp. and *Kalanchoe* sp. removed 280 µg/hr of formaldehyde after five months, while the *Spathiphyllum* sp. removed 939 µg/hr after five months. It is interesting to observe that after five months the *Kalanchoe* sp. only removed 188 µg/hr which is less than the potting soil control. Although the soil in which the *Kalanchoe* sp. had grown contained more bacteria than the soil from the *Spathiphyllum* sp., they were a different type of bacteria. The predominant bacteria found in the *Spathiphyllum* sp. soil were gram-negative rods, whereas the predominant bacteria found in the *Kalanchoe* sp. soil and soil control were gram-positive rods. The ability of gram-negative bacteria, such as *Pseudomonas*, to degrade organic chemicals is well documented. (Davies and Evans, 1964; Evans et al., 1965; and LaPat-Polasko et al., 1984). Therefore, the ability of certain plants

to stimulate the production of gram-negative micro organisms such as *Pseudomonas* on and around the roots appears to be a major factor contributing to their increased chemical removal rates. Different plant roots appear to stimulate the production of microbes that may have negative or positive effects on removing chemicals from sealed chambers.

To determine the importance of plant leaf surfaces in removing formaldehyde and xylene from sealed chambers, plants with exposed potting soil and soil covered with sterilized sand were tested. Plant leaves contributed 33-49.5 percent of formaldehyde and xylene removal from the sealed chambers (Tables 6 and 7). The ability of plant leaves and stems to absorb, translocate and oxidize insecticides and other organic substances has been known and studied for many years. It is possible that the plant leaves are absorbing formaldehyde and xylene from the air and translocating them via the phloem/xylem to the plant roots where they are degraded by microorganisms.

The ability of *Rhapis excelsa* (lady palm) and *Nephrolepis exalta* (Boston fern) to not only continuously remove outgassing formaldehyde from sections of interior paneling, but improve with exposure time, suggests that the microorganisms are improving their ability to degrade formaldehyde through adaptation (Figures 1 and 2).

Conclusions.

(a) Low-light-requiring, interior plants have demonstrated the ability to remove significant quantities of formaldehyde, xylene, and ammonia from sealed chambers. (b) Experimental data indicates that both plant leaves and soil microorganisms are involved in removing these chemicals from sealed chambers. (c) Experimental data indicates that plants that culture large numbers of gram-negative bacteria on and around their roots are more effective in removing volatile organic chemicals from sealed chambers than plants that culture predominantly gram-positive bacteria. (d) Other factors, such as leaf structures which influence transpiration rates, may also be important factors in determining why some plants are more effective in removing organic chemicals from air than others.

Acknowledgements

The authors gratefully acknowledge the financial support of the Plants for Clean Air Council, 10210 Bald Hill Road, Mitchellville, Maryland 20721.

LITERATURE CITED

- Davies, J. R. and W. C. Evans. 1964. Oxidative metabolism of naphthalene by soil *Pseudomonads*. *Biochem. J.* 91:251-261.
- Evans, W. C., H. N. Fernley, and E. Griffiths. 1965. Oxidation metabolism of phenanthrene and anthracene by soil *Pseudomonads*. *Biochem. J.* 95:813-819.
- Katznelson, H. 1970. Nature and importance of the rhizosphere. In: K. W. Baker and W. C. Snyder [Eds.] *Ecology of soil-born plant pathogens*. Univ. of Cal. Press, pp. 187-209.
- LaPat-Polasko, L. T., P. L. McCarthy, and A. J. B. Zehnder. 1984. Secondary substrate utilization of methylene chloride by an isolated strain of *Pseudomonas* sp. *Appl. Environ. Microbiol.* 47(4):825-830.
- Nelson, M. J. R., S. O. Montgomery, E. J. O'Neill, and P. H. Pritchard. 1986. Aerobic metabolism of trichloroethylene by a bacterial isolate. *Appl. Environ. Microbiol.* 52(2):383-384.
- Rovira, A. D. 1959. Root excretions in relation to the rhizosphere effect. IV. Influence of plant species, age of plant, light, temperature and calcium nutrition on exudation. *Plant Soil* 11:53-64.
- Rovira, A. D. 1963. Microbial inoculation of plants. I. Establishment of free living nitrogen fixing bacteria in the rhizosphere and their effects on maize, tomato and wheat. *Plant Soil* 19:304-314.
- Rovira, A. D. 1970. Plant root exudates and their influence upon soil microorganisms. In: K. W. Baker and W. C. Snyder [Eds.] *Ecology of soil-born plant pathogens*. Univ. of Cal. Press, pp. 170-182.
- Rovira, A. D. and C. B. Davey. 1974. Biology of the rhizosphere. In: E. W. Carson [Ed.] *The plant root and its environment*. Univ. Press of Virginia, Charlottesville, VA, pp. 153-204.
- Saber, D. L. and R. L. Crawford. 1985. Isolation and characterization of *Flavobacterium* strains that degrade pentachlorophenol. *Appl. Environ. Microbiol.* 50(6):1512-1518.
- Tabak, H. H., S. A. Quave, C. I. Mashni, and F. F. Barth. 1981. Biodegradability studies with organic priority pollutant compounds. *J. Water Pollut. Control Fed.* 53(10):1503-1518.
- U. S. Environmental Protection Agency. 1989. Environmental Protection Agency report to Congress on indoor air quality. Executive summary and recommendations. EPA/400/1-89/001A. U. S. EPA, Washington, D.C.
- U. S. Environmental Protection Agency. 1988. Environmental Protection Agency report on indoor air quality in public buildings: Volume 1 and II. EPA/600/6-88ab, August, 1988.
- Wolverton, B. C. and D. D. Harrison. 1973. Aquatic plants for removal of mevinphos from the aquatic environment. *J. Miss. Academy Sciences*, 19:84-88.
- Wolverton, B. C., R. McDonald, and E. A. Watkins, Jr. 1984. Foliage plants for removing indoor air pollutants from energy-efficient homes. *Economic Botany*. 38(2):224-228.
- Wolverton, B. C., R. McDonald, and H. H. Mesick. 1985. Foliage plants for indoor removal of the primary combustion gases carbon monoxide and nitrogen dioxide. *J. Miss. Academy Sciences*. 30:1-8.

Table 1. Removal of **Formaldehyde** by Plants

Plants	μg Removed Per Hour	Temperature (°C)	Pot Size (cm)
<i>Nephrolepis exaltata</i> "Bostoniensis"	1863	26.1	20.3
<i>Chrysanthemum morifolium</i>	1450	26.0	15.2
<i>Phoenix roebelenii</i>	1385	22.8	35.6
<i>Dracaena deremensis</i> "Janet Craigs"	1361	24.7	25.4
<i>Nephrolepis obliterata</i>	1328	25.7	25.4
<i>Hedera helix</i>	1120	25.7	20.3
<i>Ficus benjamina</i>	940	25.7	15.2
<i>Spathiphyllum</i> sp. "Clevelandii"	939	26.0	15.2
<i>Dracaena fragrans</i>	938	26.2	20.3
<i>Rhapis excelsa</i>	876	23.9	25.4
<i>Dracaena marginata</i>	772	25.6	20.3
<i>Dracaena deremensis</i> "Warneckeii"	760	25.7	25.4
<i>Liriope spicata</i>	758	23.4	15.2
<i>Dendrobium</i> sp.	756	24.4	20.3
<i>Dieffenbachia</i> sp. "Exotica compacta"	754	24.7	15.2
Tulip "Yellow Present"	717	24.5	15.2
<i>Ficus sabre</i>	692	22.9	25.4
<i>Homalomena</i> sp.	668	26.1	20.3
<i>Chamaedorea elegans</i>	660	22.4	15.2
<i>Rhododendron indicum</i>	617	23.6	15.2
<i>Aglaonema</i> sp. "Silver Queen"	564	25.5	15.2
<i>Chlorophytum comosum</i> "Vittatum"	560	25.9	20.3
<i>Dieffenbachia camille</i>	469	24.3	15.2
<i>Cissus rhombifolia</i>	376	25.8	15.2
<i>Syngonium podophyllum</i>	341	22.2	20.3
<i>Anthurium andraeanum</i>	336	26.6	25.4
<i>Calathea ornata</i>	334	26.4	20.3
<i>Euphorbia pulcherrima</i>	309	23.6	15.2
<i>Cyclamen persicum</i>	295	21.9	15.2
<i>Phalaenopsis</i> sp.	240	26.8	15.2
<i>Aechmea fasciata</i>	234	24.9	15.2
<i>Sansevieria trifasciata</i>	189	26.2	15.2
<i>Aloe barbadensis</i>	188	26.2	15.2

Table 2. Removal of **Xylene** by Plants

Plants	μg Removed Per Hour	Temperature (°C)	Pot Size (cm)
<i>Phoenix roebelenii</i>	610	23.1	35.6
<i>Dieffenbachia camille</i>	341	26.6	15.2
<i>Dracaena marginata</i>	338	26.9	20.3
<i>Dieffenbachia maculata</i>	325	26.4	25.4
<i>Homalomena</i> sp.	325	26.0	20.3
<i>Nephrolepis obliterata</i>	323	26.0	25.4
<i>Dracaena deremensis</i> "Warneckeii"	295	25.6	25.4
<i>Anthurium andraeanum</i>	276	25.8	25.4
<i>Dracaena fragrans</i>	274	23.9	20.3
<i>Ficus benjamina</i>	271	25.5	15.2
<i>Spathiphyllum</i> "Clevelandii"	268	25.6	15.2
<i>Chlorophytum comosum</i>	247	27.2	20.3
<i>Liriope spicata</i>	230	22.9	15.2
Tulip "Yellow Present"	229	24.5	15.2
<i>Chamaedorea elegans</i>	223	22.6	16.5
<i>Syngonium podophyllum</i>	220	22.0	15.2
<i>Rhapis excelsa</i>	217	25.5	25.4
<i>Nephrolepis exaltata</i> "Bostoniensis"	208	26.8	20.3
<i>Chrysanthemum morifolium</i>	201	24.0	15.2
<i>Dendrobium</i> sp.	200	26.9	8.0
<i>Cyclamen persicum</i>	173	22.0	15.2
<i>Kalanchoe</i>	170	25.9	15.2
<i>Rhododendron indicum</i>	168	22.2	15.2
<i>Sansevieria trifasciata</i>	157	26.6	15.2
<i>Dracaena deremensis</i> "Janet Craigs"	154	25.6	25.4
<i>Guzmania</i> cherry	146	24.5	12.7
<i>Hedera helix</i>	131	25.6	20.3
<i>Euphorbia pulcherrima</i>	116	21.2	15.2
<i>Senecio cruentus</i>	115	23.0	15.2
<i>Neoregelia</i> cv.	47	26.5	12.7

Table 3. Removal of **Ammonia** by Plants

Plants	μg Removed Per Hour	Temperature (°C)	Pot Size (cm)
<i>Rhapis excelsa</i>	7,356	24.1	25.4
<i>Homalomena</i> sp.	5,208	24.3	20.3
<i>Liriope spicata</i>	4,308	26.4	15.2
<i>Anthurium andraeanum</i>	4,119	24.5	25.4
<i>Chrysanthemum morifolium</i>	3,641	26.5	15.2
<i>Calathea vittata</i>	3,100	26.2	20.3
Tulip "Yellow Present"	2,815	26.7	15.2
<i>Chamaedorea elegans</i>	2,453	25.8	16.5
<i>Ficus benjamina</i>	1,480	24.4	15.2
<i>Spathiphyllum</i> "Clevelandii"	1,269	24.1	15.2
<i>Rhododendron indicum</i>	984	23.3	15.2

*Greenhouse grown forced tulip

Table 4. Formaldehyde Removal Rates Of Potting Soils

Soils	μg Removed Per Hour	Ave. Temp. (°C)	Soil Bacterial Counts (cfu/g)
Sterilized Soil	< 0.05	25.8	0
Unsterilized Soil	188.0	25.0	235
Unsterilized Soil Covered With Sterilized Sand	<0.05	26.8	0*

* Sand only

Table 5. Formaldehyde Removal Rates of Fresh Potting Soil and Plants Grown In Potting Soil

Soil And Plants	μg Removed Per Hour After 7 Days	μg Removed Per Hour After 5 Mos.	Average Temp (°C)	Soil Bacterial Counts (cfu/g) After 7 Days	Soil Bacterial Counts (cfu/g) After 5 Mos.
Fresh Potting Soil	46	280	26.0	9.00×10^2	4.13×10^2
<i>Ficus benjamina</i> in Potting Soil	335	940	25.7	5.22×10^3	5.00×10^3
<i>Spathiphyllum</i> "Clevelandii" in Potting Soil	300	939	26.0	4.37×10^3	5.50×10^3
<i>Sansevieria</i> <i>trifasciata</i> in Potting Soil	97	189	26.6	7.56×10^2	5.70×10^2
<i>Kalanchoe</i> in Potting Soil	142	188	25.6	7.38×10^3	7.50×10^3

Continued on page 15

Table 6. Removal of Formaldehyde by Plants with Exposed Soil and Soil Covered with Sterilized Sand

Plants And Soil	µg Removed Per Hour	% Removed by soil Microbes	% Removed by Leaves	Average Temp (°C)
<i>Aglaonema</i> "Silver Queen" Exposed Soil	564			
		67	33	25.3
<i>Aglaonema</i> "Silver Queen" Soil Covered With Sterilized Sand	188			
<i>Dieffenbachia</i> "Exotica Compacta" Exposed Soil	754			
		63	37	25.4
<i>Dieffenbachia</i> "Exotica Compacta" Soil Covered With Sterilized Sand	281			
<i>Nephrolepis exaltata</i> "Bostoniensis" Exposed Soil	1027			
		60	40	25.3
<i>Nephrolepis exaltata</i> "Bostoniensis" Soil Covered With Sterilized Sand	409			

Table 7. Removal of Xylene by Plants with Exposed Soil and Soil Covered with Sterilized Sand

Plants And Soil	µg Removed Per Hour	% Removed by soil Microbes	% Removed by Leaves	Average Temp (°C)
<i>Dieffenbachia maculata</i> Exposed Soil	325			
		53.0	47.0	26.3
<i>Dieffenbachia maculata</i> Soil Covered With Sterilized Sand	154			
<i>Nephrolepis exaltata</i> "Bostoniensis" Exposed Soil	208			
		50.5	49.5	26.5
<i>Nephrolepis exaltata</i> "Bostoniensis" Soil Covered With Sterilized Sand	103			