Proof of the Clustered Hadwiger Conjecture

David Wood Monash University Melbourne, Australia

joint work with

Vida Dujmović



Louis Esperet




Pat Morin




arXiv:2306.06224 and FOCS 2023

4-colour theorem: every planar graph is 4-colourable



4-colour theorem: every planar graph is 4-colourable



a graph H is a minor of a graph G if H can be obtained from a subgraph of G by contracting edges




a graph H is a minor of a graph G if H can be obtained from a subgraph of G by contracting edges



complete K_h minor \equiv


h pairwise-disjoint pairwise-adjacent connected subgraphs



a graph H is a minor of a graph G if H can be obtained from a subgraph of G by contracting edges



complete K_h minor \equiv h pairwise-disjoint pairwise-adjacent connected subgraphs



planar graphs are 4-colourable and are K_5 -minor-free

excluded minor	graph family	colourable	
K_1	no vertices	0	
K_2	no edges	1	
K_3	forests	2	Same.
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
			-

excluded minor	graph family	colourable	
K_1	no vertices	0	
K_2	no edges	1	
K_3	forests	2	See
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
K_5	planar++	4	[4CT & Wagner '37]

excluded minor	graph family	coloura	ble
K_1	no vertices	0	
K_2	no edges	1	
K_3	forests	2	
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
K_5	planar++	4	[4CT & Wagner '37]
K_6	???	5 [4	4CT & Rob., Seymour & Thomas '93]

	•		
K_1	no vertices	0	
K_2	no edges	1	
K_3	forests	2	
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
K_5	planar++	4	[4CT & Wagner '37]
K_6	???	5	[4CT & Rob., Seymour & Thomas '93]

colourable

Hadwiger's Conjecture [1943] every K_h -minor-free graph is (h-1)-colourable

excluded

minor

graph family



κ_1	no vertices	U	
K_2	no edges	1	
K_3	forests	2	
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
K_5	planar++	4	[4CT & Wagner '37]
K_6	???	5	[4CT & Rob., Seymour & Thomas '93]

colourable

Hadwiger's Conjecture [1943]

graph

family

excluded

minor

every K_h -minor-free graph is (h-1)-colourable

• $O(h(\log h)^{1/2})$ colours [Kostochka '84, Thomason '84]

	,		
K_1	no vertices	0	
K_2	no edges	1	
K_3	forests	2	
K_4	series parallel	3	[Hadwiger '43, Dirac '52]
K_5	planar++	4	[4CT & Wagner '37]
K_6	???	5	[4CT & Rob., Seymour & Thomas '93]

colourable

Hadwiger's Conjecture [1943]

every K_h -minor-free graph is (h-1)-colourable

graph family

excluded

minor

- $O(h(\log h)^{1/2})$ colours
- $O(h(\log h)^{1/4+\epsilon})$ colours

olours [Kostochka '84, Thomason '84]

[Kostochka '84, Thomason '84] [Norin–Song–Postle '19]

K_2	no edges	1	The state of the s	
K_3	forests	2	Same of the same o	
K_4	series parallel	3	[Hadwiger '43, Dirac '52]	
K_5	planar++	4	[4CT & Wagner '37]	
K_6	???	5	[4CT & Rob., Seymour & Thomas '93]	
Hadwiger's Conjecture [1943]				
every k	K _h -minor-free grap	oh is	(h-1)-colourable	

[Kostochka '84, Thomason '84]

[Norin-Song-Postle '19]

[Delcourt-Postle '21]

colourable

0

excluded

minor

 K_1

graph

family no vertices


• $O(h(\log h)^{1/2})$ colours

• $O(h \log \log h)$ colours

• $O(h(\log h)^{1/4+\epsilon})$ colours

k-colouring with clustering *c*:

- each v is assigned one of k colours
- ullet each monochromatic component has $\leqslant c$ vertices



k-colouring with clustering *c*:

- each v is assigned one of k colours
- each monochromatic component has $\leq c$ vertices

planar graphs are 4-colourable with clustering 1 [4CT]

k-colouring with clustering *c*:

- each v is assigned one of k colours
- each monochromatic component has $\leq c$ vertices


planar graphs are 4-colourable with clustering 1 [4CT] or clustering 2 [Cowen, Cowen, Woodall '86]

k-colouring with clustering *c*:

- each v is assigned one of k colours
- each monochromatic component has $\leq c$ vertices

planar graphs are 4-colourable with clustering 1 [4CT] or clustering 2 [Cowen, Cowen, Woodall '86]

but planar graphs are not 3-colourable with bounded clustering



k-colouring with clustering *c*:

- \bullet each v is assigned one of k colours
- each monochromatic component has ≤ c vertices

clustered chromatic number

 $\chi^{\mathsf{CLUS}}(\mathcal{G}) := \mathsf{minimum} \ k \ \mathsf{such that} \ \exists c \ \mathsf{and}$ every graph in \mathcal{G} is k-colourable with clustering c

k-colouring with clustering *c*:

- each v is assigned one of k colours
- ullet each monochromatic component has $\leqslant c$ vertices

clustered chromatic number

 $\chi^{\mathsf{CLUS}}(\mathcal{G}) := \mathsf{minimum}\ k \ \mathsf{such}\ \mathsf{that}\ \exists c \ \mathsf{and}$ every graph in \mathcal{G} is k-colourable with clustering c

$$\chi^{\mathsf{CLUS}}(\mathsf{planar\ graphs}) = 4$$

k-colouring with clustering *c*:

- each v is assigned one of k colours
- ullet each monochromatic component has $\leqslant c$ vertices

clustered chromatic number

 $\chi^{\mathsf{CLUS}}(\mathcal{G}) := \mathsf{minimum} \ k \ \mathsf{such} \ \mathsf{that} \ \exists c \ \mathsf{and}$ every graph in \mathcal{G} is k-colourable with clustering c


$$\chi^{\mathsf{CLUS}}(\mathsf{planar\ graphs}) = 4$$

 $\chi^{\rm CLUS}({\rm graphs\ embeddable\ on\ any\ fixed\ surface})=4$ [Dvořák & Norin '17]




$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs}) \geqslant h-1$$

$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs})\geqslant h-1$$



$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs})\geqslant h-1$$



 $S_{h,d}$ is K_h -minor-free and has no (h-2)-colouring with clustering $\leqslant d$

lower bound

$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs}) \geqslant h-1$$

upper bounds

$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs})\geqslant h-1$$

upper bounds	# colours	clustering
[Kawarabayashi & Mohar '07]	$\lceil \frac{31}{2}h \rceil$	c(h)
[W. '10]*	$\lceil \frac{7h-3}{2} \rceil$	<i>c</i> (<i>h</i>)
[Edwards, Kang, Kim, Oum, Seymour '14]	4 <i>h</i> – 4	<i>c</i> (<i>h</i>)
[Liu & Oum '15]	3h - 3	<i>c</i> (<i>h</i>)
[Norin '15]	2h - 2	<i>c</i> (<i>h</i>)
[van den Heuvel & W. '17]	2h - 2	$\lceil \frac{h-2}{2} \rceil$
[Liu & W. '19]	h	<i>c</i> (<i>h</i>)

lower bound

$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs})\geqslant h-1$$

upper bounds

clustered Hadwiger theorem [Dujmović, Esperet, Morin, W. '23]

- K_h -minor-free graphs are (h-1)-colourable with clustering c(h)
- $\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs}) = h 1$

lower bound

$$\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs})\geqslant h-1$$

upper bounds

clustered Hadwiger theorem [Dujmović, Esperet, Morin, W. '23]

- K_h -minor-free graphs are (h-1)-colourable with clustering c(h)
- $\chi^{\mathsf{CLUS}}(K_h\text{-minor-free graphs}) = h-1$

(announced by Dvořák & Norin '17)

 $K_{s,t}$ -minor-free graphs $(s\leqslant t)$

Hadwiger's Conjecture $K_{s,t}$ -minor-free graphs are (s+t-1)-colourable

 $K_{s,t}$ -minor-free graphs $(s\leqslant t)$

Hadwiger's Conjecture $K_{s,t}$ -minor-free graphs are (s+t-1)-colourable

 $| \text{lower bound} \quad \chi^{\text{CLUS}}(\textit{K}_{s,t}\text{-minor-free graphs}) \geqslant s+1$

$$K_{s,t}$$
-minor-free graphs $(s\leqslant t)$

Hadwiger's Conjecture $K_{s,t}$ -minor-free graphs are (s+t-1)-colourable

lower bound $\chi^{\mathsf{CLUS}}(K_{s,t}\text{-minor-free graphs})\geqslant s+1$ upper bounds


oounds	# colours	clustering
[van den Heuvel & W. '17]	3 <i>s</i>	$O(t^s)$
[Dvořák & Norin '17]	2s + 2	c(s,t)
[Liu & W. '19]	s+2	c(s,t)

$$K_{s,t}$$
-minor-free graphs $(s\leqslant t)$


Hadwiger's Conjecture $K_{s,t}$ -minor-free graphs are (s+t-1)-colourable

upper bo	ounds 	# colours	clustering
	[van den Heuvel & W. '17]	3 <i>s</i>	$O(t^s)$
	[Dvořák & Norin '17]	2s + 2	c(s,t)
	[Liu & W. '19]	s+2	c(s,t)


- $oldsymbol{\epsilon}_{s,t}$ -minor-free graphs are (s+1)-colourable with clustering c(s,t)
- $\chi^{\mathsf{CLUS}}(K_{s,t}\text{-minor-free graphs}) = s+1 \text{ for } t \geqslant \max\{s,3\}$



let $\mathcal{J}_{s,t}$ be the class of all graphs with




let $\mathcal{J}_{s,t}$ be the class of all graphs with



main theorem [Dujmović, Esperet, Morin, W. '22] $\mathcal{J}_{s,t}$ -minor-free graphs are (s+1)-colourable with clustering c(s,t)


let $\mathcal{J}_{s,t}$ be the class of all graphs with



main theorem [Dujmović, Esperet, Morin, W. '22] $\mathcal{J}_{s,t}$ -minor-free graphs are (s+1)-colourable with clustering c(s,t)

implies previous theorems since $\mathcal{J}_{h-2,2} = \{K_h\}$

let $\mathcal{J}_{s,t}$ be the class of all graphs with




main theorem [Dujmović, Esperet, Morin, W. '22] $\mathcal{J}_{s,t}$ -minor-free graphs are (s+1)-colourable with clustering c(s,t)

implies previous theorems since $\mathcal{J}_{h-2,2} = \{K_h\}$ and every graph in $\mathcal{J}_{s,t}$ contains $K_{s,t}$



1. treewidth


measures how similar a graph is to a tree



- 1. treewidth
- 2. graph product structure theory

- 1. treewidth
- 2. graph product structure theory

theorem [Dujmović, Joret, Micek, Morin, Ueckerdt, W. '19] every planar graph G is a subgraph of $H \boxtimes P$ for some graph H with treewidth ≤ 8 and some path P



- 1. treewidth
- 2. graph product structure theory
- 3. graph minor structure theorem

theorem [Robertson & Seymour '80s] graphs excluding a fixed minor can be constructed from graphs embedded on surfaces, vortices, apex vertices, and clique-sums

- 1. treewidth
- 2. graph product structure theory
- 3. graph minor structure theorem
- 4. islands, curtains, drapes, etc.

K_h -minor-free	K_h -subdivision-free
Hadwiger Conjecture: $h-1$ colours suffice	Hajós' Conjecture: $h-1$ colours suffice

K_h -minor-free	K_h -subdivision-free
Hadwiger Conjecture: $h-1$ colours suffice	Hajós' Conjecture: $\mathit{h}-1$ colours suffice
true for clustered colouring [Dujmović, Esperet, Morin, W. '23]	almost true for clustered colouring $4h - 9$ colours [Liu W. '19]

K_h -minor-free	K_h -subdivision-free
Hadwiger Conjecture: $h-1$ colours suffice	Hajós' Conjecture: $h-1$ colours suffice
true for clustered colouring [Dujmović, Esperet, Morin, W. '23]	almost true for clustered colouring $4h - 9$ colours [Liu W. '19]
Hadwiger's Conjecture is true for almost every graph [Bollobas, Catlin, Erdős '80]	Hajós' Conjecture is false for every graph! [Erdős, Fajtlowicz '81]

K_h -minor-free	K_h -subdivision-free
Hadwiger Conjecture: $h-1$ colours suffice	Hajós' Conjecture: $\mathit{h}-1$ colours suffice
true for clustered colouring [Dujmović, Esperet, Morin, W. '23]	almost true for clustered colouring $4h - 9$ colours [Liu W. '19]
Hadwiger's Conjecture is true for almost every graph [Bollobas, Catlin, Erdős '80]	Hajós' Conjecture is false for every graph! [Erdős, Fajtlowicz '81]

so do not think of the Clustered Hadwiger Theorem as evidence for Hadwiger's Conjecture