


45TH AUSTRALASIAN COMBINATORICS CONFERENCE



The University of Western Australia, December 11–15, 2023

© 2023, the organisers:

John Bamberg

Alice Devillers

Michael Giudici

Luke Morgan

Cheryl Praeger

Gordon Royle

45acc.github.io

Welcome!

This is the fifth time the ACC (formerly, ACCMCC) has been hosted in Perth, having previously been at UWA and/or Curtin University in the years 1984, 1992, 2001 and 2013. There are more than 65 registrants for this year, making it the second largest ACC/ACCMCC to be hosted in Western Australia. We are very grateful for the support from the following institutions and organisations:

- The School of Physics, Mathematics, and Computing (UWA)
- Optiver
- The Institute of Combinatorics and its Applications

We wish you an interesting and exciting conference, and a pleasant stay in Perth.

The organisers:
John Bamberg
Alice Devillers
Michael Giudici
Luke Morgan
Cheryl Praeger
Gordon Royle

Contents

Welcome!	iii
1 Invited talks	3
2 Contributed talks	13
3 List of participants	73

Sunday

	EZone
17:00 – 19:00	Welcome reception and registration

Monday

	Weatherburn LT	Blakers LT	Praeger LR
8.00 – 8.45	Registration		
8.45 – 9.00	Opening address (Prof Mark Reynolds)		
9.00 – 10:00	<i>Gabriel Verret</i> 12		
10.00 – 10.30	Morning tea		
10.30 – 11.00	Chen* 22	Bastida* 18	Santos 59
11.00 – 11.30	Ding* 26	Tangjai 64	Wang* 66
11.30 – 12.00	Mitrović* 56	Farooq* 29	Yost 69
12.00 – 12.30	Dacaymat* 24	Semple 61	Umar 65
12.30 – 14.30	Lunch break		
14.30 – 15.30	<i>CMSA Prize Winner</i>		
15.30 – 16.00	Afternoon tea		
16.00 – 16.30	Basit 17	Bunjamin* 21	
16.30 – 17.00	Liebenau 51	Mitchell* 55	
17.00 – 17.30	Hasunuma 36	Lacaze-Masmonteil* 46	

Tuesday

	Weatherburn LT	Blakers LT	Praeger LR
9.00 – 10:00	<i>Krystal Guo</i> 7		
10.00 – 10.30	Morning tea		
10.30 – 11.00	Hickingbotham* 38	Briones 20	
11.00 – 11.30	Distel* 27	Mammoliti 52	
11.30 – 12.00	Brettell 19	Ernst* 28	
12.00 – 12.30	Wood 68	Klawuhn* 45	
12.30 – 14.30	Lunch break		
14.30 – 15.30	<i>Gary Greaves</i> 6		
15.30 – 16.00	Afternoon tea		
16.00 – 16.30	Allsop* 15	Imamura 40	
16.30 – 17.00	Ghafari* 32	Kawabuchi 44	
17.00 – 17.30	CMSA AGM		

Wednesday

	Weatherburn LT	Blakers LT	Praeger LR
9.00 – 10.00	<i>André Kündgen</i> 8		
10.00 – 10.30	Morning tea		
10.30 – 11.00	Gentle* 31	Lehner 48	Gunasekara 34 Smith* 62 Li* 49
11.00 – 11.30	Syrotiuk 63	Hafidh* 35	
11.30 – 12.00	Hirao 39	Maruta 54	
12.00 – 12.30	Hawtin 37	Yasufuku 70	
12.30 – 13.30	Lunch break		
14.00 – 17.00	<i>Excursion</i>		

Thursday

	Weatherburn LT	Blakers LT	Praeger LR
9.00 – 10:00	<i>Tibor Szabó</i> 10		
10.00 – 10.30	Morning tea		
10.30 – 11.00	Arumugam* 16	Satake 60	
11.00 – 11.30	Miura 57	Zhang* 71	
11.30 – 12.00	Lia 50		
12.00 – 12.30	De Beule 25		
12.30 – 14.30	Lunch break		
14.30 – 15.30	<i>Geertrui Van de Voorde</i> 11		
15.30 – 16.00	Afternoon tea		
16.00 – 16.30	McKay 53		
16.30 – 17.00	Colbourn 23		
17.00 – 17.30	Wanless 67		

18.30: Conference dinner (UniClub)

Friday

	Weatherburn LT	Blakers LT
9.00 – 10.00	<i>Sara Davies</i> 5	
10.00 – 10.30	Morning tea	
10.30 – 11.00	Kaemawichanurat 42	Popiel 58
11.00 – 11.30	Greenhill 33	Freedman 30
11.30 – 12.00	Isaev 41	Lansdown 47
12.00 – 14.00	Lunch break	
14.30 – 15.30	<i>Padraig Ó Catháin</i> 9	
15.30 – 16.00	Afternoon tea	

1

Invited talks

<i>Sara Davies</i> – The Hamilton decomposition problem	5
<i>Gary Greaves</i> – How to design a graph with three eigenvalues	6
<i>Krystal Guo</i> – Algebraic graph theory and quantum walks	7
<i>Andre Kundgen</i> – The Saturation Spectrum of odd cycles	8
<i>Padraig Ó Catháin</i> – Quadratic forms in design theory	9
<i>Tibor Szabó</i> – New Ramsey multiplicity bounds and search heuristics	10
<i>Geertrui Van de Voorde</i> – ‘Segre-type’ theorems: combinatorial characterisations for algebraic objects	11
<i>Gabriel Verret</i> – Local actions and eigenspaces of vertex-transitive graphs	12

The Hamilton decomposition problem

Sara Davies

The University of Queensland

Determining whether an arbitrary graph has a Hamilton cycle is a classic problem in graph theory. A *Hamilton decomposition* of a graph is a set of edge-disjoint Hamilton cycles that collectively contain all of the edges of the graph. The study of Hamilton decompositions dates back to the late 1800's and has received a lot of attention since the 1980's. In this talk, I will survey some of the progress made on this problem, especially on Hamilton decompositions of Cayley graphs, infinite graphs, line graphs and graph products.

Gary Greaves

(Joint work with Jose Yip)

Muzychuk and Klin initiated the study of a graph with three distinct eigenvalues via its Weisfeiler-Leman closure (also known as the coherent closure). They classified such graphs whose Weisfeiler-Leman closure has rank at most 7. In this talk, I will provide a brief overview of the history of non-regular graphs with three distinct eigenvalues, as well as present our recent results on such graphs whose Weisfeiler-Leman closure has a small rank. Our results include the discovery of a new non-regular graph with three distinct eigenvalues obtained from a quasi-symmetric design and a new conjecturally infinite family of non-regular graphs having three distinct eigenvalues obtained by switching Latin square graphs.

Algebraic graph theory and quantum walks

Krystal Guo

Korteweg-De Vries Institute for Mathematics, University of Amsterdam and QuSoft

The interplay between the properties of graphs and the eigenvalues of their adjacency matrices is well-studied. Important graph invariants, such as diameter and chromatic number, can be understood using these eigenvalue techniques. In this talk, we bring these classical techniques in algebraic graph theory to the study of quantum walks.

A system of interacting quantum qubits can be modelled by a quantum process on an underlying graph and is, in some sense, a quantum analogue of random walk. This gives rise to a rich connection between graph theory, linear algebra and quantum computing. In this talk, I will give an overview of applications of algebraic graph theory in quantum walks, as well as various recent results on discrete-time quantum walks and strong cospectrality of vertices.

The Saturation Spectrum of odd cycles

Andre Kundgen

California State University San Marcos

(Joint work with Ronald J. Gould and Minjung Kang)

Given a graph H , we say that a graph G is H -saturated if H is not a subgraph of G , but the addition of any new edge to G creates at least one copy of H . In this talk we will discuss all pairs (n, m) for which there is a C_5 -saturated graph on n vertices and m edges. In addition, we determine all but $O(nk)$ possible sizes for n -vertex H -saturated graphs when H is an odd cycle C_{2k+1} for $k \geq 3$.

Quadratic forms in design theory

Padraig Ó Catháin

Dublin City University

(Joint work with Guillermo Nuñez Ponasso, Oliver Gnille and Oktay Olmez.)

The classification of quadratic forms over the rational numbers, due to Minkowski, Hilbert and Hasse among others, is a major achievement of mathematicians in the early twentieth century. In concrete terms, given square rational matrices A and B it yields necessary and sufficient conditions for the existence of an invertible matrix X such that $X^{\top}AX = B$. (In contrast, the Jordan Canonical Form gives necessary and sufficient conditions for solvability of $X^{-1}AX = B$ over an algebraically closed field, and the Frobenius Canonical Form solves the conjugacy problem over an arbitrary field.) The main tools in the classification of quadratic forms are Legendre and Hilbert symbols, which describe existence of solutions to certain quadratic equations.

Groundbreaking work of Bruck, Ryser and Chowla in the mid-twentieth century applied this theory to obtain non-existence of certain combinatorial designs. While in theory the application is straightforward, Marshall Hall described the computations as detailed and troublesome. This seems to have scared a substantial number of combinatorialists. In this talk, we aim to restore the reputation of the Bruck-Ryser-Chowla theorem by demonstrating that the algebraic manipulations are less familiar, but not more difficult, than Gaussian elimination.

I will motivate this talk by an application to a problem on symmetric designs which Darryn Bryant posed to me in 2013, while I was a postdoc at the University of Queensland.

'Segre-type' theorems: combinatorial characterisations for algebraic objects

Geertrui Van de Voorde

The University of Canterbury

One of the most beautiful results within finite geometry is Segre's characterisation of conics in Desarguesian projective planes of odd order. In 1955, Segre showed that in those planes, the coordinates of a point set that has the same *combinatorial* properties as a conic, must have the same *algebraic* property of satisfying a quadratic equation. In even order planes, the situation is vastly different, and the classification of ovals remains is still an open problem.

Several ‘Segre-type’ questions have been studied for objects such as *quadrics*, *Hermitian varieties*, and more generally, for sets with *few intersection numbers*.

In this talk, I'll give an overview of some of the history of this subject and present new recent results.

2

Contributed talks

<i>Jack Allsop*</i> – Latin squares without proper subsquares	15
<i>Vishnuram Arumugam*</i> – Groups of Lie Type Acting on Generalised Quadrangles	16
<i>Abdul Basit</i> – Point-box incidences and logarithmic density of semilinear graphs	17
<i>Sam Bastida*</i> – List Colouring Graphs with bounded Maximal Local Edge Connectivity	18
<i>Nick Brettell</i> – A comparison of graph width parameters	19
<i>Dom Vito A. Briones</i> – Association schemes on triples from two-transitive groups	20
<i>Yudhistira Andersen Bunjamin*</i> – Group divisible designs with block size three and two group sizes	21
<i>Lei Chen*</i> – The distinguishing number of 2-arc-transitive bipartite graphs	22
<i>Charles Colbourn</i> – Covering Arrays via Finite Fields	23
<i>John Mel Dacaymat*</i> – Diameter of some families of quotient-complete arc-transitive graphs	24
<i>Jan De Beule</i> – A strongly regular graph co-spectral and non-isomorphic to $\text{NO}^+(8, 2)$	25
<i>Zhaochen Ding*</i> – Compatible groups and inverse limits	26
<i>Marc Distel*</i> – Proper Minor-Closed Classes of Graphs have Assouad-Nagata Dimension 2	27
<i>Alena Ernst*</i> – Erdős-Ko-Rado theorems for finite general linear groups	28
<i>Muhammad Talha Farooq*</i> – Enumeration of the Hosoya Index in Anthracene System	29
<i>Saul Freedman</i> – Spreading primitive groups of diagonal type do not exist	30
<i>Dani Gentle*</i> – Levenshtein’s conjecture for sequence covering arrays	31
<i>Afsane Ghafari Baghestani*</i> – Existence of Latin Squares with Constrained Transversals	32
<i>Catherine Greenhill</i> – Enumerating dihypergraphs	33
<i>Ajani De Vas Gunasekara</i> – Transitive path decompositions of Cartesian products of complete graphs	34
<i>Yusuf Hafidh*</i> – Perfect codes in Cayley graphs on $\mathbb{Z}_p \times \mathbb{Z}_p$ and \mathbb{Z}_{p^k}	35
<i>Toru Hasunuma</i> – Connectivity Preserving Hamiltonian Cycles in k -Connected Dirac Graphs	36
<i>Dan Hawtin</i> – Large sets of infinite-dimensional q -Steiner systems	37
<i>Robert Hickingbotham*</i> – Powers of planar graphs, product structure, and blocking partitions	38
<i>Masatake Hirao</i> – Spherical designs and the D_4 lattice	39
<i>Koji Imamura</i> – Matroid representation over finite rings	40
<i>Mikhail Isaev</i> – Cumulant expansion for counting Eulerian orientations	41
<i>Pawaton Kaemawichanurat</i> – Safe Sets and Dominating Sets of Graphs	42

<i>Nina Kamčev</i> – Common and Sidorenko linear patterns	43
<i>Shinya Kawabuchi</i> – Some Properties of q -Perfect Matroid Designs	44
<i>Lukas Klawuhn*</i> – Designs in the generalised symmetric group	45
<i>Alice Lacaze-Masmonteil*</i> – On the directed Oberwolfach problem with two tables	46
<i>Jesse Lansdown</i> – Constructing witnesses for nonspreading permutation groups	47
<i>Florian Lehner</i> – Self-avoiding walks on graphs with infinitely many ends	48
<i>Yuxuan Li*</i> – The second largest eigenvalue of non-normal Cayley graphs on symmetric groups generated by cycles	49
<i>Stefano Lia</i> – Tensor representation of semifields and commuting polarities	50
<i>Anita Liebenau</i> – Universality for graphs of bounded degeneracy	51
<i>Adam Mammoliti</i> – On generalisations of The Erdős-Ko-Rado Theorem for per- mutations	52
<i>Brendan McKay</i> – Some new results in combinatorial generation	53
<i>Tatsuya Maruta</i> – On the non-existence of q -ary linear codes with minimum weight $d \equiv -1 \pmod{q}$	54
<i>Jeremy Mitchell*</i> – Equally Distributed 1-Factorisations of Graphs	55
<i>Dorđe Mitrović*</i> – Automorphisms of direct products of circulant graphs	56
<i>Yusuke Miura</i> – On the minimal 2-blocking sets in $PG(5, 2)$	57
<i>Tomasz Popiel</i> – Computing with the Monster group (a public service announce- ment)	58
<i>Rovin B. Santos</i> – Prime labeling of some graphs with Eisenstein integers	59
<i>Shohei Satake</i> – Explicit $K_{3,3}$ -subdivisions of Markoff mod p graphs	60
<i>Charles Semple</i> – Optimising phylogenetic diversity on phylogenetic networks	61
<i>Jacob Smith*</i> – New 2-closed groups that are not automorphism groups of di- graphs	62
<i>Violet Syrotiuk</i> – The Screening Effectiveness of Locating Arrays	63
<i>Wipawee Tangjai</i> – On a coloring of a δ -complement graph	64
<i>Abdullahi Umar</i> – Combinatorial results for certain semigroups of contraction mappings of a finite chain	65
<i>Jie Wang*</i> – Some lower bounds on conditionally decomposable polytopes	66
<i>Ian Wanless</i> – Automorphisms of quadratic quasigroups	67
<i>David Wood</i> – Proof of the Clustered Hadwiger Conjecture	68
<i>David Yost</i> – Polytopes with minimal number of edges	69
<i>Keita Yasufuku</i> – On the non-existence of Griesmer linear codes	70
<i>Chuanqi Zhang*</i> – On linear-algebraic notions of expansion	71

Latin squares without proper subsquares

*Jack Allsop**

Monash University

(Joint work with Ian Wanless)

A Latin square of order n is an $n \times n$ matrix of n symbols, such that each symbol occurs exactly once in each row and column. A subsquare of order k is a $k \times k$ submatrix of a Latin square that is itself a Latin square. Every Latin square of order n contains n^2 subsquares of order one, and one subsquare of order n . All other subsquares are called proper. If a Latin square contains no proper subsquares then it is called N_∞ . Around 50 years ago Hilton conjectured that an N_∞ Latin square of order n exists for all sufficiently large n . Hilton's conjecture was previously known to hold for all integers n not of the form $2^a 3^b$ for integers $a \geq 1$ and $b \geq 0$. We resolve Hilton's conjecture by constructing N_∞ Latin squares for all previously unresolved orders.

Groups of Lie Type Acting on Generalised Quadrangles

*Vishnuram Arumugam**

The University of Western Australia

Incidence geometry is the study of geometric structures involving a collection of points and lines along with a relation (called incidence) which tells us whether a point lies on a line. A generalised polygon is a type of point-line incidence structure that was introduced by Jacques Tits in 1959 to study the groups of Lie type as the symmetries of geometric objects. Since then, these objects have been studied extensively in the areas of group theory and finite geometry. The classification of these objects started from Weiss and Tits and many results about the existence (and non-existence) of generalised polygons under various symmetry conditions (point primitivity, flag transitivity and so on) since then. I will provide a survey of the work that has been done and some recent progress in this classification.

Point-box incidences and logarithmic density of semilinear graphs

Abdul Basit

Monash University

(Joint work with Artëm Chernikov, Sergei Starchenko, Terence Tao, and Chieu-Minh Tran)

Zarankiewicz’s problem in extremal graph theory asks for the maximum number of edges in a bipartite graph on n vertices which does not contain a copy of $K_{k,k}$, the complete bipartite graph with k vertices in both classes. We will consider this question for incidence graphs of geometric objects. Significantly better bounds are known in this setting, in particular when the geometric objects are defined by systems of algebraic inequalities. We show even stronger bounds under the additional constraint that the defining inequalities are linear. We will also discuss connections of these results to combinatorial geometry and model theory.

*Sam Bastida**

(Joint work with Nick Brettell)

List colouring is a generalisation of the traditional notion of colouring where each vertex of the graph can have a different palette. A *proper colouring* of a graph G maps each vertex of G to a colour such that adjacent vertices have different colours. A *k-list assignment* L is an assignment of a list of k colours to each vertex of G . A graph is *L-colourable* if it has a proper colouring where the colour for each vertex v is in the list $L(v)$. A graph G is *k-choosable* if for every k -list assignment L , the graph G is *L-colourable*. This notion generalises *k-colouring*: a graph is *k-colourable* if it is ϕ -colourable where ϕ maps each vertex to the same list of k colours. While some results about *k-colourability* generalise to *k-choosability*, such as Brooks' Theorem, others, such as the Four Colour Theorem, do not. Brooks' Theorem states that a connected graph G with maximum degree Δ is Δ -colourable, except when G is a complete graph or odd cycle. Stiebitz and Toft (2018) generalised Brooks' Theorem, showing that a graph G is *k-colourable*, where k is the maximum number of edge-disjoint paths between two vertices of G , except when each block of G can be obtained from complete graphs or odd cycles using Hajós joins. We consider an extension of this result to *k-choosability*, specifically in the case where $k = 3$.

A comparison of graph width parameters

Nick Brettell

Victoria University of Wellington

(Joint work with Andrea Munaro, Daniel Paulusma, and Shizhou Yang.)

The classic example of a width parameter is treewidth, which, loosely speaking, gives a measure of how tree-like a graph is. Due to Courcelle’s theorem, many problems are known to be polynomial-time solvable for a class of graphs with bounded treewidth. Say that a parameter p is *less restrictive* than a parameter q if there exists a function f such that $p(G) \leq f(q(G))$ for every graph G (it is “less restrictive” in the sense that a class may have bounded p -width but unbounded q -width). These days, there is a rich landscape of width parameters that are less restrictive than treewidth, but, like treewidth, facilitate efficient algorithms. In this talk, we’ll be interested in clique-width, mim-width, sim-width, and tree-independence number. I’ll give a brief introduction to each of these parameters, and touch on why they are of interest. We’ll then compare them when restricted to a class of graphs with no $K_{t,t}$ subgraph, the class of line graphs, and the common generalisation of the class of graphs with no induced $K_{t,t}$ subgraph. In particular, Gurski and Wanke (2000) showed that although clique-width is less restrictive than treewidth, these parameters are equivalent for graphs with no $K_{t,t}$ subgraph. Gurski and Wanke (2007) also showed that a class of graphs has bounded treewidth if and only if the corresponding class of line graphs has bounded clique-width. We generalise these results to mim-width, sim-width, and tree-independence number.

*Yudhistira Andersen Bunjamin**

(Joint work with Oden Petersen)

In this talk, we will present constructions for some 3-GDDs with two group sizes where one group size is a multiple of the other group size. The talk will have a particular focus on how some recent advancements regarding the existence of 4-GDDs with two group sizes have enabled the construction of some infinite families of 3-GDDs with two group sizes.

Covering Arrays via Finite Fields

Charles Colbourn

Arizona State University

In order to construct covering arrays of strength t and index λ on q symbols, one effective and well-studied method forms a base array with “few” rows whose entries are elements of \mathbb{F}_q^t . Each row of the base array underlies q^t rows of the covering array. A t -tuple T of columns is covering in a row of the base array when the corresponding q^t rows of the covering array contain each of the q^t symbol tuples in T . When every t -tuple of columns is covering in at least λ rows, the base array is a covering perfect hash family (CPHF_λ). When λ is ‘small’ and q is ‘large’, CPHFs yield the best probabilistic upper bounds on sizes of covering arrays and the best current construction algorithms. In this talk we revise the conditions on CPHFs to account for the partial coverage arising from non-covering t -tuples of columns. This improves the quality of the bounds on covering array sizes, particularly when λ is ‘large’ or q is ‘small’.

A strongly regular graph co-spectral and non-isomorphic to $\text{NO}^+(8, 2)$

Ian De Beule

Vrije Universiteit Brussel

The graph $\text{NO}^+(8, 2)$ is strongly regular with parameters $(120, 63, 30, 36)$. It can be constructed using a quadratic form of Witt index 4 on $\text{GF}(2)^8$. Then its vertices are the set of non-singular vectors. Two vertices are adjacent if and only if they are orthogonal with relation to the quadratic form. Its automorphism group is $\text{P}\Gamma\text{O}^+(8, 2)$.

In their recent book – Strongly Regular Graphs – Brouwer and Van Maldeghem mention the existence of a non-isomorphic, strongly regular graph with the same parameters, admitting $\text{Sym}(7)$ as automorphism group. In this talk we discuss how the adjacency relation of $\text{NO}^+(8, 2)$ can be modified to obtain this graph, it turns out that the unique ovoid (and spread) of the triality quadric $\text{Q}^+(7, 2)$ plays a central role. We also discuss further interesting properties such as that fact the cliques and co-cliques get switched by modifying the adjacency relation of $\text{NO}^+(8, 2)$.

Zhaochen Ding*

Two finite groups L_1 and L_2 are called compatible if there is a group G with two isomorphic normal subgroups N_1 and N_2 such that $G/N_1 \cong L_1$ and $G/N_2 \cong L_2$. In this talk, we will discuss some recent work (joint with Gabriel Verret) on compatibility of groups, including a new construction based on inverse limits.

Proper Minor-Closed Classes of Graphs have Assouad-Nagata Dimension 2

Marc Distel*

Monash University

Asymptotic dimension and Assouad-Nagata dimension are measures of the large-scale shape of a class of graphs. Bonamy et al. [J. Eur. Math. Society] showed that any proper minor-closed class has asymptotic dimension 2, dropping to 1 only if the treewidth is bounded. We improve this result by showing it also holds for the stricter Assouad-Nagata dimension. We also characterise when subdivision-closed classes of graphs have bounded Assouad-Nagata dimension.

Erdős-Ko-Rado theorems for finite general linear groups

Alena Ernst*

Paderborn University

(Joint work with Kai-Uwe Schmidt)

We call a subset Y of the finite general linear group $\mathrm{GL}(n, q)$ *t-intersecting* if $\mathrm{rk}(x - y) \leq n - t$ for all $x, y \in Y$. In this talk we give upper bounds on the size of *t-intersecting* sets and characterise the extremal cases that attain the bound. This is a q -analog of the corresponding result for the symmetric group, which was conjectured by Deza and Frankl in 1977 and proved by Ellis, Friedgut, and Pilpel in 2011. The results are obtained by using eigenvalue techniques and the theory of association schemes plays a crucial role.

Enumeration of the Hosoya Index in Anthracene System

*Muhammad Talha Farooq**

King Mongkut's University of Technology Thonburi, Bangkok, Thailand

In chemical graph theory, topological indices play an important role and have various uses in quantitative structure–property relationship (QSPR) as well as quantitative structure–activity relationships (QSARs). The Hosoya index is one of them, performing a molecular descriptor in mathematical chemistry. Therefore, progressing with investigations requires computing the Hosoya index of distinct molecular graphs. This research presents a computational method for determining the Hosoya index of triple benzenoid systems (Anthracene) by using transfer matrix approach and the Hosoya vector.

Saul Freedman

(Joint work with John Bamberg and Michael Giudici)

The synchronisation hierarchy of finite permutation groups, introduced by Araújo, Cameron and Steinberg in 2017, consists of classes of groups lying between 2-transitive groups and primitive groups. This includes the classes of synchronising and separating groups, defined in terms of combinatorial properties of related graphs, and the class of spreading groups, defined in terms of sets and multisets of permuted points. Araújo et al. proved that the members of these classes are primitive of almost simple, affine or diagonal type. In addition, Bray, Cai, Cameron, Spiga and Zhang showed in 2020 that any such diagonal type group must have socle $T \times T$ for some non-abelian finite simple group T . In this talk, we prove that no spreading group of diagonal type exists, by considering transitive actions (and several character tables) of the non-abelian finite simple groups.

Levenshtein's conjecture for sequence covering arrays

*Dani Gentle**

Monash University

A sequence covering array is a set of permutations of the v -element alphabet $\{0, \dots, v-1\}$ such that every sequence of t distinct symbols of the alphabet appears in the specified order in at least one permutation. A key conjecture in this area attributed to Levinshtein concerns when it is possible to build such an array in which each sequence appears in exactly one permutation. In this talk, I will discuss existing results on this conjecture, and present new results for the next open case of the conjecture.

Existence of Latin Squares with Constrained Transversals

*Afsane Ghafari Baghestani**

Monash University

A Latin Square is an $n \times n$ array where entries are chosen from the set $\{1, 2, \dots, n\}$ with the property that every symbol appears exactly once in every row and column. A transversal of such a square is defined to be a selection of n entries, one from each row and each column, where we choose every symbol exactly once. Let k be any positive integer. We construct infinitely many latin squares of even order that have at least one transversal, yet all transversals coincide on k entries

UNSW Sydney

I will discuss some work on finding asymptotic enumeration formulae for directed hypergraphs where the in-degrees and out-degrees of the vertices, and the head and tail sizes for the edges are all specified. If at least one of these four sequences is regular and the entries are not too large then the result follows easily from asymptotic enumeration formulae for sparse bipartite graphs. Otherwise we need a stricter assumption on the maximum degrees and maximum head/tail sizes, and the proof involves a martingale argument.

Monash University

An H -decomposition of a graph Γ is a partition of its edge set into subgraphs isomorphic to H . A transitive decomposition is a special kind of H -decomposition that is highly symmetrical in the sense that the subgraphs (copies of H) are preserved and transitively permuted by a group of automorphisms of Γ . This paper concerns transitive H -decompositions of the graph $K_n \square K_n$ where H is a path. When n is an odd prime, we present a construction for a transitive path decomposition where the paths in the decomposition are arbitrary large.

Large sets of infinite-dimensional q -Steiner systems

Dan Hawtin

University of Rijeka

Let V be a vector space over the finite field \mathbb{F}_q . An $S(t, k, V)_q$ is a collection \mathcal{B} of k -spaces of V such that every t -space of V is contained in a unique element of \mathcal{B} . An $LS(t, k, V)_q$ is a partition of the k -dimensional subspaces of V into $S(t, k, V)_q$ systems. In 1995, Cameron proved that if V has infinite dimension then an $LS(t, k, V)_q$ exists for all positive integers t, k with $t < k$. We give an explicit construction of an $LS(t, t + 1, V)_q$ for all prime powers q , all positive integers t , and where V has countably infinite dimension.

Robert Hickingbotham*

Graph product structure theory describes complex graphs in terms of products of simpler graphs. In this talk, I will introduce this subject and talk about a new tool called ‘blocking partitions.’ I’ll show how this tool can be used to prove stronger product structure theorems for powers of planar graphs as well as k -planar graphs, resolving open problems of Dujmović, Morin and Wood, and Ossona de Mendez.

Contributed talks
Schedule

Kumamoto University

In this talk, we propose some representations of non-representable matroids by using matrices over finite rings. For this end, we adopted modular independence, introduced by Y.H. Park as one of the generalizations of linearly independence. It was originally defined over the ring \mathbb{Z}_{p^e} of integers modulo p^e , where p is a prime and $e \in \mathbb{Z}_{>0}$, and then generalized to the case of Frobenius rings by S.T. Dougherty and H. Liu. We restrict ourselves to local rings R with the unique maximal ideal \mathfrak{m} , where the vectors $\mathbf{v}_1, \dots, \mathbf{v}_k \in R^n$ are said to be *modular independent* if $\sum a_i \mathbf{v}_i = \mathbf{0}$ implies $a_i \in \mathfrak{m}$ for all i . We will provide some conditions for a matrix over a finite ring to yield some matroid using modular independence. We also show that some well-known non-representable matroids can be represented in this way.

Cumulant expansion for counting Eulerian orientations

Mikhail Isaev

Monash University

We consider the problem of enumerating Eulerian orientations of a given graphs, that is, the orientations of its edges such that every vertex has the same in-degree and out-degree. This problem is $\#P$ -hard and corresponds to the crucial partition function in so-called "ice-type models" in statistical physics. In this work, we derive an asymptotic formula for approximating the number of Eulerian orientations of a graph with good expansion properties up to a multiplicative error $O(n^{-c})$, where c is an arbitrary fixed constant. The answer is in terms of cumulants of a multidimensional polynomial of Gaussian random variables. The proof relies on the new tail bound for the cumulant expansion series, which is of independent interest.

Safe Sets and Dominating Sets of Graphs

Pawaton Kaemarwichanurat

King Mongkut's University of Technology Thonburi, Bangkok, Thailand

(Joint work with Shinya Fujita and Furuya Michitaka (Yokohama City University))

A subset S of vertices of a graph G is a safe set if, for a component H of $G - S$ and a component C of $G[S]$, we have $|V(H)| \leq |V(C)|$ whenever there is an edge joining vertices between H and C . Moreover, if the subgraph of G induced by safe set S , $G[S]$, is connected, then S is a connected safe set. The minimum cardinality of a safe set of G is called the safe number of G and is denoted by $s(G)$. Similarly, the minimum cardinality of a connected safe set of G is called the connected safe number of G and is denoted by $s_c(G)$. A subset D of vertices of a graph G is a dominating set of G if every vertex in $V(G) - D$ is adjacent to a vertex in D . Moreover, if $G[D]$ is connected, then D is called a connected dominating set of G . The minimum cardinality of a dominating set of G is called the domination number of G and is denoted by $\gamma(G)$. Similarly, the minimum cardinality of a connected dominating set of G is called the connected domination number of G and is denoted by $\gamma_c(G)$. In this paper, we prove that if G is a graph with the maximum degree Δ , then

$$f(\Delta) \leq s(G) \leq \lceil \frac{\gamma(G)(\Delta + 1)}{2} \rceil$$

where $f(\Delta) = \frac{\gamma+6}{3}$ when $\Delta = 2$ and $f(\Delta) = \frac{\Delta^2 - 2\Delta - 3 + \sqrt{(2\Delta - \Delta^2 + 3)^2 + 4(3\Delta + \gamma(G))(\Delta - 2)}}{2(\Delta - 2)}$ when $\Delta \geq 3$. Moreover, for a connected graph G , we have

$$g(\Delta) \leq s_c(G) \leq \lceil \frac{\gamma_c(G)(\Delta - 1) + 2}{2} \rceil$$

where $g(\Delta) = \frac{\gamma_c(G)+2}{3}$ when $\Delta = 2$ and $g(\Delta) = \frac{\Delta - 5 + \sqrt{\Delta^2 - 2\Delta + 4(\Delta - 2)\gamma_c(G) + 9}}{2(\Delta - 2)}$ when $\Delta \geq 3$. The upper bounds are shown to be sharp for some $\gamma(G)$, $\gamma_c(G)$ and Δ . We also characterize all graphs satisfying each lower bound.

Common and Sidorenko linear patterns

Nina Kamčev

University of Zagreb

(Joint work with Anita Liebenau and Natasha Morrison)

Several classical results in Ramsey theory (including famous theorems of Schur, van der Waerden, Rado) deal with finding monochromatic linear patterns in two-colourings of the integers. Our topic will be quantitative extensions of such results. A linear system L over \mathbb{F}_q is *common* if the number of monochromatic solutions to $L = 0$ in any two-colouring of \mathbb{F}_q^n is asymptotically at least the expected number of monochromatic solutions in a random two-colouring of \mathbb{F}_q^n . Motivated by existing results for specific systems (such as Schur triples and arithmetic progressions), as well as extensive research on common and Sidorenko graphs, the systematic study of common systems of linear equations was recently initiated by Saad and Wolf. Fox, Pham and Zhao characterised common linear equations. A parallel concept of *Sidorenko* systems has also been investigated.

We will survey fundamental results on linear patterns and graphs, as well as recent progress towards a classification of common systems of two or more linear equations. For instance, any system containing a four-term arithmetic progression is uncommon.

Designs in the generalised symmetric group

*Lukas Klawuhn**

Paderborn University

(Joint work with Kai-Uwe Schmidt)

It is known that the notion of a transitive subgroup of a permutation group G extends naturally to the subsets of G . We study transitive subsets of the wreath product $C_r \wr S_n$ of generalised permutations acting on subsets of $\{1, \dots, n\}$ whose elements are coloured with one of r possible colours. This includes the symmetric group for $r = 1$ and the hyperoctahedral group for $r = 2$. The group $C_r \wr S_n$ can also be interpreted as the symmetry group of a regular polytope for every r and this gives rise to an intuitively accessible definition of transitivity. We consider different notions of transitivity in $C_r \wr S_n$ and interpret these algebraically as designs in the conjugacy class association scheme of $C_r \wr S_n$ using representation theory. We also give constructions showing that there exist transitive subsets of $C_r \wr S_n$ that are small compared to the size of the group. Many of these results extend results previously known for the symmetric group S_n .

On the directed Oberwolfach problem with two tables

*Alice Lacaze-Masmonteil**

University of Ottawa

(Joint work with Daniel Horsley)

A $(\vec{C}_{m_1}, \vec{C}_{m_2})$ -factor of a directed graph G is a spanning subdigraph of G comprised of two disjoint directed cycles of lengths m_1 and m_2 . In this talk, we will be constructing a decomposition of the complete symmetric digraph K_n^* into $(\vec{C}_{m_1}, \vec{C}_{m_2})$ -factors when $m_1 + m_2 = n$, $m_1 \in \{4, 6\}$, and $m_2 \geq 8$ is even. In conjunction with recent results of Kadri and Šajna (2023+), this gives rise to a complete solution to the two-table case of the directed Oberwolfach problem.

Jesse Lansdown

(Joint work with John Bamberg, Michael Giudici, and Gordon Royle.)

The class of *spreading* permutation groups lies inbetween the 2-transitive and primitive groups. Similar to a primitive group being defined by the absence of any invariant partition, a spreading group is defined by the absence of any set-multiset pair satisfying certain properties. If however a suitable set-multiset pair exists then it is called a “witness” and the group is *nonspreading*. In this talk I will consider how to construct witnesses, in particular using techniques inspired by the “AB-Lemma” used to construct hemisystems in finite geometry.

Self-avoiding walks on graphs with infinitely many ends

Florian Lehner

The University of Auckland

(Joint work with Lindorfer and Panagiotis)

The self-avoiding walk is a model from statistical physics which has been studied extensively on integer lattices. Over the last few decades, the study of self-avoiding walks on more general graphs, in particular graphs with a high degree of symmetry such as Cayley graphs of finitely generated groups, has received increasing attention.

In this talk, we focus on graphs with more than one end; intuitively these can be thought of as having some large-scale tree structure. This tree structure allows us to decompose self-avoiding walks into smaller, more manageable pieces, and answer questions for graphs with more than one end whose answers for lattices currently seem out of reach.

The talk will be aggressively non-technical. No prior knowledge of self-avoiding walks will be assumed.

The second largest eigenvalue of non-normal Cayley graphs on symmetric groups generated by cycles

Yuxuan Li*

The University of Melbourne

Aldous' Spectral Gap Conjecture states that the second largest eigenvalue of each connected Cayley graph on the symmetric group S_n with respect to a set of transpositions is attained by the standard representation of S_n . This celebrated conjecture, which was proposed in 1992 and completely proved in 2010, has inspired much interest in determining the second largest eigenvalue of Cayley graphs on S_n . For $1 \leq r < k < n$, let $C(n, k; r)$ be the set of k -cycles of S_n which move every $i \in \{1, 2, \dots, r\}$. It is conjectured that the non-normal Cayley graph $\text{Cay}(S_n, C(n, k; r))$ has the Aldous property, that is, its strictly second largest eigenvalue is achieved by the standard representations of S_n . In this talk, I will introduce the latest research developments about this conjecture, which is based on collaborative work with Binzhou Xia and Sanming Zhou.

Tensor representation of semifields and commuting polarities

Stefano Lia

University College Dublin

Finite semifields correspond to nonsingular threefold tensors and as such they admit different representation in projective spaces. In this joint work with John Sheekey, we exploit the cyclic model for threefold tensors to obtain results on a semifield invariant called BEL-rank. We show that the cyclic model allows to represent in the same space both tensors and their contraction spaces, providing a geometric interpretation of the contraction. This provides a purely geometrical proof of Dickson classification of semifields two dimensional over their center. The investigation of the nonsingularity of tensors in this model also leads to the construction of new quasi-hermitian surfaces, arising from a pair of commuting polarities related to the semifields.

(Joint work with Joint with Peter Allen and Julia Böttcher.)

What is the smallest number of edges that a graph G can have if it contains all D -degenerate graphs on n vertices as subgraphs? A counting argument shows that this number is at least of order $n^{2-1/D}$, assuming n is large enough. We show that this is tight up to a polylogarithmic factor.

On generalisations of The Erdős-Ko-Rado Theorem for permutations

Adam Mammoliti

UNSW Sydney

The celebrated Erdős-Ko-Rado Theorem states that if $n \geq 2k$ and \mathcal{F} is a family of k -subsets of $[n]$ such that $A \cap B \neq \emptyset$ for all sets $A, B \in \mathcal{F}$, then $|\mathcal{F}| \leq \binom{n-1}{k-1}$, with equality for $n > 2k$ occurring precisely when \mathcal{F} is a family of all k -subsets containing a fixed element of $[n]$. Since its discovery, the Erdős-Ko-Rado Theorem has been generalised extensively and analogous results have been shown for structures other than sets. In particular, an analogue of the Erdős-Ko-Rado Theorem has been shown for families of permutations of $[n]$.

Brendan McKay

The exhaustive generation of classes of combinatorial objects has been a hobby of mine since my student days. After my arrival at ANU in 1983, among my first projects were to generate cubic graphs and vertex transitive graphs with Gordon Royle. Like Gordon, I'm still addicted to the field and will discuss two recent projects. One is to compile a list of graphs extremal under not containing cycles of specified lengths, to as large an order as possible. The other is to compile a library of combinatorial 2-designs.

On the non-existence of q -ary linear codes with minimum weight $d \equiv -1 \pmod{q}$

Tatsuya Maruta

Osaka Metropolitan University

(Joint work with Hitoshi Kanda and Atsuya Kato)

A q -ary linear code is an $[n, k, d]_q$ code, which is a linear code of length n , dimension k and minimum weight d over the field of order q . When an $[n, k, d]_q$ code with d divisible by q does not exist, it is most likely that an $[n - 1, k, d - 1]_q$ code does not exist as well. We give a result on a new notion “ e -locally 2-weight \pmod{q} ” for q -ary linear codes which help us to prove the non-existence of such a code applying the well-known Extension Theorem by Hill and Lizak(1995).

Equally Distributed 1-Factorisations of Graphs

Jeremy Mitchell*

The University of Queensland

The union of a pair of edge-disjoint 1-factors of a graph forms a collection of even length cycles. If t cycles formed by the union of two edge-disjoint 1-factors have lengths a_1, a_2, \dots, a_t we say the pair of 1-factors have type (a_1, a_2, \dots, a_t) , if all the pairs of 1-factors of some 1-factorisation have the same type then it is a uniform 1-factorisation. Consider a 1-factorisation \mathcal{F} of some graph and let t_1, t_2, \dots, t_m be all types of the pairs of 1-factors of \mathcal{F} . Let a_{t_i} be the number of pairs that are type t_i . If $a_{t_1} = a_{t_2} = \dots = a_{t_m} = b$ for some integer b , then we say that \mathcal{F} is an *m-equally distributed 1-factorisation* (m -ED1F) with types (t_1, t_2, \dots, t_m) . We present some results on m -ED1Fs of 3- and 4-regular circulant graphs. Finally, we impose some additional conditions on m -ED1Fs and investigate when such constrained m -ED1Fs exist for complete and complete bipartite graphs.

Đorđe Mitrović*

For a non-bipartite graph X , the automorphisms of the direct product $X \times K_2$ play an important role in understanding the automorphism group of $X \times Y$, where Y is bipartite. A graph X is unstable if $X \times K_2$ has automorphisms that do not come from automorphisms of its factors. It is non-trivially unstable if it is unstable, connected, non-bipartite and twin-free. We provide new sufficient conditions for the instability of circulant graphs, generalising previously known results. Furthermore, we classify non-trivially unstable members of several families of circulants.

Yusuke Miura

(Joint work with Koji Imamura (Kumamoto Univ.) and Tatsuya Maruta)

An n -set B in $\text{PG}(r, q)$ is a k -blocking set if every $(r - k)$ -space in $\text{PG}(r, q)$ meets B in at least one point. B is called *trivial* if B contains a k -space. Bono et al.(2021) proved that there are exactly six non-trivial minimal 2-blocking sets in $\text{PG}(4, 2)$ up to projective equivalence. We consider the non-trivial minimal 2-blocking sets in $\text{PG}(5, 2)$ and their generalizations.

Tomasz Popiel

(Joint work with Heiko Dietrich and Melissa Lee)

The Monster is the largest of the 26 sporadic finite simple groups, and is notoriously difficult to compute with, owing to a lack of sufficiently small permutation or matrix representations. As a result, various ‘basic’ facts about the Monster that are often needed for combinatorial applications of the Classification of the Finite Simple Groups have yet to be determined. In particular, the classification of the maximal subgroups of the Monster has remained uncompleted for some four decades. I shall report on recent joint work on this problem with Heiko Dietrich and Melissa Lee, involving software developed by Martin Seysen.

Prime labeling of some graphs with Eisenstein integers

Rovin B. Santos

Institute of Mathematics, University of the Philippines - Diliman

A graph on n vertices is said to admit a prime labeling if the vertices can be labeled with the first n natural numbers in a such a way that two adjacent vertices have relatively prime labels. In this paper, we define an order on the set of Eisenstein integers to extend the notion of prime labeling of graphs to the set of Eisenstein integers. Properties of the ordering are studied to come up with prime labeling of some families of graphs such as the flower, wheel, centipede, and double broom graphs.

Optimising phylogenetic diversity on phylogenetic networks

Charles Semple

University of Canterbury

(Joint work with Magnus Bordewich and Kristina Wicke)

Phylogenetic diversity (PD) is a popular measure for quantifying the biodiversity of a set of present-day taxa. This measure quantifies the extent to which the taxa spans the ‘Tree of Life’. In applications, the underlying optimisation problem is to find, for a given set S of taxa and positive integer k , a subset of S of size k that maximises the phylogenetic diversity score. Historically, PD has been typically restricted to phylogenetic trees, but it extends naturally to phylogenetic networks. In this talk, we investigate such an extension.

Arizona State University

A (d,t) -locating array is a covering array of strength t with an additional property: Any set of d level-wise t -way interactions can be distinguished from any other such set by appearing in a distinct set of rows. Locating arrays have been proposed as experimental designs for screening experiments for complex systems due to their efficiency. In this talk, we describe how a $(1,2)$ -locating array recovers main effects and two-way interactions from the measurements of a screening experiment. Preliminary results investigate the role of separation and d -efficiency in screening effectiveness.

Wipawee Tangjai

A δ -complement graph was introduced in 2022. The graph is constructed in the same way as a complement graph with a restriction on taking a complement within the set of vertices with the same degree of the graph. In this work, we give several results related to a property and a chromatic number of a δ -complement graph including bounds of the chromatic number and an exact value of the chromatic number of some special classes of graphs.

Abdullahi Umar

The study of various (sub)-semigroups of transformations/mappings has made a significant contribution to semigroup theory. The most notable classes are the THREE fundamental semigroups of transformations: the full symmetric semigroup, the partial symmetric semigroup and the symmetric inverse semigroup. In this talk, we are going to discuss some combinatorial results of some classes semigroups of (partial) contraction transformations of a finite chain, which for some curious reason(s), until very recently, little is known about.

*Jie Wang**

Suppose we have two polytopes that are combinatorially equivalent, but one decomposable, the other one indecomposable. Such polytopes are called conditionally decomposable. For a conditionally decomposable polytope, we show that the minimum number of vertices is in the range $[3d - 3, 4d - 4]$; and the minimum number of facets is obtained for $d \geq 4$. Joint work with David Yost.

Keita Yasufuku

(Joint work with Tatsuya Maruta)

We consider the problem of determining $n_q(k, d)$, the smallest possible length n for which an $[n, k, d]_q$ code of fixed dimension k and minimum weight d over the field of order q exists. We investigate the validity of Kawabata's conjecture on the achievement of the Griesmer bound for linear codes over the field of order q , especially for $q = 5$.

On linear-algebraic notions of expansion

Chuanqi Zhang*

University of Technology Sydney

A fundamental fact about bounded-degree graph expanders is that three notions of expansion—vertex expansion, edge expansion, and spectral expansion—are all equivalent. This motivates us to study to what extent such a statement is true for linear-algebraic notions of expansion.

There are two well-studied notions of linear-algebraic expansion, namely dimension expansion [1] (defined in analogy to graph vertex expansion) and quantum expansion [2, 3] (defined in analogy to graph spectral expansion). Lubotzky and Zelmanov [4] proved that the latter implies the former. We proved that the converse is false: there are dimension expanders which are not quantum expanders.

Moreover, this asymmetry is explained by the fact that there are two distinct linear-algebraic analogues of graph edge expansion. The first of these is *quantum edge expansion*, which was introduced by Hastings [5], and which he proved to be equivalent to quantum expansion. We established a new notion, termed *dimension edge expansion*, which we proved is equivalent to dimension expansion and which is implied by quantum edge expansion. Thus, the separation above is implied by a finer one: dimension edge expansion is strictly weaker than quantum edge expansion. This new notion also led to a new and more modular proof of the Lubotzky-Zelmanov result [4] that quantum expanders are dimension expanders.

[1] Boaz Barak, Russell Impagliazzo, Amir Shpilka, and Avi Wigderson. Definition and existence of dimension expanders. Discussion (no written record), 2004.

[2] Avraham Ben-Aroya and Amnon Ta-Shma. Quantum expanders and the quantum entropy difference problem. ArXiv:quant-ph/0702129, 2007.

[3] M. B. Hastings. Entropy and entanglement in quantum ground states. *Phys. Rev. B*, 76:035114, Jul 2007.

[4] Alexander Lubotzky and Efim Zelmanov. Dimension expanders. *Journal of Algebra*, 319(2):730–738, 2008.

[5] M. B. Hastings. Random unitaries give quantum expanders. *Physical Review A*, 76:032315, Sep 2007.

3

List of participants

Name	Affiliation	email address
Jack Allsop	Monash University	jack.allsop@monash.edu
Vishnuram Arumugam	UWA	vishnuram.arumugam@research.uwa.edu.au
John Bamberg	UWA	john.bamberg@uwa.edu.au
Abdul Basit	Monash University	abdul.basit@monash.edu
Samuel Bastida	Victoria University of Wellington	bastidsamu@myvuw.ac.nz
Anton Baykalov	UWA	anton.baykalov@uwa.edu.au
Nick Brettell	Victoria University of Wellington	nick.brettell@vuw.ac.nz
Thomas Britz	UNSW	britz@unsw.edu.au
Lei Chen	UWA	lei.chen@research.uwa.edu.au
Charles Colbourn	Arizona State University	colbourn@asu.edu
John Mel Dacaymat	University of the Philippines Diliman	jmdacaymat@math.upd.edu.ph
Sara Davies	The University of Queensland	sara.davies@uq.edu.au
Jan De Beule	Vrije Universiteit Brussel	Jan.De.Beule@vub.be
Ajani De Vas Gunasekara	Monash University	ajani.gunasekara@gmail.com
Alice Devillers	UWA	alice.devillers@uwa.edu.au
Zhaochen Ding	University of Auckland	dingren941@gmail.com
Marc Distel	Monash University	Marc.Distel@monash.edu
Alena Ernst	Paderborn University	alena.ernst@math.upb.de
Dani Gentle	Monash University	aidan.gentle@monash.edu
Afsane Ghafari	Monash University	afsane.ghafaribaghestani@monash.edu
Michael Giudici	UWA	michael.giudici@uwa.edu.au
Gary Greaves	Nanyang Technical University	gary@ntu.edu.sg
Catherine Greenhill	UNSW Sydney	c.greenhill@unsw.edu.au
Krystal Guo	University of Amsterdam	k.guo@uva.nl
Yusuf Hafidh	University of Melbourne	yhafidh@student.unimelb.edu.au
Hao Chuien Hang	The University of Queensland	hanghc@hotmail.com
Toru Hasunuma	Tokushima University	hasunuma@tokushima-u.ac.jp
Daniel Hawtin	University of Rijeka	dan.hawtin@gmail.com
Robert Hickingbotham	Monash University	robert.hickingbotham1@monash.edu
Masatake Hirao	Aichi Prefectural University	hirao@ist.aichi-pu.ac.jp
Koji Imamura	Kumamoto University	211d9321@st.kumamoto-u.ac.jp
Mikhail Isaev	Monash University	mikhail.isaev@monash.edu
Nina Kamčev	University of Zagreb	nina.kamcev@math.hr
Pawaton Kaemawichanurat	King Mongkut's University of Technology Thonburi	pawaton.kae@kmutt.ac.th
Shinya Kawabuchi	Kumamoto University	230d8554@st.kumamoto-u.ac.jp
Lukas Klawuhn	Paderborn University	klawuhn@math.upb.de
André Kündgen	California State University San Marcos	akundgen@csusm.edu
Alice Lacaze-Masmonteil	University of Ottawa	alaca054@uottawa.ca
Jesse Lansdown	University of Canterbury	jesse.lansdown@canterbury.ac.nz
Melissa Lee	Monash University	melissa.lee@monash.edu
Florian Lehner	University of Auckland	florian.lehner@auckland.ac.nz
Thomas Lesgourgues	UNSW	tlesgourgues@gmail.com
Yuxuan Li	The University of Melbourne	yuxuan11@student.unimelb.edu.au
Stefano Lia	University College Dublin	stefano.lia@ucd.ie
Hongyi Lyu	Monash University	hongyi.lyu1@monash.edu
Adam Mammoliti	UNSW Sydney	adam.mammoliti@outlook.com.au
Tatsuya Maruta	Osaka Metropolitan University	maruta@omu.ac.jp
Brendan McKay	Australian National University	brendan.mckay@anu.edu.au
Jeremy Mitchell	The University of Queensland	jeremy.mitchell@uq.net.au
Đorđe Mitrović	University of Auckland	dmit755@aucklanduni.ac.nz
Yusuke Miura	Osaka Metropolitan University	sd22525u@st.omu.ac.jp
Luke Morgan	UWA	luke.morgan@uwa.edu.au
Padraig Ó Catháin	Dublin City University	padraig.ocathain@dcu.ie
Tomasz Popiel	Monash University	tomasz.popiel@monash.edu
Cheryl Praeger	UWA	cheryl.praeger@uwa.edu.au
Gordon Royle	UWA	gordon.royle@uwa.edu.au

Joe Ryan	University of Newcastle	joe.ryan@newcastle.edu.au
Shohei Satake	Kumamoto University	shohei-satake@kumamoto-u.ac.jp
George Savvoudis	The University of Adelaide	george.savvoudis@adelaide.edu.au
Charles Semple	University of Canterbury	charles.semple@canterbury.ac.nz
Jacob Smith	UWA	jacob.smith@research.uwa.edu.au
Keisuke Shiromoto	Kumamoto University	keisuke@kumamoto-u.ac.jp
Violet Syrotiuk	Arizona State University	syrotiuk@asu.edu
Tibor Szabó	Freie Universität Berlin	szabo@zedat.fu-berlin.de
Wipawee Tangjai	Maharakham University	wipawee.t@msu.ac.th
Abdullahi Umar	Khalifa University of Science and Technology	abdullahi.umar@ku.ac.ae
Geertrui Van de Voorde	University of Canterbury	geertrui.vandevoorde@canterbury.ac.nz
Gabriel Verret	University of Auckland	g.verret@auckland.ac.nz
Jie Wang	Federation University	jiawang@students.federation.edu.au
Ian Wanless	Monash University	ian.wanless@monash.edu
David Wood	Monash University	david.wood@monash.edu
Binzhou Xia	University of Melbourne	binzhoux@unimelb.edu.au
Keita Yasufuku	Osaka Metropolitan University	sd23270s@st.omu.ac.jp
David Yost	Federation University	d.yost@federation.edu.au
Zhishuo Zhang	The University of Melbourne	zhishuoz@student.unimelb.edu.au
Chuanqi Zhang	University of Technology Sydney	chuanqi.zhang@student.uts.edu.au