

Computer Organization and Architecture

Arithmetic Operation

Veena Thenkanidiyoor

National Institute of Technology Goa



1

1

Recap

- Need for number and character representation
- Unsigned number representation
- Signed number representation
 - Sign-and-magnitude
 - 1's complement
 - 2's complement
- Character representation

2

2

Addition of 1-bit Numbers

$$\begin{array}{r} 0 \\ + 0 \\ \hline 0 \end{array}$$

$$\begin{array}{r} 1 \\ + 0 \\ \hline 1 \end{array}$$

$$\begin{array}{r} 0 \\ + 1 \\ \hline 1 \end{array}$$

$$\begin{array}{r} 1 \\ + 1 \\ \hline 10 \end{array}$$

2-base.

$$\begin{array}{r} 253 \\ + 77 \\ \hline 330 \end{array}$$

Sum is 0 ←
Carry-out is 1 ←

- To add 2 n-bit numbers
 - Add bit pairs starting from the LSB
 - Propagate carries toward the MSB

3

3

2's Complement Arithmetic

$n = 4$

7	0111	4	0100	-4	1100	4	0100	-4	1100
+ (-3)	+ 1101	+ 3	0011	+ (-3)	1101	+ (-3)	1101	+ 3	0011
4	10100	7	0111	-7	11001	1	10001	-1	1111
*				*		*			

- Ignoring the carry-out from the MSB in the above addition(s), gives the correct answer

4

4

Addition and Subtraction of n-bit Signed Numbers Represented in 2's Complement

- To add two numbers, add their n-bit representations, **ignoring the carry-out signal** from the most significant bit (MSB) position. The sum will be algebraically correct value in the 2's complement representation as long as the answer is in the range $-(2^{n-1})$ to $(2^{n-1} - 1)$
- To subtract two numbers X and Y, that is, to perform $X - Y$, form the **2's complement of Y** and then **add it to X**, as in rule 1. Again the result will be algebraically correct value in the 2's complement representation as long as the answer is in the range $-(2^{n-1})$ to $(2^{n-1} - 1)$

5

5

Example

$$\begin{array}{rcl}
 \text{(a)} & 0010 & (+2) \\
 & + 0011 & (+3) \\
 \hline
 & 0101 & (+5)
 \end{array}$$

$$\begin{array}{rcl}
 \text{(b)} & 0100 & (+4) \\
 & + 1010 & (-6) \\
 \hline
 & 1110 & (-2)
 \end{array}$$

$$\begin{array}{rcl}
 \text{(c)} & 1011 & (-5) \\
 & + 1110 & (-2) \\
 \hline
 & 1001 & (-7)
 \end{array}$$

$$\begin{array}{rcl}
 \text{(d)} & 0111 & (+7) \\
 & + 1101 & (-3) \\
 \hline
 & 0100 & (+4)
 \end{array}$$

Rule 1

6

6

Example

(e)

$$\begin{array}{r}
 1101 \quad (-3) \\
 - 1001 \quad (-7) \\
 \hline
 \end{array}
 \xRightarrow{+4}
 \begin{array}{r}
 1101 \\
 + 0111 \\
 \hline
 0100 \quad (+4)
 \end{array}$$

(f)

$$\begin{array}{r}
 0010 \quad (+2) \\
 - 0100 \quad (+4) \\
 \hline
 \end{array}
 \xRightarrow{-2}
 \begin{array}{r}
 0010 \\
 + 1100 \\
 \hline
 1110 \quad (-2)
 \end{array}$$

Rule 2

7

7

Example

(g)

$$\begin{array}{r}
 0110 \quad (+6) \\
 - 0011 \quad (+3) \\
 \hline
 \end{array}
 \xRightarrow{-9}
 \begin{array}{r}
 0110 \\
 + 1101 \\
 \hline
 0011 \quad (+3)
 \end{array}$$

(h)

$$\begin{array}{r}
 1001 \quad (-7) \\
 - 1011 \quad (-5) \\
 \hline
 \end{array}
 \xRightarrow{+2}
 \begin{array}{r}
 1001 \\
 + 0101 \\
 \hline
 1110 \quad (-2)
 \end{array}$$

Rule 2

8

8

Example

(i) 1 0 0 1 (- 7) - 0 0 0 1 (+1) <hr style="width: 100%;"/>		1 0 0 1 + 1 1 1 1 <hr style="width: 100%;"/> 1 0 0 0 (-8)
(j) 0 0 1 0 (+2) - 1 1 0 1 (-3) <hr style="width: 100%;"/>		0 0 1 0 + 0 0 1 1 <hr style="width: 100%;"/> 0 1 0 1 (+5)

Rule 2

9

9

2's Complement Representation

- Subtrahend to be 2's-complemented
- Subtraction is proceeded like how addition done
- Simplicity of adding or subtracting signed numbers in 2's-complement representation

(i) 1 0 0 1 (- 7) - 0 0 0 1 (+1) <hr style="width: 100%;"/>		1 0 0 1 + 1 1 1 1 <hr style="width: 100%;"/> 1 0 0 0 (-8)
(j) 0 0 1 0 (+2) - 1 1 0 1 (-3) <hr style="width: 100%;"/>		0 0 1 0 + 0 0 1 1 <hr style="width: 100%;"/> 0 1 0 1 (+5)

10

10

Arithmetic Overflow in Integer Arithmetic

- When the result of an arithmetic operation is outside the representable range, an arithmetic overflow has occurred

- Addition

- Unsigned integers: 0 to $(2^n - 1)$

$n=4$ 0 to 2^4-1
 0 to 15

1) $8 + 9 = 17$
 Binary: $1000 + 1001 = 10001$
 Carry - MSB

2) $7 + 7 = 14$
 Binary: $0111 + 0111 = 1110$

3) $15 + 3 = 18$
 Binary: $1111 + 0011 = 10010$
 Carry - MSB
 ⇒ Arithmetic Overflow ✓

11

11

Arithmetic overflow-2's Complement Addition

- Range: $-(2^{n-1})$ to $(2^{n-1} - 1)$

$n=4$ $-8 \rightarrow 7$

① $(+7) + (+4)$
 Binary: $0111 + 0100 = 1011$
 Result: -5

② $(-4) + (-6)$
 Binary: $1100 + 1010 = 10110$
 Result: $+6$

12

12

Arithmetic overflow-2's Complement Addition

- Range: $-(2^{n-1})$ to $(2^{n-1} - 1)$

3) (-8)
 $+ (+7)$

 -1

4) (-7)
 $+ (+7)$

 0

5) (-8)
 $+ (+7)$

 -1

6) (-7)
 $+ (+7)$

 0

7) (-8)
 $+ (+7)$

 -1

8) (-7)
 $+ (+7)$

 0

9) (-8)
 $+ (+7)$

 -1

10) (-7)
 $+ (+7)$

 0

11) (-8)
 $+ (+7)$

 -1

12) (-7)
 $+ (+7)$

 0

13) (-8)
 $+ (+7)$

 -1

14) (-7)
 $+ (+7)$

 0

15) (-8)
 $+ (+7)$

 -1

16) (-7)
 $+ (+7)$

 0

17) (-8)
 $+ (+7)$

 -1

18) (-7)
 $+ (+7)$

 0

19) (-8)
 $+ (+7)$

 -1

20) (-7)
 $+ (+7)$

 0

21) (-8)
 $+ (+7)$

 -1

22) (-7)
 $+ (+7)$

 0

23) (-8)
 $+ (+7)$

 -1

24) (-7)
 $+ (+7)$

 0

25) (-8)
 $+ (+7)$

 -1

26) (-7)
 $+ (+7)$

 0

27) (-8)
 $+ (+7)$

 -1

28) (-7)
 $+ (+7)$

 0

29) (-8)
 $+ (+7)$

 -1

30) (-7)
 $+ (+7)$

 0

31) (-8)
 $+ (+7)$

 -1

32) (-7)
 $+ (+7)$

 0

33) (-8)
 $+ (+7)$

 -1

34) (-7)
 $+ (+7)$

 0

35) (-8)
 $+ (+7)$

 -1

36) (-7)
 $+ (+7)$

 0

37) (-8)
 $+ (+7)$

 -1

38) (-7)
 $+ (+7)$

 0

39) (-8)
 $+ (+7)$

 -1

40) (-7)
 $+ (+7)$

 0

41) (-8)
 $+ (+7)$

 -1

42) (-7)
 $+ (+7)$

 0

43) (-8)
 $+ (+7)$

 -1

44) (-7)
 $+ (+7)$

 0

45) (-8)
 $+ (+7)$

 -1

46) (-7)
 $+ (+7)$

 0

47) (-8)
 $+ (+7)$

 -1

48) (-7)
 $+ (+7)$

 0

49) (-8)
 $+ (+7)$

 -1

50) (-7)
 $+ (+7)$

 0

51) (-8)
 $+ (+7)$

 -1

52) (-7)
 $+ (+7)$

 0

53) (-8)
 $+ (+7)$

 -1

54) (-7)
 $+ (+7)$

 0

55) (-8)
 $+ (+7)$

 -1

56) (-7)
 $+ (+7)$

 0

57) (-8)
 $+ (+7)$

 -1

58) (-7)
 $+ (+7)$

 0

59) (-8)
 $+ (+7)$

 -1

60) (-7)
 $+ (+7)$

 0

61) (-8)
 $+ (+7)$

 -1

62) (-7)
 $+ (+7)$

 0

63) (-8)
 $+ (+7)$

 -1

64) (-7)
 $+ (+7)$

 0

65) (-8)
 $+ (+7)$

 -1

66) (-7)
 $+ (+7)$

 0

67) (-8)
 $+ (+7)$

 -1

68) (-7)
 $+ (+7)$

 0

69) (-8)
 $+ (+7)$

 -1

70) (-7)
 $+ (+7)$

 0

71) (-8)
 $+ (+7)$

 -1

72) (-7)
 $+ (+7)$

 0

73) (-8)
 $+ (+7)$

 -1

74) (-7)
 $+ (+7)$

 0

75) (-8)
 $+ (+7)$

 -1

76) (-7)
 $+ (+7)$

 0

77) (-8)
 $+ (+7)$

 -1

78) (-7)
 $+ (+7)$

 0

79) (-8)
 $+ (+7)$

 -1

80) (-7)
 $+ (+7)$

 0

81) (-8)
 $+ (+7)$

 -1

82) (-7)
 $+ (+7)$

 0

83) (-8)
 $+ (+7)$

 -1

84) (-7)
 $+ (+7)$

 0

85) (-8)
 $+ (+7)$

 -1

86) (-7)
 $+ (+7)$

 0

87) (-8)
 $+ (+7)$

 -1

88) (-7)
 $+ (+7)$

 0

89) (-8)
 $+ (+7)$

 -1

90) (-7)
 $+ (+7)$

 0

91) (-8)
 $+ (+7)$

 -1

92) (-7)
 $+ (+7)$

 0

93) (-8)
 $+ (+7)$

 -1

94) (-7)
 $+ (+7)$

 0

95) (-8)
 $+ (+7)$

 -1

96) (-7)
 $+ (+7)$

 0

97) (-8)
 $+ (+7)$

 -1

98) (-7)
 $+ (+7)$

 0

99) (-8)
 $+ (+7)$

 -1

100) (-7)
 $+ (+7)$

 0

13

13

Arithmetic overflow-2's Complement Addition

- Range: $-(2^{n-1})$ to $(2^{n-1} - 1)$
- The carry-out signal from the sign-bit position is not a sufficient indicator of overflow when adding signed numbers
- Overflow can occur only when adding two numbers that have the same sign

14

14

Arithmetic overflow-2's Complement Addition

- Examine the signs of 2 summands X and Y and the sign of the result
 - When both operands X and Y have same sign, an overflow occurs when the sign of sum S is not the same as the signs of X and Y indicated

- Observe carry to $C(n-1)$ and $C(n)$
 - Different: Overflow
 - Same: No overflow

$$\begin{array}{r} +2 \\ + (+2) \\ \hline +4 \end{array}$$

$$\begin{array}{r} 0010 \\ 0010 \\ \hline 0100 \end{array}$$

15

15

To Summarize

- Adding 1- bit numbers
- Addition operation on signed numbers represented in 2's complement
- Addition and Subtraction of n-bit Signed Numbers Represented in 2's Complement
- Arithmetic overflow

16

16

Reference

- Carl Hamacher, Zvonko Vranesic and Safwat Zaky, "**Computer Organization**", 5th Edition, Tata McGraw Hill, 2002

17

17

Thank You

18

18