

## Improper Integrals of the Second Kind, Continued

We'll continue our discussion of integrals of functions which have singularities at finite values; for example,  $f(x) = \frac{1}{x}$ . If  $f(x)$  has a singularity at 0 we **define**

$$\int_0^1 f(x) dx = \lim_{a \rightarrow 0^+} \int_a^1 f(x) dx.$$

As before, we say the integral *converges* if this limit exists and *diverges* if not.




Figure 1: Area under the graph of  $y = \frac{1}{x}$ .

We treat this infinite vertical “tail” the same way we treated horizontal tails. Figure 1 shows a function whose value goes to positive infinity as  $x$  goes to zero from the right hand side. We don't know whether the area under its graph between 0 and 1 is going to be infinite or finite, so we cut it off at some point  $a$  where we know it will be finite. Then we let  $a$  go to zero from above ( $a \rightarrow 0^+$ ) and see whether the area under the curve between  $a$  and 1 goes to infinity or to some finite limit.

**Example:**  $\int_0^1 \frac{dx}{\sqrt{x}}$

$$\begin{aligned} \int_0^1 \frac{dx}{\sqrt{x}} &= \int_0^1 x^{-1/2} dx \\ &= \left. \frac{1}{1/2} x^{1/2} \right|_0^1 \\ &= \left. 2x^{1/2} \right|_0^1 \end{aligned}$$

$$\begin{aligned}
&= 2 \cdot 1^{1/2} - 2 \cdot 0^{1/2} \\
&= 2.
\end{aligned}$$

This is a convergent integral.

**Example:**  $\int_0^1 \frac{dx}{x}$

$$\begin{aligned}
\int_0^1 \frac{dx}{x} &= \ln x \Big|_0^1 \\
&= \ln 1 - \ln 0^+ \\
&= 0 - (-\infty) \\
&= +\infty.
\end{aligned}$$

This integral diverges.

**In general:**

$$\begin{aligned}
\int_0^1 \frac{dx}{x^p} &= \left. \frac{x^{-p+1}}{-p+1} \right|_0^1 \quad (\text{for } p \neq 1) \\
&= \frac{1^{-p+1}}{-p+1} - \frac{0^{-p+1}}{-p+1} \\
&= \begin{cases} \frac{1}{1-p} & \text{if } p < 1 \\ \text{diverges} & \text{if } p \geq 1. \end{cases}
\end{aligned}$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.01SC Single Variable Calculus  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.