

1 Categories

Definition 1.1 (Category). A category, \mathcal{A} is defined to have each of the following,

- (i) A collection of objects, denoted by $\text{ob}(\mathcal{A})$ and written $A, B, C \in \mathcal{A}$.
Such that each object has an ‘identity’, $1_A \in \mathcal{A}(A, A)$, $1_B \in \mathcal{A}(B, B)$, $1_C \in \mathcal{A}(C, C)$
- (ii) For each pair of objects, a collection of ‘links’/morphisms between them, denoted by $\mathcal{A}(A, B)$ and written as $f \in \mathcal{A}(A, B)$ $g \in \mathcal{A}(B, C)$. Such that,
 - (a) morphisms with matching domain, co-domain can be ‘chained’/composed $(g, f) = g \circ f$
 - (b) with this composition being associative, $(h \circ g) \circ f = h \circ (g \circ f)$
 - (c) and they are ‘fixed’ by the identity $f \circ 1_A = f = 1_B \circ f$

Example 1.1. Non-trivial Identity Consider the objects to be groups, and morphisms to be direct product between them:

- i $\text{ob}(\mathcal{A}) = \{G \mid G \text{ is a group}\}$
- ii $\mathcal{A}(A, B) := A \times B$
- iii $\mathcal{A}(B, C) \circ \mathcal{A}(A, B) \mapsto \mathcal{A}(A, C)$

So, there’s a unique morphism between any two objects i.e groups. And the identity morphism,

$\forall A, B \in \mathcal{A}$, if $f \in \mathcal{A}(A, B)$, then $f \circ 1_A \in \mathcal{A}(A, B) \times \mathcal{A}(A, A) \mapsto \mathcal{A}(A, B)$ and $1_B \circ f \in \mathcal{A}(B, B) \times \mathcal{A}(A, B) \mapsto \mathcal{A}(A, B)$

Thus, $\text{ob}(\mathcal{A})$ along with \circ is actually a group. And hence has a unique inverse. [But how exactly?](#)

Example 1.2. Set The objects are defined to be sets, and morphisms are the functions between them, with the usual composition law:

- i $\text{ob}(\mathcal{A}) = \{S \mid S \text{ is a set}\}$
- ii $(f : A \mapsto B) \in \mathcal{A}(A, B)$
- iii $(g \in \mathcal{A}(B, C)) \circ (f \in \mathcal{A}(A, B)) \mapsto g(f) \in \mathcal{A}(A, C)$

Example 1.3. Grp

- i $\text{ob}(\mathcal{A}) = G \mid G \text{ is a group}$
- ii

2 Functors

Definition 2.1. Functor A functor is a map between categories, written $F : \mathcal{A} \mapsto \mathcal{B}$, consists :

- (i) function taking objects of \mathcal{A} to those of \mathcal{B} i.e. $\text{ob}(\mathcal{A}) \mapsto \text{ob}(\mathcal{B})$. Written as $A \mapsto F(A)$.
- (ii) associative, identity-preserving function taking links between objects of \mathcal{A} to those for \mathcal{B} , $f \mapsto F(f)$, i.e.

$$\forall A, B \in \mathcal{A}, \mathcal{A}(A, B) \mapsto \mathcal{B}(F(A), F(B)) \text{ such that } (a) f \in \mathcal{A}(A, B), g \in \mathcal{A}(B, C) \implies F(g \circ f) = F(g) \circ F(f) = F(g \circ f)$$

$$(b) A \in \mathcal{A} \implies F(1_A) = 1_{F(A)}$$

Example 2.1. Forgetful Functors

- (a)

3 Natural Isomorphisms

Example 3.1.