

# Yoneda Lemma and Quasi-Uniform Spaces

Ayush Rawat

Shiv Nadar University

22 June 2020

# Abstract

We work out the details of the proof for Yoneda Lemma using the text from [3]. Roughly speaking, Yoneda Lemma allows us to embed locally small categories into *Set* via representable functors. We then give two consequences of the Lemma: first is to show that Cayley's theorem from group theory is a particular case of Yoneda Lemma, and second is to derive Yoneda Embedding, a fully faithful functor from locally small categories to their presheaf category. Further, we discuss quasi-uniform spaces from the paper [1]. Here we discuss categories of quasi-uniform spaces and Promodules. We define the Yoneda embedding and prove a (weak) Yoneda Lemma for quasi-uniform spaces. We stop our work here; though the paper goes on a step further to discuss the Cauchy completion monad for quasi-uniform spaces.

# Representables

## Definition

For any category  $\mathcal{A}$ , it's opposite category,  $\mathcal{A}^{op}$  is the category having the objects of  $\mathcal{A}$ . And for objects  $A, B \in \mathcal{A}$ , a morphism  $f \in \mathcal{A}^{op}(A, B)$  if and only if there is a morphism  $g \in \mathcal{A}(B, A)$ .

# Representables

## Definition

For any category  $\mathcal{A}$ , it's opposite category,  $\mathcal{A}^{op}$  is the category having the objects of  $\mathcal{A}$ . And for objects  $A, B \in \mathcal{A}$ , a morphism  $f \in \mathcal{A}^{op}(A, B)$  if and only if there is a morphism  $g \in \mathcal{A}(B, A)$ .

## Proposition

For a locally small category  $\mathcal{A}$ , fixing an object  $A \in \mathcal{A}$  gives a functor,  $H_A : \mathcal{A}^{op} \rightarrow \mathbf{Set}$  defined as:

- ① For any object  $B \in \mathcal{A}$ ,  $H_A(B) := \mathcal{A}(B, A)$ .
- ② For any morphism,  $g : X \rightarrow Y$  in  $\mathcal{A}$ ,

$H_A(g) : \mathcal{A}(Y, A) \rightarrow \mathcal{A}(X, A)$  is given by  $p \mapsto p \circ g$ .

# Representables

## Definition

For any category  $\mathcal{A}$ , it's opposite category,  $\mathcal{A}^{op}$  is the category having the objects of  $\mathcal{A}$ . And for objects  $A, B \in \mathcal{A}$ , a morphism  $f \in \mathcal{A}^{op}(A, B)$  if and only if there is a morphism  $g \in \mathcal{A}(B, A)$ .

## Proposition

For a locally small category  $\mathcal{A}$ , fixing an object  $A \in \mathcal{A}$  gives a functor,  $H_A : \mathcal{A}^{op} \rightarrow \mathbf{Set}$  defined as:

- ① For any object  $B \in \mathcal{A}$ ,  $H_A(B) := \mathcal{A}(B, A)$ .
- ② For any morphism,  $g : X \rightarrow Y$  in  $\mathcal{A}$ ,

$H_A(g) : \mathcal{A}(Y, A) \rightarrow \mathcal{A}(X, A)$  is given by  $p \mapsto p \circ g$ .

A functor  $\mathcal{A}^{op} \rightarrow \mathbf{Set}$  that is isomorphic to  $H_A$  is called a representable.

# Required Results

If a transformation is natural in two individual variables simultaneously, then it is natural their pair.

## Lemma

Let  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$  be categories. Suppose there are functors  $F, G : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ .

For every  $A \in \mathcal{A}$ , there are functors,  $F^A, G^A : \mathcal{B} \rightarrow \mathcal{C}$  defined as taking  $B \in \mathcal{B}$  to  $F(A, B)$ ,  $G(A, B)$  and morphism  $f$  to  $F((1_A, f))$ ,  $G((1_A, f))$ . And, for every  $B \in \mathcal{B}$ , there are functors  $F_B, G_B : \mathcal{A} \rightarrow \mathcal{C}$  defined similarly.

A family of maps,  $(\alpha_{A,B} : F(A, B) \rightarrow G(A, B))_{A \in \mathcal{A}, B \in \mathcal{B}}$  is a natural transformation  $F \rightarrow G$  if the following conditions are satisfied:

- ① For each  $A \in \mathcal{A}$ , the family  $(\alpha_{A,B} : F^A(B) \rightarrow G^A(B))_{B \in \mathcal{B}}$  is a natural transformation  $F^A \rightarrow G^A$ ;
- ② For each  $B \in \mathcal{B}$ , the family  $(\alpha_{A,B} : F_B(A) \rightarrow G_B(A))_{A \in \mathcal{A}}$  is a natural transformation  $F_B \rightarrow G_B$ .

## Required Results

If a transformation is natural in two individual variables simultaneously, then it is natural their pair.

### Lemma

Let  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$  be categories. Suppose there are functors  $F, G : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ .

For every  $A \in \mathcal{A}$ , there are functors,  $F^A, G^A : \mathcal{B} \rightarrow \mathcal{C}$  defined as taking  $B \in \mathcal{B}$  to  $F(A, B)$ ,  $G(A, B)$  and morphism  $f$  to  $F((1_A, f))$ ,  $G((1_A, f))$ . And, for every  $B \in \mathcal{B}$ , there are functors  $F_B, G_B : \mathcal{A} \rightarrow \mathcal{C}$  defined similarly.

A family of maps,  $(\alpha_{A,B} : F(A, B) \rightarrow G(A, B))_{A \in \mathcal{A}, B \in \mathcal{B}}$  is a natural transformation  $F \rightarrow G$  if the following conditions are satisfied:

- ① For each  $A \in \mathcal{A}$ , the family  $(\alpha_{A,B} : F^A(B) \rightarrow G^A(B))_{B \in \mathcal{B}}$  is a natural transformation  $F^A \rightarrow G^A$ ;
- ② For each  $B \in \mathcal{B}$ , the family  $(\alpha_{A,B} : F_B(A) \rightarrow G_B(A))_{A \in \mathcal{A}}$  is a natural transformation  $F_B \rightarrow G_B$ .

Following is an equivalent definition of natural isomorphism.

### Lemma

Let  $\mathcal{A} \begin{array}{c} \xrightarrow{F} \\ \Downarrow \alpha \\ \xrightarrow{G} \end{array} \mathcal{B}$  be a natural transformation. If for every  $A \in \mathcal{A}$ ,  $\alpha_A : F(A) \rightarrow G(A)$  is an isomorphism then  $\alpha$  is a natural isomorphism.

# Yoneda Lemma

## Theorem

*If  $\mathcal{A}$  is a locally small category then, for any object  $A \in \mathcal{A}$  and  $X \in [\mathcal{A}^{op}, \text{Set}]$ , there exists an isomorphism,*

$$[\mathcal{A}^{op}, \text{Set}](H_A, X) \cong X(A) \text{ which is natural in } A \text{ and } X. \quad (1)$$



# Yoneda Lemma

## Theorem

*If  $\mathcal{A}$  is a locally small category then, for any object  $A \in \mathcal{A}$  and  $X \in [\mathcal{A}^{op}, Set]$ , there exists an isomorphism,*

$$[\mathcal{A}^{op}, Set](H_A, X) \cong X(A) \text{ which is natural in } A \text{ and } X. \quad (1)$$

Outline of the proof:

- Define a map from  $[\mathcal{A}^{op}, Set](H_A, X)$  to  $X(A)$ .
- Define a map from  $X(A)$  to  $[\mathcal{A}^{op}, Set](H_A, X)$ .

# Yoneda Lemma

## Theorem

*If  $\mathcal{A}$  is a locally small category then, for any object  $A \in \mathcal{A}$  and  $X \in [\mathcal{A}^{op}, Set]$ , there exists an isomorphism,*

$$[\mathcal{A}^{op}, Set](H_A, X) \cong X(A) \text{ which is natural in } A \text{ and } X. \quad (1)$$

Outline of the proof:

- Define a map from  $[\mathcal{A}^{op}, Set](H_A, X)$  to  $X(A)$ .
- Define a map from  $X(A)$  to  $[\mathcal{A}^{op}, Set](H_A, X)$ .
- Show the maps to be mutually inverse.

# Yoneda Lemma

## Theorem

*If  $\mathcal{A}$  is a locally small category then, for any object  $A \in \mathcal{A}$  and  $X \in [\mathcal{A}^{op}, \text{Set}]$ , there exists an isomorphism,*

$$[\mathcal{A}^{op}, \text{Set}](H_A, X) \cong X(A) \text{ which is natural in } A \text{ and } X. \quad (1)$$

Outline of the proof:

- Define a map from  $[\mathcal{A}^{op}, \text{Set}](H_A, X)$  to  $X(A)$ .
- Define a map from  $X(A)$  to  $[\mathcal{A}^{op}, \text{Set}](H_A, X)$ .
- Show the maps to be mutually inverse.
- Prove the resulting isomorphism natural in  $A$ .

# Yoneda Lemma

## Theorem

*If  $\mathcal{A}$  is a locally small category then, for any object  $A \in \mathcal{A}$  and  $X \in [\mathcal{A}^{op}, \text{Set}]$ , there exists an isomorphism,*

$$[\mathcal{A}^{op}, \text{Set}](H_A, X) \cong X(A) \text{ which is natural in } A \text{ and } X. \quad (1)$$

Outline of the proof:

- Define a map from  $[\mathcal{A}^{op}, \text{Set}](H_A, X)$  to  $X(A)$ .
- Define a map from  $X(A)$  to  $[\mathcal{A}^{op}, \text{Set}](H_A, X)$ .
- Show the maps to be mutually inverse.
- Prove the resulting isomorphism natural in  $A$ .
- Prove it natural in  $X$ .

# Cayley's Theorem

## Theorem

**Cayley's Theorem** *Every group,  $G$  is isomorphic to a subgroup of symmetric group on  $G$ .*

Outline of the proof: Take a group  $G$ .

- Define category  $\mathcal{A}$  to have a single object,  $\star$ . With morphisms corresponding to elements of  $G$ . With their product also being as elements of  $G$ . So that  $\mathcal{A}(\star, \star)$  is isomorphic to group  $G$ .

# Cayley's Theorem

## Theorem

**Cayley's Theorem** *Every group,  $G$  is isomorphic to a subgroup of symmetric group on  $G$ .*

Outline of the proof: Take a group  $G$ .

- Define category  $\mathcal{A}$  to have a single object,  $\star$ . With morphisms corresponding to elements of  $G$ . With their product also being as elements of  $G$ . So that  $\mathcal{A}(\star, \star)$  is isomorphic to group  $G$ .
- Prove that each member of the collection  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  can be considered as a bijection on  $G$ .

# Cayley's Theorem

## Theorem

**Cayley's Theorem** *Every group,  $G$  is isomorphic to a subgroup of symmetric group on  $G$ .*

Outline of the proof: Take a group  $G$ .

- Define category  $\mathcal{A}$  to have a single object,  $\star$ . With morphisms corresponding to elements of  $G$ . With their product also being as elements of  $G$ . So that  $\mathcal{A}(\star, \star)$  is isomorphic to group  $G$ .
- Prove that each member of the collection  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  can be considered as a bijection on  $G$ .
- Show that the above collection of bijections is actually a group.

# Cayley's Theorem

## Theorem

**Cayley's Theorem** *Every group,  $G$  is isomorphic to a subgroup of symmetric group on  $G$ .*

Outline of the proof: Take a group  $G$ .

- Define category  $\mathcal{A}$  to have a single object,  $\star$ . With morphisms corresponding to elements of  $G$ . With their product also being as elements of  $G$ . So that  $\mathcal{A}(\star, \star)$  is isomorphic to group  $G$ .
- Prove that each member of the collection  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  can be considered as a bijection on  $G$ .
- Show that the above collection of bijections is actually a group.
- Use Yoneda Lemma to get a set isomorphism between  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  and  $\mathcal{A}(\star, \star)$ .



# Cayley's Theorem

## Theorem

**Cayley's Theorem** *Every group,  $G$  is isomorphic to a subgroup of symmetric group on  $G$ .*

Outline of the proof: Take a group  $G$ .

- Define category  $\mathcal{A}$  to have a single object,  $\star$ . With morphisms corresponding to elements of  $G$ . With their product also being as elements of  $G$ . So that  $\mathcal{A}(\star, \star)$  is isomorphic to group  $G$ .
- Prove that each member of the collection  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  can be considered as a bijection on  $G$ .
- Show that the above collection of bijections is actually a group.
- Use Yoneda Lemma to get a set isomorphism between  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  and  $\mathcal{A}(\star, \star)$ .
- Prove this set isomorphism to be a group isomorphism.

# Yoneda Embedding

## Definition

A category  $\mathcal{A}$  is said to be embedded in a category  $\mathcal{B}$  if and only if there exists a functor  $F : \mathcal{A} \rightarrow \mathcal{B}$  such that  $F$  is full and faithful.

# Yoneda Embedding

## Definition

A category  $\mathcal{A}$  is said to be embedded in a category  $\mathcal{B}$  if and only if there exists a functor  $F : \mathcal{A} \rightarrow \mathcal{B}$  such that  $F$  is full and faithful.

## Proposition

For a locally small category  $\mathcal{A}$ , fixing an object  $A \in \mathcal{A}$  gives a functor,  $H_{\bullet} : \mathcal{A} \rightarrow [\mathcal{A}^{op}, \text{Set}]$  defined as:

- 1 For any object  $B \in \mathcal{A}$ ,  $H_{\bullet}(B) := H_A(B)$ ,
- 2 For any morphism,  $g : X \rightarrow Y$  in  $\mathcal{A}$ , for any  $K \in \mathcal{A}$ ,  $H_{\bullet}(g)$  takes  $p \in H_X(K)$  to  $g \circ p$  in  $H_Y(K)$ .

# Yoneda Embedding

## Definition

A category  $\mathcal{A}$  is said to be embedded in a category  $\mathcal{B}$  if and only if there exists a functor  $F : \mathcal{A} \rightarrow \mathcal{B}$  such that  $F$  is full and faithful.

## Proposition

For a locally small category  $\mathcal{A}$ , fixing an object  $A \in \mathcal{A}$  gives a functor,  $H_\bullet : \mathcal{A} \rightarrow [\mathcal{A}^{op}, \text{Set}]$  defined as:

- ① For any object  $B \in \mathcal{A}$ ,  $H_\bullet(B) := H_A(B)$ ,
- ② For any morphism,  $g : X \rightarrow Y$  in  $\mathcal{A}$ , for any  $K \in \mathcal{A}$ ,  $H_\bullet(g)$  takes  $p \in H_X(K)$  to  $g \circ p$  in  $H_Y(K)$ .

## Theorem

**Yoneda Embedding** Any locally small category can be embedded in the presheaf category on it.

# Yoneda Embedding

## Definition

A category  $\mathcal{A}$  is said to be embedded in a category  $\mathcal{B}$  if and only if there exists a functor  $F : \mathcal{A} \rightarrow \mathcal{B}$  such that  $F$  is full and faithful.

## Proposition

For a locally small category  $\mathcal{A}$ , fixing an object  $A \in \mathcal{A}$  gives a functor,  $H_\bullet : \mathcal{A} \rightarrow [\mathcal{A}^{op}, \text{Set}]$  defined as:

- ① For any object  $B \in \mathcal{A}$ ,  $H_\bullet(B) := H_A(B)$ ,
- ② For any morphism,  $g : X \rightarrow Y$  in  $\mathcal{A}$ , for any  $K \in \mathcal{A}$ ,  $H_\bullet(g)$  takes  $p \in H_X(K)$  to  $g \circ p$  in  $H_Y(K)$ .

## Theorem

**Yoneda Embedding** Any locally small category can be embedded in the presheaf category on it.

The proof proceeds by showing functor  $H_\bullet$  to be full and faithful.

# Prorelation

## Definition

A prorelation is a partially ordered, down-directed, up-set of relations  $X \rightarrow Y$ . That is,  $P \subseteq \mathcal{P}(X \times Y)$  is a prorelation if it satisfies the following conditions:

- 1 Partial Order: Containment of relations defines a partial order. That is,  $r \subseteq s$  meaning that for any  $(x, y) \in X \times Y$ , if  $(x, y) \in r$  then  $(x, y) \in s$ .
- 2 Down-directed: For any  $r, s \in P$ , there exists  $t \in P$  such that  $t \subseteq r$  and  $t \subseteq s$ .
- 3 Up-set: For any relation  $u : X \rightarrow Y$ , if there exists  $p \in P$  such that  $p \subseteq u$  then  $u \in P$ .

# Prorelation

## Definition

A prorelation is a partially ordered, down-directed, up-set of relations  $X \rightarrow Y$ . That is,  $P \subseteq \mathcal{P}(X \times Y)$  is a prorelation if it satisfies the following conditions:

- 1 Partial Order: Containment of relations defines a partial order. That is,  $r \subseteq s$  meaning that for any  $(x, y) \in X \times Y$ , if  $(x, y) \in r$  then  $(x, y) \in s$ .
- 2 Down-directed: For any  $r, s \in P$ , there exists  $t \in P$  such that  $t \subseteq r$  and  $t \subseteq s$ .
- 3 Up-set: For any relation  $u : X \rightarrow Y$ , if there exists  $p \in P$  such that  $p \subseteq u$  then  $u \in P$ .

## Example

For any positive real number  $\epsilon$ , define a relation on  $\mathbb{R}$  as  $A_\epsilon = \{(x, y) \mid |x - y| < \epsilon\}$ . The collection of all relations on  $\mathbb{R}$  that contains some  $A_\epsilon$  will be a prorelation,  $K$  on  $\mathbb{R}$ . That is,  $K = \{a : \mathbb{R} \rightarrow \mathbb{R} \mid a \supseteq A_\epsilon \text{ for some } \epsilon > 0\}$  forms a prorelation. If  $k, l \in K$ , then there exist  $\delta, \epsilon > 0$  such that  $k \supseteq A_\delta$  and  $l \supseteq A_\epsilon$ . Thus, the relation  $A_{\frac{\delta+\epsilon}{2}}$  is in both  $k$  and  $l$ . Moreover,  $K$  is an up-set by definition.

# Prorelation

## Definition

A prorelation is a partially ordered, down-directed, up-set of relations  $X \rightarrow Y$ . That is,  $P \subseteq \mathcal{P}(X \times Y)$  is a prorelation if it satisfies the following conditions:

- ① Partial Order: Containment of relations defines a partial order. That is,  $r \subseteq s$  meaning that for any  $(x, y) \in X \times Y$ , if  $(x, y) \in r$  then  $(x, y) \in s$ .
- ② Down-directed: For any  $r, s \in P$ , there exists  $t \in P$  such that  $t \subseteq r$  and  $t \subseteq s$ .
- ③ Up-set: For any relation  $u : X \rightarrow Y$ , if there exists  $p \in P$  such that  $p \subseteq u$  then  $u \in P$ .

## Example

For any positive real number  $\epsilon$ , define a relation on  $\mathbb{R}$  as  $A_\epsilon = \{(x, y) \mid |x - y| < \epsilon\}$ . The collection of all relations on  $\mathbb{R}$  that contains some  $A_\epsilon$  will be a prorelation,  $K$  on  $\mathbb{R}$ . That is,  $K = \{a : \mathbb{R} \rightarrow \mathbb{R} \mid a \supseteq A_\epsilon \text{ for some } \epsilon > 0\}$  forms a prorelation. If  $k, l \in K$ , then there exist  $\delta, \epsilon > 0$  such that  $k \supseteq A_\delta$  and  $l \supseteq A_\epsilon$ . Thus, the relation  $A_{\frac{\delta+\epsilon}{2}}$  is in both  $k$  and  $l$ . Moreover,  $K$  is an up-set by definition.

## Lemma

*Composition of two prorelations is a prorelation.*



# Quasi-Uniform Space

Quasi-uniformity is a particular kind of prorelation.

## Definition

A prorelation  $P$  on a set  $X$  is said to be a quasi-uniformity if it satisfies the following conditions:

- 1 Every relation in  $P$  is reflexive. That is, for each  $p \in P$ , if  $x \in X$  then  $(x, x) \in p$ .
- 2 For each  $p$  in  $P$ , there exists  $p'$  in  $P$  such that  $p' \circ p' \subseteq p$ .

# Quasi-Uniform Space

Quasi-uniformity is a particular kind of prerelation.

## Definition

A prerelation  $P$  on a set  $X$  is said to be a quasi-uniformity if it satisfies the following conditions:

- ① Every relation in  $P$  is reflexive. That is, for each  $p \in P$ , if  $x \in X$  then  $(x, x) \in p$ .
- ② For each  $p$  in  $P$ , there exists  $p'$  in  $P$  such that  $p' \circ p' \subseteq p$ .

## Example

As earlier, define  $A_\epsilon = \{(x, y) \mid |x - y| < \epsilon\}$ . The collection of all relations on  $\mathbb{R}$  that contains some  $A_\epsilon$ , that is,  $K = \{a : \mathbb{R} \rightarrow \mathbb{R} \mid a \supseteq A_\epsilon \text{ for some } \epsilon > 0\}$  forms a prerelation. If  $k, l \in K$ , then there exist  $\delta, \epsilon > 0$  such that  $k \supseteq A_\delta$  and  $l \supseteq A_\epsilon$ . Thus, the relation  $A_{\frac{\delta+\epsilon}{2}}$  is in both  $k$  and  $l$ . Moreover,  $K$  is an up-set by definition.

# Quasi-Uniform Space

Quasi-uniformity is a particular kind of prorelation.

## Definition

A prorelation  $P$  on a set  $X$  is said to be a quasi-uniformity if it satisfies the following conditions:

- ① Every relation in  $P$  is reflexive. That is, for each  $p \in P$ , if  $x \in X$  then  $(x, x) \in p$ .
- ② For each  $p$  in  $P$ , there exists  $p'$  in  $P$  such that  $p' \circ p' \subseteq p$ .

## Example

As earlier, define  $A_\epsilon = \{(x, y) \mid |x - y| < \epsilon\}$ . The collection of all relations on  $\mathbb{R}$  that contains some  $A_\epsilon$ , that is,  $K = \{a : \mathbb{R} \rightarrow \mathbb{R} \mid a \supseteq A_\epsilon \text{ for some } \epsilon > 0\}$  forms a prorelation. If  $k, l \in K$ , then there exist  $\delta, \epsilon > 0$  such that  $k \supseteq A_\delta$  and  $l \supseteq A_\epsilon$ . Thus, the relation  $A_{\frac{\delta+\epsilon}{2}}$  is in both  $k$  and  $l$ . Moreover,  $K$  is an up-set by definition.

## Definition

If  $X$  is a set, and  $A$  is a quasi-uniformity on  $X$ , then  $(X, A)$  is a quasi-uniform space.

The following gives us a partial order on prorelations:

### Definition

For prorelations  $P, Q : X \rightarrow Y$ , if for each  $q \in Q$ , there exists  $p \in P$  such that  $p \subseteq q$ , then we write  $P \leq Q$ .

The following gives us a partial order on prorelations:

### Definition

For prorelations  $P, Q : X \rightarrow Y$ , if for each  $q \in Q$ , there exists  $p \in P$  such that  $p \subseteq q$ , then we write  $P \leq Q$ .

We will require the following definition in the next slide.

### Definition

A function,  $f : (X, A) \rightarrow (Y, B)$  is said to be uniformly continuous if and only if  $f.A \leq B.f$ . That is, for each  $b \in B$ , there exists  $a \in A$  such that  $f \circ a \subseteq b \circ f$ .

# Category definition

On this slide, we define two categories.

## Definition

The collection of quasi-uniform spaces can be given a categorical structure by considering the uniformly continuous maps between two spaces as the morphisms between them. We call this category QUnif. Composition is as that of functions, and identity morphisms are the identity functions.

# Category definition

On this slide, we define two categories.

## Definition

The collection of quasi-uniform spaces can be given a categorical structure by considering the uniformly continuous maps between two spaces as the morphisms between them. We call this category QUnif. Composition is as that of functions, and identity morphisms are the identity functions.

## Definition

A prorelation,  $\phi : X \rightarrow Y$  is called a promodule  $\phi : (X, A) \rightarrow (Y, B)$  if it satisfies:

$$\phi.A \leq \phi \text{ and } B.\phi \leq \phi.$$

# Category definition

On this slide, we define two categories.

## Definition

The collection of quasi-uniform spaces can be given a categorical structure by considering the uniformly continuous maps between two spaces as the morphisms between them. We call this category QUnif. Composition is as that of functions, and identity morphisms are the identity functions.

## Definition

A prerelation,  $\phi : X \rightarrow Y$  is called a promodule  $\phi : (X, A) \rightarrow (Y, B)$  if it satisfies:

$$\phi.A \leq \phi \text{ and } B.\phi \leq \phi.$$

## Definition

Now, we define a 2-category called ProMod as having quasi-uniform spaces as 0-cells and the promodules between them being 1-cells. The promodule  $A$  will work as the identity of  $(X, A)$ .

Let promodules  $P, Q : (X, A) \rightarrow (Y, B)$ . Then, there is a 2-cell from  $P$  to  $Q$  if and only if  $P \leq Q$  as prerelations.



# Functors from QUnif to ProMod

We have a covariant functor from QUnif to ProMod:

## Proposition

*Functor  $(-)_* : QUnif \rightarrow ProMod$  defined as:*

- ① *for  $(X, A) \in QUnif$ ,  $(X, A)_* := (X, A) \in ProMod$ ,*
- ② *for  $f : (X, A) \rightarrow (Y, B)$  in  $QUnif$ ,  $f_* := B.f$ ,*

# Functors from QUnif to ProMod

We have a covariant functor from QUnif to ProMod:

## Proposition

Functor  $(-)_* : QUnif \rightarrow ProMod$  defined as:

- ① for  $(X, A) \in QUnif$ ,  $(X, A)_* := (X, A) \in ProMod$ ,
- ② for  $f : (X, A) \rightarrow (Y, B)$  in  $QUnif$ ,  $f_* := B.f$ ,

as well as a contravariant functor from QUnif to ProMod,

## Proposition

Functor  $(-)^* : QUnif^{op} \rightarrow ProMod$  defined as as:

- ① for  $(X, A) \in QUnif$ ,  $(X, A)^* := (X, A) \in ProMod$ ,
- ② for  $f : (X, A) \rightarrow (Y, B)$  in  $QUnif$ ,  $f^* := f \circ B$ .

# Topological definitions

## Definition

For any quasi-uniform space  $(X, A)$ , an element  $x \in X$  is said to belong in the topological closure of set  $M \subseteq X$  if and only if for each  $a \in A$ , there exists  $y \in M$  such that  $x a y$  and  $y a x$ .

# Topological definitions

## Definition

For any quasi-uniform space  $(X, A)$ , an element  $x \in X$  is said to belong in the topological closure of set  $M \subseteq X$  if and only if for each  $a \in A$ , there exists  $y \in M$  such that  $x a y$  and  $y a x$ .

## Definition

Let  $f : (X, A) \rightarrow (Y, B)$  be a uniformly continuous function.

- ①  $f$  is said to be fully faithful if and only if  $f^*.f_* = A$ .
- ②  $f$  is said to be fully dense if and only if  $f_*.f^* = B$ .
- ③  $f$  is said to be topologically dense if and only if  $\overline{f(X)} = Y$ .

# Topological definitions

## Definition

For any quasi-uniform space  $(X, A)$ , an element  $x \in X$  is said to belong in the topological closure of set  $M \subseteq X$  if and only if for each  $a \in A$ , there exists  $y \in M$  such that  $x a y$  and  $y a x$ .

## Definition

Let  $f : (X, A) \rightarrow (Y, B)$  be a uniformly continuous function.

- ①  $f$  is said to be fully faithful if and only if  $f^*.f_* = A$ .
- ②  $f$  is said to be fully dense if and only if  $f_*.f^* = B$ .
- ③  $f$  is said to be topologically dense of and only if  $\overline{f(X)} = Y$ .

The following proposition gives us easier to apply versions of previous definitions.

## Proposition

Let  $f : (X, A) \rightarrow (Y, B)$  be a uniformly continuous map.

- ①  $f$  is fully faithful if and only if  $A = f^\circ.B.f$ , that is  $A \geq f^\circ.B.f$ .
- ②  $f$  is fully dense if and only if for any  $b \in B$ ,  $\exists b' \in B$  such that  $b' \subseteq b f f^\circ b$ .
- ③  $f$  is topologically dense if and only if for any  $b \in B$ ,  $b f f^\circ b$  is reflexive.
- ④  $f$  is fully dense if and only if  $f$  is topologically dense.

## Quasi-uniform space of promodules

### Definition

The set  $PX$  is defined to be the collection of all promodules from the quasi-uniform space  $(X, A)$  to the quasi-uniform space  $1$ .

$$PX := \{\psi : (X, A) \rightarrow 1 \mid \psi \text{ is a promodule}\}$$

# Quasi-uniform space of promodules

## Definition

The set  $PX$  is defined to be the collection of all promodules from the quasi-uniform space  $(X, A)$  to the quasi-uniform space  $1$ .

$$PX := \{\psi : (X, A) \rightarrow 1 \mid \psi \text{ is a promodule}\}$$

On this set, we can define a quasi-uniformity.

## Proposition

For any  $a \in A$ ,  $\tilde{a}$  is defined to be a relation on  $PX$  as:

$$\text{for } \phi, \psi \in PX, \phi \tilde{a} \psi \text{ if and only if } \phi \leq \psi.a .$$

The set,  $\tilde{A} := \{\tilde{a} : a \in A\}$  defines a quasi-uniformity on  $PX$ .

# Quasi-uniform space of promodules

## Definition

The set  $PX$  is defined to be the collection of all promodules from the quasi-uniform space  $(X, A)$  to the quasi-uniform space  $1$ .

$$PX := \{\psi : (X, A) \rightarrow 1 \mid \psi \text{ is a promodule}\}$$

On this set, we can define a quasi-uniformity.

## Proposition

For any  $a \in A$ ,  $\tilde{a}$  is defined to be a relation on  $PX$  as:

$$\text{for } \phi, \psi \in PX, \phi \tilde{a} \psi \text{ if and only if } \phi \leq \psi.a.$$

The set,  $\tilde{A} := \{\tilde{a} : a \in A\}$  defines a quasi-uniformity on  $PX$ .

Just like we could embed any category into its presheaf category via  $H_\bullet$ , we can embed any quasi-uniform space into its 'quasi-uniform space of promodules':

## Proposition

For a quasi-uniform space  $(X, A)$ , function  $y_X : X \rightarrow PX$  is defined by  $x \mapsto x^*$  for  $x \in X$ .

- $y_X : (X, A) \rightarrow (PX, \tilde{A})$  is a uniformly continuous map.
- $y_X : (X, A) \rightarrow (PX, \tilde{A})$  is fully faithful.



## Yoneda Lemma for Quasi-Uniform Spaces

To think of Yoneda Lemma in quasi-uniform spaces,

- consider the promodule  $(y_X)_*$  to be the representable  $H_A$  in our initial statement of Yoneda Lemma,
- consider the promodule  $\psi^*$  to be the presheaf  $X$  in our initial statement of Yoneda Lemma.

This gives us a weak version of Yoneda Lemma that holds in quasi-uniform space:

# Yoneda Lemma for Quasi-Uniform Spaces

To think of Yoneda Lemma in quasi-uniform spaces,

- consider the promodule  $(y_X)_*$  to be the representable  $H_A$  in our initial statement of Yoneda Lemma,
- consider the promodule  $\psi^*$  to be the presheaf  $X$  in our initial statement of Yoneda Lemma.

This gives us a weak version of Yoneda Lemma that holds in quasi-uniform space:

## Theorem

*The following statements hold for any  $\psi \in PX$ :*

- ①  $\psi \geq \psi^* \cdot (y_X)_*$ ,
- ②  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^* \cdot (y_X)_*$ .

# Yoneda Lemma for Quasi-Uniform Spaces

To think of Yoneda Lemma in quasi-uniform spaces,

- consider the promodule  $(y_X)_*$  to be the representable  $H_A$  in our initial statement of Yoneda Lemma,
- consider the promodule  $\psi^*$  to be the presheaf  $X$  in our initial statement of Yoneda Lemma.

This gives us a weak version of Yoneda Lemma that holds in quasi-uniform space:

## Theorem

*The following statements hold for any  $\psi \in PX$ :*

- ①  $\psi \geq \psi^* \cdot (y_X)_*$ ,
- ②  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^* \cdot (y_X)_*$ .

And the full strength Yoneda Lemma holds only for elements of  $PX$  that are also in the topological closure of  $y_X(X)$ :

## Corollary

*For  $\psi \in PX$ ,  $\psi \in \overline{y_X(X)}$  if and only if  $\psi$  is a right-adjoint.*

# References



Maria Manuel Clementino and Dirk Hofmann. “On the completion monad via the Yoneda embedding in quasi-uniform spaces”. In: *Topology and its Applications* 158.17 (Nov. 2011), p. 24232430. ISSN: 01668641. DOI: 10.1016/j.topol.2011.01.026.



Peter Fletcher and William F. Lindgren. *Quasi-uniform spaces*. Lecture notes in pure and applied mathematics. M. Dekker, 1982. ISBN: 9780824718398.



Tom Leinster. *Basic category theory*. Cambridge studies in advanced mathematics. Cambridge University Press, 2014. ISBN: 9781107044241.



Emily Riehl. *Category theory in context*. Aurora: Dover modern math originals. Dover Publications, 2016. ISBN: 9780486809038.

# Showing $\psi \geq \psi^*.(y_X)^*$

- By definition,  $(y_X)_* = \tilde{A}.y_X$ , and  $\psi^* = \psi^\circ.\tilde{A}$ ,  
so, we need to show that  $\psi \geq (y_X)_*.\psi^* = \psi^\circ.\tilde{A}.\tilde{A}.y_X$ .  
As  $\tilde{A}.\tilde{A} = \tilde{A}$ , we need  $\psi \geq \psi^\circ.\tilde{A}.y_X$ .  
Fix  $p \in \psi$ . By definition of 2-cells in ProMod, we require the following.
- For this  $p$  we need to find  $a \in A$  such that  $p \supseteq \psi^\circ \tilde{a} y_X$ .

# Showing $\psi \geq \psi^*.(y_X)^*$

- By definition,  $(y_X)_* = \tilde{A}.y_X$ , and  $\psi^* = \psi^\circ.\tilde{A}$ ,  
so, we need to show that  $\psi \geq (y_X)_*.\psi^* = \psi^\circ.\tilde{A}.\tilde{A}.y_X$ .  
As  $\tilde{A}.\tilde{A} = \tilde{A}$ , we need  $\psi \geq \psi^\circ.\tilde{A}.y_X$ .  
Fix  $p \in \psi$ . By definition of 2-cells in ProMod, we require the following.
- For this  $p$  we need to find  $a \in A$  such that  $p \supseteq \psi^\circ \tilde{a} y_X$ .
- Examining the right side of the condition, for any  $a \in A$ ,  $x \in X$

$$(\psi^\circ.\tilde{a}.y_X)(x) == \psi^\circ(\tilde{a}(x^*)) = \begin{cases} \phi & \text{if } \psi \notin \tilde{a}(x^*) \\ \star & \text{if } \psi \in \tilde{a}(x^*) \end{cases}.$$

So, if  $\psi \notin \tilde{a}(x^*)$ , then, as  $p \supseteq \phi$ , we're done. Thus, let  $\psi \in \tilde{a}(x^*)$ , that is,  $x^* \tilde{a} \psi$ .

- Hence,  $x(\psi^\circ.\tilde{a}.y_X)\star$ , we now need to show  $x p \star$

# Showing $\psi \geq \psi^*.(y_X)^*$

- By definition,  $(y_X)_* = \tilde{A}.y_X$ , and  $\psi^* = \psi^\circ.\tilde{A}$ ,  
so, we need to show that  $\psi \geq (y_X)_*.\psi^* = \psi^\circ.\tilde{A}.\tilde{A}.y_X$ .  
As  $\tilde{A}.\tilde{A} = \tilde{A}$ , we need  $\psi \geq \psi^\circ.\tilde{A}.y_X$ .  
Fix  $p \in \psi$ . By definition of 2-cells in ProMod, we require the following.
- For this  $p$  we need to find  $a \in A$  such that  $p \supseteq \psi^\circ \tilde{a} y_X$ .
- Examining the right side of the condition, for any  $a \in A$ ,  $x \in X$

$$(\psi^\circ.\tilde{a}.y_X)(x) == \psi^\circ(\tilde{a}(x^*)) = \begin{cases} \phi & \text{if } \psi \notin \tilde{a}(x^*) \\ \star & \text{if } \psi \in \tilde{a}(x^*) \end{cases}.$$

So, if  $\psi \notin \tilde{a}(x^*)$ , then, as  $p \supseteq \phi$ , we're done. Thus, let  $\psi \in \tilde{a}(x^*)$ , that is,  $x^* \tilde{a} \psi$ .

- Hence,  $x(\psi^\circ.\tilde{a}.y_X)\star$ , we now need to show  $x p \star$
- As  $\psi$  is a promodule,  $\psi.A \leq \psi$  gives us the the existence of  $q \in \psi$  and  $a \in A$  such that  $qa \subseteq p$ . Using the definition of  $\tilde{a}$ ,  $x^* \tilde{a} \psi$  gives:

$$x^\circ.A \leq \psi.a \implies \exists b \in A : x^\circ b \subseteq qa \implies \forall z \in X, (x^\circ b)(z) \subseteq (qa)(z).$$

# Showing $\psi \geq \psi^*.(y_X)^*$

- By definition,  $(y_X)_* = \tilde{A}.y_X$ , and  $\psi^* = \psi^\circ.\tilde{A}$ ,  
so, we need to show that  $\psi \geq (y_X)_*.\psi^* = \psi^\circ.\tilde{A}.\tilde{A}.y_X$ .  
As  $\tilde{A}.\tilde{A} = \tilde{A}$ , we need  $\psi \geq \psi^\circ.\tilde{A}.y_X$ .  
Fix  $p \in \psi$ . By definition of 2-cells in ProMod, we require the following.
- For this  $p$  we need to find  $a \in A$  such that  $p \supseteq \psi^\circ \tilde{a} y_X$ .
- Examining the right side of the condition, for any  $a \in A$ ,  $x \in X$

$$(\psi^\circ.\tilde{a}.y_X)(x) = \psi^\circ(\tilde{a}(x^*)) = \begin{cases} \phi & \text{if } \psi \notin \tilde{a}(x^*) \\ \star & \text{if } \psi \in \tilde{a}(x^*) \end{cases}.$$

So, if  $\psi \notin \tilde{a}(x^*)$ , then, as  $p \supseteq \phi$ , we're done. Thus, let  $\psi \in \tilde{a}(x^*)$ , that is,  $x^* \tilde{a} \psi$ .

- Hence,  $x(\psi^\circ.\tilde{a}.y_X)\star$ , we now need to show  $x p \star$
- As  $\psi$  is a promodule,  $\psi.A \leq \psi$  gives us the the existence of  $q \in \psi$  and  $a \in A$  such that  $q a \subseteq p$ . Using the definition of  $\tilde{a}$ ,  $x^* \tilde{a} \psi$  gives:

$$x^\circ.A \leq \psi.a \implies \exists b \in A : x^\circ b \subseteq q a \implies \forall z \in X, (x^\circ b)(z) \subseteq (q a)(z).$$

- By quasi-uniformity of  $A$ ,  $b$  is reflexive, giving that in particular, for  $x b x$ ,

$$(x^\circ b)(x) \subseteq (q a)(x) \implies x^\circ x \subseteq (q a)(x) \implies \star \in (q a)(x) \implies \star \in p(x)$$



Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- 1 By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- 2 So, need that  $xp^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $xp^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b}y_X$ . Thus, for any  $z$  such that  $zcw$ ,  

$$(y_X c)(z) \subseteq (\tilde{b}y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*) \text{ that is, } z^* \tilde{b} w^* .$$

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $xp^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b}y_X$ . Thus, for any  $z$  such that  $zcw$ ,  

$$(y_X c)(z) \subseteq (\tilde{b}y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*) \text{ that is, } z^* \tilde{b} w^* .$$
- ④ As  $A$  is a quasi-uniformity,  $\exists d \in A : dd \subseteq c$ . Also, because  $A$  is a down-directed set,  $\exists a' \in A : a' \subseteq b, d$ . Thus, using the above chain of implications, for any  $x, y \in X$ ,  $x(a'a')y \implies x(dd)y \implies xcy \implies x^* \tilde{b} y^* .$

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $xp \star \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b}y_X$ . Thus, for any  $z$  such that  $zcw$ ,  

$$(y_X c)(z) \subseteq (\tilde{b}y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*) \text{ that is, } z^* \tilde{b} w^* .$$
- ④ As  $A$  is a quasi-uniformity,  $\exists d \in A : dd \subseteq c$ . Also, because  $A$  is a down-directed set,  $\exists a' \in A : a' \subseteq b, d$ . Thus, using the above chain of implications, for any  $x, y \in X$ ,  $x(a'a')y \implies x(dd)y \implies xcy \implies x^* \tilde{b} y^* .$
- ⑤ Now, because  $\psi \in \overline{y_X(X)}$ , we get  $\exists x \in X$  such that  $\psi \tilde{a}' x^*$  and  $x^* \tilde{a}' \psi$ .

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $xp^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b}y_X$ . Thus, for any  $z$  such that  $zcw$ ,  
 $(y_X c)(z) \subseteq (\tilde{b}y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*)$  that is,  $z^*\tilde{b}w^*$ .
- ④ As  $A$  is a quasi-uniformity,  $\exists d \in A : dd \subseteq c$ . Also, because  $A$  is a down-directed set,  $\exists a' \in A : a' \subseteq b, d$ . Thus, using the above chain of implications, for any  $x, y \in X$ ,  $x(a'a')y \implies x(dd)y \implies xcy \implies x^*\tilde{b}y^*$ .
- ⑤ Now, because  $\psi \in \overline{y_X(X)}$ , we get  $\exists x \in X$  such that  $\psi\tilde{a}'x^*$  and  $x^*\tilde{a}'\psi$ .
- ⑥ Using the definition of  $\tilde{a}'$ , from  $\psi\tilde{a}'x^*$ , we have  $\psi \leq x^o.A.a'$ , i.e.  
 $\exists p \in \psi : p \subseteq x^o a' a'$ .

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*. (y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o. \tilde{a}. y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o. \tilde{a}. y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $x p^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b} y_X$ . Thus, for any  $z$  such that  $z c w$ ,  
 $(y_X c)(z) \subseteq (\tilde{b} y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*)$  that is,  $z^* \tilde{b} w^*$ .
- ④ As  $A$  is a quasi-uniformity,  $\exists d \in A : dd \subseteq c$ . Also, because  $A$  is a down-directed set,  $\exists a' \in A : a' \subseteq b, d$ . Thus, using the above chain of implications, for any  $x, y \in X$ ,  $x(a' a') y \implies x(dd) y \implies x c y \implies x^* \tilde{b} y^*$ .
- ⑤ Now, because  $\psi \in \overline{y_X(X)}$ , we get  $\exists x \in X$  such that  $\psi \tilde{a}' x^*$  and  $x^* \tilde{a}' \psi$ .
- ⑥ Using the definition of  $\tilde{a}'$ , from  $\psi \tilde{a}' x^*$ , we have  $\psi \leq x^o.A.a'$ , i.e.  
 $\exists p \in \psi : p \subseteq x^o a' a'$ .
- ⑦ Fix any  $z \in X$  such that  $z p^*$ . Using (6), definition of  $x^o$  and then (4), we get that we need to show:

$$z p^* \implies z(x^o a' a')^* \implies z(a' a') x \implies z^* \tilde{b} x^* .$$

Showing  $\psi \in \overline{y_X(X)} \implies \psi \leq \psi^*.(y_X)^*$

- ① By definition, we need to show that for any  $a \in A$ ,  $\exists p \in \psi : p \subseteq \psi^o.\tilde{a}.y_X$ .  
Fix any  $x \in \text{Dom}(p)$ , will show  $p(x) \subseteq \psi^o.\tilde{a}.y_X(x) = \psi^o(\tilde{a}(x^*))$
- ② So, need that  $xp^* \implies \psi \in \tilde{a}(x^*)$  Fix any  $a \in A$ , let's find  $p \in \psi$  such that the above inequality holds.
- ③ Quasi-uniformity of  $A$  gives that  $\exists b \in A : bb \subseteq a$ . As we know that  $y_X$  is uniformly continuous, from Yoneda Embedding in QUS, we have that  $y_X.A \leq \tilde{A}.y_X$ . So,  $\exists c \in A$  such that  $y_X c \subseteq \tilde{b}y_X$ . Thus, for any  $z$  such that  $zcw$ ,  
 $(y_X c)(z) \subseteq (\tilde{b}y_X)(z) \implies y_X(c(z)) \subseteq \tilde{b}(z^*) \implies w^* \in \tilde{b}(z^*)$  that is,  $z^*\tilde{b}w^*$ .
- ④ As  $A$  is a quasi-uniformity,  $\exists d \in A : dd \subseteq c$ . Also, because  $A$  is a down-directed set,  $\exists a' \in A : a' \subseteq b, d$ . Thus, using the above chain of implications, for any  $x, y \in X$ ,  $x(a'a')y \implies x(dd)y \implies xcy \implies x^*\tilde{b}y^*$ .
- ⑤ Now, because  $\psi \in \overline{y_X(X)}$ , we get  $\exists x \in X$  such that  $\psi\tilde{a}'x^*$  and  $x^*\tilde{a}'\psi$ .
- ⑥ Using the definition of  $\tilde{a}'$ , from  $\psi\tilde{a}'x^*$ , we have  $\psi \leq x^o.A.a'$ , i.e.  
 $\exists p \in \psi : p \subseteq x^o.a'a'$ .
- ⑦ Fix any  $z \in X$  such that  $zp^*$ . Using (6), definition of  $x^o$  and then (4), we get that we need to show:

$$zp^* \implies z(x^o.a'a')^* \implies z(a'a')x \implies z^*\tilde{b}x^*.$$

- ⑧ Finally, as  $\tilde{a}' \subseteq \tilde{b}$ , From (5), we have that  $z^*\tilde{b}x^*$  and  $x^*\tilde{b}\psi$ . Using composition,  $z^*\tilde{b}x^*$ .



Showing  $y_X$  is uniformly continuous

- ① As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .

Showing  $y_X$  is uniformly continuous

- ① As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .
- ② So, need that for each  $a \in A$ , there exists  $b \in A$  such that  $y_X \circ b \subseteq \tilde{a} \circ y_X$ .

# Showing $y_X$ is uniformly continuous

- 1 As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .
- 2 So, need that for each  $a \in A$ , there exists  $b \in A$  such that  $y_X \circ b \subseteq \tilde{a} \circ y_X$ .
- 3 Applying the relations to some element,  $x$  of the set  $X$  gives us:

$$(y_X \circ b)(x) \subseteq (\tilde{a} \circ y_X)(x) \implies y_X(b(x)) \subseteq \tilde{a}(x^*).$$

# Showing $y_X$ is uniformly continuous

- ① As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .
- ② So, need that for each  $a \in A$ , there exists  $b \in A$  such that  $y_X \circ b \subseteq \tilde{a} \circ y_X$ .
- ③ Applying the relations to some element,  $x$  of the set  $X$  gives us:

$$(y_X \circ b)(x) \subseteq (\tilde{a} \circ y_X)(x) \implies y_X(b(x)) \subseteq \tilde{a}(x^*).$$

- ④ So, for the condition given above to hold, if  $y \in b(x)$ , then it's required that  $y^* = y_X(y) \in \tilde{a}(x^*)$  i.e.  $x^* \tilde{a} y^*$ . Using the definition of  $x^*, y^*$  and  $\tilde{a}$ ,

$$x^* \tilde{a} y^* \iff x^\circ.A \leq y^\circ.A.a \iff \forall a' \in A, \exists a'' \in A : x^\circ a'' \subseteq y^\circ a' a$$

# Showing $y_X$ is uniformly continuous

- ① As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .
- ② So, need that for each  $a \in A$ , there exists  $b \in A$  such that  $y_X \circ b \subseteq \tilde{a} \circ y_X$ .
- ③ Applying the relations to some element,  $x$  of the set  $X$  gives us:

$$(y_X \circ b)(x) \subseteq (\tilde{a} \circ y_X)(x) \implies y_X(b(x)) \subseteq \tilde{a}(x^*).$$

- ④ So, for the condition given above to hold, if  $y \in b(x)$ , then it's required that  $y^* = y_X(y) \in \tilde{a}(x^*)$  i.e.  $x^* \tilde{a} y^*$ . Using the definition of  $x^*, y^*$  and  $\tilde{a}$ ,

$$x^* \tilde{a} y^* \iff x^\circ.A \leq y^\circ.A.a \iff \forall a' \in A, \exists a'' \in A : x^\circ a'' \subseteq y^\circ a' a$$

- ⑤ Fix any  $a \in A, x \in X$ . Thus, quasi-uniformity of  $A$ , gives  $\exists a'' \in A$  such that  $a'' a'' \subseteq a$ . Now, choose  $y \in a''(x)$ . Hence, in order to show that the condition above holds, need that  $\forall b \in A, x^\circ a'' \subseteq y^\circ b a$ . Applying the relations to an element  $z \in X$  gives the following condition:

$$\forall b \in B, \forall x \in X, (x^\circ a'')(z) \subseteq (y^\circ b a)(z). \quad (2)$$

# Showing $y_X$ is uniformly continuous

- ① As  $y_X : (X, A) \rightarrow (PX, \tilde{A})$ , we need to show  $y_X.A \leq \tilde{A}.y_X$ .
- ② So, need that for each  $a \in A$ , there exists  $b \in A$  such that  $y_X \circ b \subseteq \tilde{a} \circ y_X$ .
- ③ Applying the relations to some element,  $x$  of the set  $X$  gives us:

$$(y_X \circ b)(x) \subseteq (\tilde{a} \circ y_X)(x) \implies y_X(b(x)) \subseteq \tilde{a}(x^*).$$

- ④ So, for the condition given above to hold, if  $y \in b(x)$ , then it's required that  $y^* = y_X(y) \in \tilde{a}(x^*)$  i.e.  $x^* \tilde{a} y^*$ . Using the definition of  $x^*, y^*$  and  $\tilde{a}$ ,

$$x^* \tilde{a} y^* \iff x^\circ.A \leq y^\circ.A.a \iff \forall a' \in A, \exists a'' \in A : x^\circ a'' \subseteq y^\circ a' a$$

- ⑤ Fix any  $a \in A, x \in X$ . Thus, quasi-uniformity of  $A$ , gives  $\exists a'' \in A$  such that  $a'' a'' \subseteq a$ . Now, choose  $y \in a''(x)$ . Hence, in order to show that the condition above holds, need that  $\forall b \in A, x^\circ a'' \subseteq y^\circ ba$ . Applying the relations to an element  $z \in X$  gives the following condition:

$$\forall b \in B, \forall x \in X, (x^\circ a'')(z) \subseteq (y^\circ ba)(z). \quad (2)$$

- ⑥ Examining the left side,

$$(x^\circ a'')(z) = x^\circ(a''(z)) = \begin{cases} \phi & \text{if } x \notin a''(z) \\ \star & \text{if } x \in a''(z) \end{cases}.$$

Showing  $y_X$  is uniformly continuous

④ Thus, to show that (1) holds, need to show that, for any  $b \in A$  and  $z \in X$ :

$$x \in a''(z) \implies z(y \circ ba) \star \text{ i.e. } y \in (ba)(z)$$

Showing  $y_X$  is uniformly continuous

- ⑦ Thus, to show that (1) holds, need to show that, for any  $b \in A$  and  $z \in X$ :

$$x \in a''(z) \implies z(y \circ ba) \star \text{ i.e. } y \in (ba)(z)$$

- ⑧ Fix any  $z \in X : x \in a''(z)$ . Also, by our choice of  $y$ , have that  $y \in a''(x)$ . Now, going to show that the above condition holds: As  $A$  is a quasi-uniformity,  $b$  is reflexive, giving that  $y \in b(y)$ .



# Showing $y_X$ is uniformly continuous

- ⑦ Thus, to show that (1) holds, need to show that, for any  $b \in A$  and  $z \in X$ :

$$x \in a''(z) \implies z(y \circ ba) \star \text{ i.e. } y \in (ba)(z)$$

- ⑧ Fix any  $z \in X : x \in a''(z)$ . Also, by our choice of  $y$ , have that  $y \in a''(x)$ . Now, going to show that the above condition holds: As  $A$  is a quasi-uniformity,  $b$  is reflexive, giving that  $y \in b(y)$ .
- ⑨ Hence, by composition of relations, we get:

$$za''x, xa''y \text{ and } yby \implies z(a''a''b)y \implies z(ab)y \text{ i.e. } y \in (ba)(z).$$

## Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^\circ \cdot \tilde{A} \cdot y_X$ .

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^\circ \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^\circ \tilde{b} y_X$ .

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^o \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^o \tilde{b} y_X$ .
- Applying the relations to an element,  $x \in X$  gives the condition:

$$(y_X^o \tilde{b} y_X)(x) \subseteq a(x) \implies (y_X^o \tilde{b})(x^*) = y_x^o(\tilde{b}(x^*)) \subseteq a(x). \quad (3)$$

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^o \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^o \tilde{b} y_X$ .
- Applying the relations to an element,  $x \in X$  gives the condition:

$$(y_X^o \tilde{b} y_X)(x) \subseteq a(x) \implies (y_X^o \tilde{b})(x^*) = y_X^o(\tilde{b}(x^*)) \subseteq a(x). \quad (3)$$

- Thus, if  $y^* \in PX$  such that  $x^* \tilde{b} y^*$ , then  $y \in y_X^o(\tilde{b}(x^*))$ . For the above condition to hold, need that  $y \in a(x)$ , that is,  $xay$ .

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^o \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^o \tilde{b} y_X$ .
- Applying the relations to an element,  $x \in X$  gives the condition:

$$(y_X^o \tilde{b} y_X)(x) \subseteq a(x) \implies (y_X^o \tilde{b})(x^*) = y_X^o(\tilde{b}(x^*)) \subseteq a(x). \quad (3)$$

- Thus, if  $y^* \in PX$  such that  $x^* \tilde{b} y^*$ , then  $y \in y_X^o(\tilde{b}(x^*))$ . For the above condition to hold, need that  $y \in a(x)$ , that is,  $xay$ .
- Thus, need only to show that for any  $a \in A$ , there exists  $b \in A$  such that for any  $x, y \in X$ ,  $x^* \tilde{b} y^*$  implies  $xay$ .

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^o \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^o \tilde{b} y_X$ .
- Applying the relations to an element,  $x \in X$  gives the condition:

$$(y_X^o \tilde{b} y_X)(x) \subseteq a(x) \implies (y_X^o \tilde{b})(x^*) = y_X^o(\tilde{b}(x^*)) \subseteq a(x). \quad (3)$$

- Thus, if  $y^* \in PX$  such that  $x^* \tilde{b} y^*$ , then  $y \in y_X^o(\tilde{b}(x^*))$ . For the above condition to hold, need that  $y \in a(x)$ , that is,  $xay$ .
- Thus, need only to show that for any  $a \in A$ , there exists  $b \in A$  such that for any  $x, y \in X$ ,  $x^* \tilde{b} y^*$  implies  $xay$ .
- So, fix  $a \in A$ , and take  $b \in A : bb \subseteq a$ . Now, let  $x^* \tilde{b} y^*$  i.e.  $x^o \cdot A \leq y^o \cdot A \cdot b$ .

# Showing $y_X$ is full and faithful

- By using the four-part proposition's first result, we just need to show that  $A \geq y_X^o \cdot \tilde{A} \cdot y_X$ .
- That amounts to showing  $\forall a \in A, \exists \tilde{b} \in \tilde{A} : a \supseteq y_X^o \tilde{b} y_X$ .
- Applying the relations to an element,  $x \in X$  gives the condition:

$$(y_X^o \tilde{b} y_X)(x) \subseteq a(x) \implies (y_X^o \tilde{b})(x^*) = y_X^o(\tilde{b}(x^*)) \subseteq a(x). \quad (3)$$

- Thus, if  $y^* \in PX$  such that  $x^* \tilde{b} y^*$ , then  $y \in y_X^o(\tilde{b}(x^*))$ . For the above condition to hold, need that  $y \in a(x)$ , that is,  $xay$ .
- Thus, need only to show that for any  $a \in A$ , there exists  $b \in A$  such that for any  $x, y \in X$ ,  $x^* \tilde{b} y^*$  implies  $xay$ .
- So, fix  $a \in A$ , and take  $b \in A : bb \subseteq a$ . Now, let  $x^* \tilde{b} y^*$  i.e.  $x^o \cdot A \leq y^o \cdot A \cdot b$ .
- Hence,  $\exists c \in A : x^o c \subseteq y^o bb$ . And as  $c$  is reflexive,

$$xcx \implies x(cx^o) \star \implies x(bby^o) \star \implies x(bb)y \implies xay.$$



# Proof: Yoneda Corollary I

Fix any  $\psi \in PX$ .

① ( $\implies$ ) Let  $\psi \in \overline{y_X(X)}$ , from Theorem 5.4, we get that  $\psi = \psi^*.(y_X)_*$ . In order to show  $\psi$  is a right-adjoint, we will show that  $\psi^*$  is a right adjoint and that  $(y_X)_*$  is an equivalence.

① In order to show that  $(y_X)_*$  is an equivalence, we need that  $A = (y_X)^*.(y_X)_*$  and  $\tilde{A} = (y_X)_*.(y_X)^*$ .

From proposition 5.3 (b), we have that  $y_X$  is fully faithful, and by Proposition 4.14 (a), this gives us that  $A = (y_X)^*.(y_X)_*$ .

- We are now going to show that  $\tilde{A} \leq (y_X)_*.(y_X)^*$ . Fix any  $a, b \in A$ , we need to find  $c \in A$  such that  $\tilde{c} \subseteq \tilde{a} y_X y_X^o \tilde{b}$ .

$$(\tilde{a} y_X y_X^o \tilde{b})(\psi) = (\tilde{a} \tilde{b})(\psi) \supseteq \tilde{c} \tilde{c}(\psi) \supseteq \tilde{c}(\psi)$$

In the above equation, the equality holds because  $\psi \in \overline{y_X(X)}$ , gives the existence of  $x^* = \tilde{b}(\psi)$ . And the first inequality is given by down-directedness of  $\tilde{A}$ , whereas the second one holds because  $\tilde{c}$  is reflexive, as  $\tilde{A}$  is a quasi-uniformity.

- To show that  $\tilde{A} \geq (y_X)_*.(y_X)^*$ , fix any  $a \in A$ . By quasi-uniformity of  $\tilde{A}$ , there exists  $\tilde{b} \in \tilde{A}$  such that  $\tilde{b} \tilde{b} \subseteq a$ . We will show that  $\tilde{a} \supseteq \tilde{b} y_X y_X^o \tilde{b}$ :

$$\psi(\tilde{b} y_X y_X^o \tilde{b}) \phi \implies \psi(\tilde{b} \tilde{b}) \phi \implies \psi \tilde{a} \phi.$$

② In order to show that  $\psi^*$  is a right adjoint to  $\psi_*$ , due to the 2-categorical structure of ProMod, we need to show that  $\tilde{A} \geq \psi_*.\psi^*$  and  $\psi_*.\psi^* \geq 1$ .

# Proof: Yoneda Corollary II

- To show that  $\tilde{A} \geq \psi_* \cdot \psi^* = \psi_* \cdot \psi^\circ \cdot \tilde{A}$ , fix any  $a \in A$ . We will show that  $\psi_* \cdot \psi^\circ \cdot \tilde{a} \subseteq \tilde{a}$ . Using definition of  $\psi_*$ , for any  $\phi \in y_X(X)$ , we get:

$$(\psi_* \cdot \psi^\circ \cdot \tilde{a})(\phi) = \psi_* \cdot \psi^\circ(\tilde{a}(\phi)) = \begin{cases} \phi & \text{if } \tilde{a}(\phi) \neq \psi \\ \psi = \psi_* \cdot \psi^\circ(\psi) & \text{if } \tilde{a}(\phi) = \psi \end{cases}.$$

The above equation gives that  $\phi(\psi_* \cdot \psi^\circ \cdot \tilde{a})\psi$  implies  $\phi\tilde{a}\psi$ . Hence, we have that  $\tilde{a} \supseteq \psi_* \cdot \psi^\circ \cdot \tilde{a}$ .

- We will show that  $\psi_* \cdot \psi^* \geq 1$ , that is  $\star(\psi^\circ \cdot \tilde{a} \cdot \psi_*)\star$ . Using definition of  $\psi_*$ ,

$$(\psi^\circ \cdot \tilde{a} \cdot \psi_*)(\star) = (\psi^\circ \cdot \tilde{a})(\psi_*(\star)) = (\psi^\circ \cdot \tilde{a})(\psi) = \psi^\circ(\tilde{a}(\psi)).$$

By the quasi-uniformity of  $\tilde{A}$ , we get that  $\tilde{a}$  is reflexive, and hence,  $\psi\tilde{a}\psi$ . So, from the above equation, we have that  $\star \in \psi^\circ(\psi) \subseteq (\psi^\circ \cdot \tilde{a} \cdot \psi_*)(\star)$ .

- ② (  $\Leftarrow$  ) Suppose  $\psi$  is a right adjoint. Need to show that for any  $a \in A$ ,  $\exists x^* \in y_X(X)$  such that  $\psi \tilde{a} x^* \tilde{a} \psi$ . Fix  $a \in A$ . Because  $\psi$  is a right-adjoint, there exists a promodule  $\phi : 1 \rightarrow X$  such that  $\phi \cdot \psi \leq A$  and  $1 \leq \psi \cdot \phi$ . From  $\phi \cdot \psi \leq A$ , we get that:

$$\exists p \in \phi, q \in \psi \text{ such that } a \supseteq p \cdot q. \quad (1)$$

Because  $\phi$  and  $\psi$  are promodules,

$$A \cdot \phi \leq \phi \text{ gives the existence of } p' \in \phi \text{ such that } p \supseteq a' p', \quad (2)$$

$$A \cdot \psi \leq \psi \text{ gives the existence of } q' \in \psi \text{ and } a'' \in A \text{ such that } q \supseteq a'' q'. \quad (3)$$

# Proof: Yoneda Corollary III

Now, from  $1 \leq \psi.\phi$ , we get that  $q' p'$  is reflexive i.e.  $\star(q' p')\star$ . By the definition of composition we get the existence of an  $x \in X$  such that  $\star p' x q' \star$ . Now, considering  $x$  as a map,  $x : 1 \rightarrow X$  defined as  $\star \mapsto x$ ,

$$x q' \star \text{ i.e. } \star \in q'(x) \text{ gives that } q' \supseteq x^\circ, \quad (4)$$

$$\star p' x \text{ i.e. } x \in p'(\star) \text{ gives that } p' \supseteq x. \quad (5)$$

Thus, by using inequalities (1),(2) and (3), we get:

$$a \supseteq p q \supseteq a' p' q' a'' . \quad (6)$$

By definition of  $\tilde{a}$ , to show  $\psi \tilde{a} x^*$ , we need that  $\psi \leq x^* a = x^\circ.A.a$ . We are now going to show that for any  $b \in A$ ,  $x^\circ b a \supseteq q'$ :

$$x^\circ b a \supseteq x^\circ b a' p' q' \supseteq x^\circ b a' x q' \supseteq x^\circ x q' = q' .$$

Where the first inequality comes from (6) by using reflexivity of  $a''$  and then left-multiplying by  $x^\circ$ . The second inequality comes from (5), third one from reflexivity of  $b$  and  $a'$ , and the last one is given by Lemma 3.7.

In order to show  $x^* \tilde{a} \psi$ , by definition of  $\tilde{a}$ , need that  $x^\circ.A = x^* \leq \psi a$ . Fix  $k \in \psi$ . We will now show  $k a \supseteq x^\circ a''$ :

$$a \supseteq a' p' q' a'' \supseteq p' q' a'' \supseteq p' x^\circ a'' . \quad (7)$$

## Proof: Yoneda Corollary IV

Where the first inequality is given by (6), second one is due to reflexiveness of  $a'$  and the third inequality comes by using (4). Left-multiplying (7) with  $k$  gives the following:

$$ka \supseteq k p' x^{\circ} a'' \text{ that is, for any } z \in X, \quad z(ka) \star \implies z(k p' x^{\circ} a'') \star . \quad (8)$$

As  $\psi$  is a right adjoint to  $\phi$ , we have  $1 \leq \psi.\phi$ , giving that  $\star(k p') \star$ . So, using the implication in (8), we get that  $z(ka) \star$  implies  $z(x^{\circ} a'') \star (k p') \star$ , which in turn gives that  $z(x^{\circ} a'') \star$ . Hence, we get that  $ka \supseteq x^{\circ} a''$

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

Let  $\mathcal{A}$  be a locally small category. Fix an object  $A \in \mathcal{A}$  and a presheaf  $X$  on  $\mathcal{A}$ .

- Define  $\hat{\alpha} : \mathcal{C}(H_A, X) \rightarrow X(A)$  for any  $\alpha : H_A \rightarrow X$ , as  $\hat{\alpha} := \alpha_A(1_A)$ . As  $1_A \in Set(A, A) = H_A(A)$ , definition of  $\alpha_A$  gives that  $\alpha_A(1_A) \in X(A)$ .

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

Let  $\mathcal{A}$  be a locally small category. Fix an object  $A \in \mathcal{A}$  and a presheaf  $X$  on  $\mathcal{A}$ .

- Define  $\hat{\cdot} : \mathcal{C}(H_A, X) \rightarrow X(A)$  for any  $\alpha : H_A \rightarrow X$ , as  $\hat{\alpha} := \alpha_A(1_A)$ . As  $1_A \in Set(A, A) = H_A(A)$ , definition of  $\alpha_A$  gives that  $\alpha_A(1_A) \in X(A)$ .
- Define  $\tilde{\cdot} : X(A) \rightarrow \mathcal{C}(H_A, X)$  for any  $x \in X(A)$  as the natural transformation  $\tilde{x} : H_A \rightarrow X$  whose  $K$ -component is the function mapping each morphism  $p \in \mathcal{A}(K, A)$  to  $(X(p))(x)$ . That is,  $\tilde{x}_K(p) := (X(p))(x)$ .

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

Let  $\mathcal{A}$  be a locally small category. Fix an object  $A \in \mathcal{A}$  and a presheaf  $X$  on  $\mathcal{A}$ .

- Define  $\hat{\cdot} : \mathcal{C}(H_A, X) \rightarrow X(A)$  for any  $\alpha : H_A \rightarrow X$ , as  $\hat{\alpha} := \alpha_A(1_A)$ . As  $1_A \in Set(A, A) = H_A(A)$ , definition of  $\alpha_A$  gives that  $\alpha_A(1_A) \in X(A)$ .
- Define  $\tilde{\cdot} : X(A) \rightarrow \mathcal{C}(H_A, X)$  for any  $x \in X(A)$  as the natural transformation  $\tilde{x} : H_A \rightarrow X$  whose  $K$ -component is the function mapping each morphism  $p \in \mathcal{A}(K, A)$  to  $(X(p))(x)$ . That is,  $\tilde{x}_K(p) := (X(p))(x)$ .
- We are going to show that  $\tilde{x}$  is a natural transformation.

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

Let  $\mathcal{A}$  be a locally small category. Fix an object  $A \in \mathcal{A}$  and a presheaf  $X$  on  $\mathcal{A}$ .

- Define  $\hat{\alpha} : \mathcal{C}(H_A, X) \rightarrow X(A)$  for any  $\alpha : H_A \rightarrow X$ , as  $\hat{\alpha} := \alpha_A(1_A)$ . As  $1_A \in Set(A, A) = H_A(A)$ , definition of  $\alpha_A$  gives that  $\alpha_A(1_A) \in X(A)$ .
- Define  $\tilde{x} : X(A) \rightarrow \mathcal{C}(H_A, X)$  for any  $x \in X(A)$  as the natural transformation  $\tilde{x} : H_A \rightarrow X$  whose  $K$ -component is the function mapping each morphism  $p \in \mathcal{A}(K, A)$  to  $(X(p))(x)$ . That is,  $\tilde{x}_K(p) := (X(p))(x)$ .
- We are going to show that  $\tilde{x}$  is a natural transformation.
- Fix objects  $K, L \in \mathcal{A}$  and morphism  $q \in \mathcal{A}^{op}(K, L)$ .

$$\begin{array}{ccc} \mathcal{A}(K, A) & \xrightarrow{- \circ q} & \mathcal{A}(L, A) \\ \tilde{x}_K \downarrow & & \downarrow \tilde{x}_L \\ X(K) & \xrightarrow{X(q)} & X(L) \end{array} \quad \text{commutes .}$$

So, for any  $f : K \rightarrow A$ , need that  $\tilde{x}_L(f \circ q) = X(q) \circ \tilde{x}_K(f)$ . Using the definition of  $\tilde{x}$  gives the following.

$$LHS = \tilde{x}_L(f \circ q) = (X(f \circ q))(x)$$

$$RHS = X(q) \circ \tilde{x}_K(f) = (X(q))(X(f)(x)) = (X(q) \circ X(f))(x)$$

And as  $X$  is a contravariant functor,  $X(f \circ q) = X(q) \circ X(f)$ , giving that LHS=RHS.



## Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

- Need to show that  $\hat{\phantom{x}}$  and  $\sim$  are mutually inverse.

## Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

- Need to show that  $\hat{\phantom{x}}$  and  $\tilde{\phantom{x}}$  are mutually inverse.
- For any  $x \in X(A)$ ,  $\hat{\tilde{x}} = \tilde{x}_A(1_A) = \left(X(1_A)\right)(x) = 1_{X(A)}(x) = x$ . For any  $\alpha \in \mathcal{C}(H_A, X)$ , need to show that  $\tilde{\hat{\alpha}} = \alpha$ . So, it's required that each of their component are equal.

## Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

- Need to show that  $\hat{\phantom{x}}$  and  $\tilde{\phantom{x}}$  are mutually inverse.
- For any  $x \in X(A)$ ,  $\hat{\tilde{x}} = \tilde{x}_A(1_A) = \left(X(1_A)\right)(x) = 1_{X(A)}(x) = x$ . For any  $\alpha \in \mathcal{C}(H_A, X)$ , need to show that  $\tilde{\tilde{\alpha}} = \alpha$ . So, it's required that each of their component are equal.
- As both  $\tilde{\tilde{\alpha}}$  and  $\alpha$  are natural transformations between functors that go to the category  $Set$ , each of their components is a function. So, need to show that for any  $f \in \mathcal{A}(K, A) = H_A(K)$ ,  $\left(\tilde{\tilde{\alpha}}\right)_K(f) = \alpha_K(f)$ .

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

- Need to show that  $\hat{\cdot}$  and  $\sim$  are mutually inverse.
- For any  $x \in X(A)$ ,  $\hat{\tilde{x}} = \tilde{x}_A(1_A) = \left(X(1_A)\right)(x) = 1_{X(A)}(x) = x$ . For any  $\alpha \in \mathcal{C}(H_A, X)$ , need to show that  $\tilde{\tilde{\alpha}} = \alpha$ . So, it's required that each of their component are equal.
- As both  $\tilde{\tilde{\alpha}}$  and  $\alpha$  are natural transformations between functors that go to the category  $Set$ , each of their components is a function. So, need to show that for any  $f \in \mathcal{A}(K, A) = H_A(K)$ ,  $\left(\tilde{\tilde{\alpha}}\right)_K(f) = \alpha_K(f)$ .
- Using first the definition of  $\sim$  and then that of  $\hat{\alpha}$  gives:

$$LHS = \tilde{\tilde{\alpha}}_K(f) = \left(X(f)\right)(\hat{\alpha}) = \left(X(f)\right)(\alpha_A(1_A)) \quad (1)$$

And as  $f \in \mathcal{A}(K, A)$ , we also have the following.

$$RHS = \alpha_K(f) = \alpha_K(1_A \circ f) \quad (2)$$

# Showing isomorphism between $[\mathcal{A}^{op}, Set](H_A, X)$ and $X(A)$

- Need to show that  $\hat{\cdot}$  and  $\tilde{\cdot}$  are mutually inverse.
- For any  $x \in X(A)$ ,  $\hat{\tilde{x}} = \tilde{x}_A(1_A) = \left(X(1_A)\right)(x) = 1_{X(A)}(x) = x$ . For any  $\alpha \in \mathcal{C}(H_A, X)$ , need to show that  $\tilde{\hat{\alpha}} = \alpha$ . So, it's required that each of their components are equal.
- As both  $\tilde{\hat{\alpha}}$  and  $\alpha$  are natural transformations between functors that go to the category  $Set$ , each of their components is a function. So, need to show that for any  $f \in \mathcal{A}(K, A) = H_A(K)$ ,  $\left(\tilde{\hat{\alpha}}\right)_K(f) = \alpha_K(f)$ .
- Using first the definition of  $\tilde{\cdot}$  and then that of  $\hat{\cdot}$  gives:

$$LHS = \tilde{\hat{\alpha}}_K(f) = \left(X(f)\right)(\hat{\alpha}) = \left(X(f)\right)(\alpha_A(1_A)) \quad (1)$$

And as  $f \in \mathcal{A}(K, A)$ , we also have the following.

$$RHS = \alpha_K(f) = \alpha_K(1_A \circ f) \quad (2)$$

- Because  $\alpha$  is a natural transformation, the following square commutes for  $1_A$ :

$$\begin{array}{ccc} \mathcal{A}(A, A) & \xrightarrow{- \circ f} & \mathcal{A}(K, A) \\ \alpha_A \downarrow & & \downarrow \alpha_K \\ X(A) & \xrightarrow{X(f)} & X(K) \end{array} ,$$

which gives that  $\alpha_K(1_A \circ f) = \left(X(f)\right)(\alpha_A(1_A))$ . Hence, we get that

$RHS = LHS$ , giving us that  $\hat{\cdot}$  and  $\tilde{\cdot}$  define a set isomorphism, as  $\alpha_K$  being a function,  $RHS$  is a set.

## Showing natural in $X$

- 1 By using the very first two Lemmas, it's enough to show that  $\hat{\phantom{x}}$  is natural in  $X$  and natural in  $A$ .

Showing natural in  $X$ 

- 1 By using the very first two Lemmas, it's enough to show that  $\hat{\phantom{x}}$  is natural in  $X$  and natural in  $A$ .
- 2 Fix any  $A \in \mathcal{A}$ . Need, for presheaves  $X, Y \in \mathcal{C}$  and natural transformation  $\beta \in \mathcal{C}(X, Y)$ , the following square to commute:

$$\begin{array}{ccc}
 \mathcal{C}(H_A, X) & \xrightarrow{\beta \circ -} & \mathcal{C}(H_A, Y) \\
 \hat{\phantom{x}} \downarrow & & \downarrow \hat{\phantom{x}} \\
 X(A) & \xrightarrow{\beta_A} & Y(A)
 \end{array}
 .$$

# Showing natural in $X$

- 1 By using the very first two Lemmas, it's enough to show that  $\hat{\phantom{x}}$  is natural in  $X$  and natural in  $A$ .
- 2 Fix any  $A \in \mathcal{A}$ . Need, for presheaves  $X, Y \in \mathcal{C}$  and natural transformation  $\beta \in \mathcal{C}(X, Y)$ , the following square to commute:

$$\begin{array}{ccc}
 \mathcal{C}(H_A, X) & \xrightarrow{\beta \circ -} & \mathcal{C}(H_A, Y) \\
 \hat{\phantom{x}} \downarrow & & \downarrow \hat{\phantom{x}} \\
 X(A) & \xrightarrow{\beta_A} & Y(A)
 \end{array}
 .$$

- 3 So, for any  $\alpha : H_A \rightarrow X$ , we need that  $(\hat{\phantom{x}} \circ (\beta \circ -))(\alpha) = (\beta_A \circ \hat{\phantom{x}})(\alpha)$ .



# Showing natural in $X$

- By using the very first two Lemmas, it's enough to show that  $\hat{\phantom{x}}$  is natural in  $X$  and natural in  $A$ .
- Fix any  $A \in \mathcal{A}$ . Need, for presheaves  $X, Y \in \mathcal{C}$  and natural transformation  $\beta \in \mathcal{C}(X, Y)$ , the following square to commute:

$$\begin{array}{ccc}
 \mathcal{C}(H_A, X) & \xrightarrow{\beta \circ -} & \mathcal{C}(H_A, Y) \\
 \hat{\phantom{x}} \downarrow & & \downarrow \hat{\phantom{x}} \\
 X(A) & \xrightarrow{\beta_A} & Y(A)
 \end{array}$$

- So, for any  $\alpha : H_A \rightarrow X$ , we need that  $(\hat{\phantom{x}} \circ (\beta \circ -))(\alpha) = (\beta_A \circ \hat{\phantom{x}})(\alpha)$ .
- Using the definition of  $(\beta \circ -)$  and  $\hat{\phantom{x}}$  gives:

$$\begin{aligned}
 LHS &= (\hat{\phantom{x}} \circ (\beta \circ -))(\alpha) = ((\widehat{\beta \circ -})(\alpha)) = (\widehat{\beta \circ \alpha}) = (\beta \circ \alpha)_A(1_A) \\
 RHS &= (\beta_A \circ \hat{\phantom{x}})(\alpha) = \beta_A(\hat{\alpha}) = (\beta_A \circ \alpha_A)(1_A)
 \end{aligned}$$

As  $\alpha \in \mathcal{C}(H_A, X)$  and  $\beta \in \mathcal{C}(X, Y)$  are morphisms in  $\mathcal{C}$ , composition in  $\mathcal{C}$  gives  $(\beta \circ \alpha)_A = \beta_A \circ \alpha_A$ .

Showing natural in  $A$ 

- ① Fix any  $X \in \mathcal{C}$  Need that for objects  $A, B \in \mathcal{A}$  and morphism  $f \in \mathcal{A}^{op}(A, B)$ , the following square commutes:

$$\begin{array}{ccc} \mathcal{C}(H_A, X) & \xrightarrow{- \circ H_f} & \mathcal{C}(H_B, X) \\ \downarrow \hat{\phantom{f}} & & \downarrow \hat{\phantom{f}} \\ X(A) & \xrightarrow{X(f)} & X(B) \end{array},$$

Where  $H_f$  denotes  $(f \circ \_)$ . So, for any  $\alpha : H_A \rightarrow X$ , we need that  $(\hat{\phantom{f}} \circ H_f)(\alpha) = ((X(f)) \circ \hat{\phantom{f}})(\alpha)$ .

Showing natural in  $A$ 

- ① Fix any  $X \in \mathcal{C}$ . Need that for objects  $A, B \in \mathcal{A}$  and morphism  $f \in \mathcal{A}^{op}(A, B)$ , the following square commutes:

$$\begin{array}{ccc} \mathcal{C}(H_A, X) & \xrightarrow{- \circ H_f} & \mathcal{C}(H_B, X) \\ \downarrow \hat{\phantom{f}} & & \downarrow \hat{\phantom{f}} \\ X(A) & \xrightarrow{X(f)} & X(B) \end{array},$$

Where  $H_f$  denotes  $(f \circ \_)$ . So, for any  $\alpha : H_A \rightarrow X$ , we need that  $(\hat{\phantom{f}} \circ H_f)(\alpha) = ((X(f)) \circ \hat{\phantom{f}})(\alpha)$ .

- ② Using definition of  $H_f$  and  $\hat{\phantom{f}}$ , we get:

$$\begin{aligned} LHS &= (\hat{\phantom{f}} \circ H_f)(\alpha) = \widehat{\alpha \circ H_f} = (\alpha \circ H_f)_B(1_B) = \alpha_B(f \circ 1_B) = \alpha_B(1_A \circ f) \\ RHS &= ((X(f)) \circ \hat{\phantom{f}})(\alpha) = (X(f))(\hat{\alpha}) = (X(f))(\alpha_A(1_A)) \end{aligned}$$

Showing natural in  $A$ 

- ① Fix any  $X \in \mathcal{C}$ . Need that for objects  $A, B \in \mathcal{A}$  and morphism  $f \in \mathcal{A}^{op}(A, B)$ , the following square commutes:

$$\begin{array}{ccc} \mathcal{C}(H_A, X) & \xrightarrow{- \circ H_f} & \mathcal{C}(H_B, X) \\ \downarrow \hat{\phantom{f}} & & \downarrow \hat{\phantom{f}} \\ X(A) & \xrightarrow{X(f)} & X(B) \end{array},$$

Where  $H_f$  denotes  $(f \circ \_)$ . So, for any  $\alpha : H_A \rightarrow X$ , we need that  $(\hat{\phantom{f}} \circ H_f)(\alpha) = ((X(f)) \circ \hat{\phantom{f}})(\alpha)$ .

- ② Using definition of  $H_f$  and  $\hat{\phantom{f}}$ , we get:

$$\begin{aligned} LHS &= (\hat{\phantom{f}} \circ H_f)(\alpha) = \widehat{\alpha \circ H_f} = (\alpha \circ H_f)_B(1_B) = \alpha_B(f \circ 1_B) = \alpha_B(1_A \circ f) \\ RHS &= ((X(f)) \circ \hat{\phantom{f}})(\alpha) = (X(f))(\hat{\alpha}) = (X(f))(\alpha_A(1_A)) \end{aligned}$$

- ③ By using equality of equations (1) and (2), for  $f \in \mathcal{A}(B, A)$ , we get that  $(X(f))(\alpha_A(1_A)) = \alpha_B(1_A \circ f)$ . Hence,  $RHS = LHS$ .

## Natural transformations from $H_\star$ to $H_\star$ are bijections on $G$ .

- Let  $G$  be a group. Define category  $\mathcal{A}$  with a single object  $\star$ . And let the morphisms of  $\mathcal{A}$  be elements of  $G$ . Then,  $G$  and  $\mathcal{A}(\star, \star)$  have the same elements and rule of composition, so there exists a group isomorphism  $\psi : \mathcal{A}(\star, \star) \rightarrow G$ .

## Natural transformations from $H_\star$ to $H_\star$ are bijections on $G$ .

- Let  $G$  be a group. Define category  $\mathcal{A}$  with a single object  $\star$ . And let the morphisms of  $\mathcal{A}$  be elements of  $G$ . Then,  $G$  and  $\mathcal{A}(\star, \star)$  have the same elements and rule of composition, so there exists a group isomorphism  $\psi : \mathcal{A}(\star, \star) \rightarrow G$ .
- As  $\mathcal{A}^{op}$  is a category with a single object, each natural transformation

$\alpha : \mathcal{A}^{op} \begin{array}{c} \xrightarrow{H_\star} \\ \Downarrow \alpha \\ \xrightarrow{H_\star} \end{array} \text{Set}$  has only one component, that is  $\alpha_\star$ . Therefore, we can identify  $\alpha$  with  $\alpha_\star$ .

## Natural transformations from $H_\star$ to $H_\star$ are bijections on $G$ .

- Let  $G$  be a group. Define category  $\mathcal{A}$  with a single object  $\star$ . And let the morphisms of  $\mathcal{A}$  be elements of  $G$ . Then,  $G$  and  $\mathcal{A}(\star, \star)$  have the same elements and rule of composition, so there exists a group isomorphism  $\psi : \mathcal{A}(\star, \star) \rightarrow G$ .
- As  $\mathcal{A}^{op}$  is a category with a single object, each natural transformation

$$\alpha : \mathcal{A}^{op} \begin{array}{c} \xrightarrow{H_\star} \\ \Downarrow \alpha \\ \xrightarrow{H_\star} \end{array} \text{Set} \quad \text{has only one component, that is } \alpha_\star. \quad \text{Therefore, we can}$$

identify  $\alpha$  with  $\alpha_\star$ .

- Using naturality of  $\alpha$ , we get that

$$\begin{array}{ccc} \mathcal{A}(\star, \star) & \xrightarrow{- \circ f} & \mathcal{A}(\star, \star) \\ \alpha_\star \downarrow & & \downarrow \alpha_\star \\ \mathcal{A}(\star, \star) & \xrightarrow{- \circ f} & \mathcal{A}(\star, \star) \end{array} \quad \text{the square commutes for any } f \in \mathcal{A}(\star, \star).$$

## Natural transformations from $H_\star$ to $H_\star$ are bijections on $G$ .

- Let  $G$  be a group. Define category  $\mathcal{A}$  with a single object  $\star$ . And let the morphisms of  $\mathcal{A}$  be elements of  $G$ . Then,  $G$  and  $\mathcal{A}(\star, \star)$  have the same elements and rule of composition, so there exists a group isomorphism  $\psi : \mathcal{A}(\star, \star) \rightarrow G$ .
- As  $\mathcal{A}^{op}$  is a category with a single object, each natural transformation

$$\alpha : \mathcal{A}^{op} \begin{array}{c} \xrightarrow{H_\star} \\ \Downarrow \alpha \\ \xrightarrow{H_\star} \end{array} \text{Set} \quad \text{has only one component, that is } \alpha_\star. \quad \text{Therefore, we can}$$

identify  $\alpha$  with  $\alpha_\star$ .

- Using naturality of  $\alpha$ , we get that

$$\begin{array}{ccc} \mathcal{A}(\star, \star) & \xrightarrow{_{\circ}f} & \mathcal{A}(\star, \star) \\ \alpha_\star \downarrow & & \downarrow \alpha_\star \\ \mathcal{A}(\star, \star) & \xrightarrow{_{\circ}f} & \mathcal{A}(\star, \star) \end{array} \quad \text{the square commutes for any } f \in \mathcal{A}(\star, \star).$$

- Applying the identity of  $\star$  in  $\mathcal{A}$  in above square gives us the following equation:

$$((_{\circ}f) \circ \alpha_\star)(1_\star) = (\alpha_\star \circ (_{\circ}f))(1_\star) \implies \alpha_\star(f) = \alpha_\star(1_\star) \circ f \implies \alpha_\star(f) = \alpha_\star(1_\star) \cdot f$$

Thus, every natural transformation  $\alpha$  is defined in terms of its value at  $1_\star$ . This can be considered as left multiplication by  $\alpha_\star(1_\star)$  in  $G$ , which we know is a bijection on  $G$ .



## Natural transformations from $H_\star$ to $H_\star$ are bijections on $G$ .

- Let  $G$  be a group. Define category  $\mathcal{A}$  with a single object  $\star$ . And let the morphisms of  $\mathcal{A}$  be elements of  $G$ . Then,  $G$  and  $\mathcal{A}(\star, \star)$  have the same elements and rule of composition, so there exists a group isomorphism  $\psi : \mathcal{A}(\star, \star) \rightarrow G$ .
- As  $\mathcal{A}^{op}$  is a category with a single object, each natural transformation

$$\alpha : \mathcal{A}^{op} \begin{array}{c} \xrightarrow{H_\star} \\ \Downarrow \alpha \\ \xrightarrow{H_\star} \end{array} \text{Set} \quad \text{has only one component, that is } \alpha_\star. \quad \text{Therefore, we can}$$

identify  $\alpha$  with  $\alpha_\star$ .

- Using naturality of  $\alpha$ , we get that

$$\begin{array}{ccc} \mathcal{A}(\star, \star) & \xrightarrow{- \circ f} & \mathcal{A}(\star, \star) \\ \text{the square } \alpha_\star \downarrow & & \downarrow \alpha_\star \\ \mathcal{A}(\star, \star) & \xrightarrow{- \circ f} & \mathcal{A}(\star, \star) \end{array} \quad \text{commutes for any } f \in \mathcal{A}(\star, \star).$$

- Applying the identity of  $\star$  in  $\mathcal{A}$  in above square gives us the following equation:

$$((\circ f) \circ \alpha_\star)(1_\star) = (\alpha_\star \circ (\circ f))(1_\star) \implies \alpha_\star(f) = \alpha_\star(1_\star) \circ f \implies \alpha_\star(f) = \alpha_\star(1_\star) \cdot f$$

Thus, every natural transformation  $\alpha$  is defined in terms of its value at  $1_\star$ . This can be considered as left multiplication by  $\alpha_\star(1_\star)$  in  $G$ , which we know is a bijection on  $G$ .

- So far we have shown that the collection  $[\mathcal{A}^{op}, \text{Set}](H_\star, H_\star)$  of all  $\alpha : H_\star \rightarrow H_\star$  is a collection of bijections on  $G$ .

The collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .

## The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.

## The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .

## The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .
- Since  $\gamma_*(1_*)$  belongs to  $\mathcal{A}(*, *)$ , let us call  $\psi(\gamma_*(1_*)) = h \in G$ . Thus, there exists  $h^{-1} \in G$ .

## The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .
- Since  $\gamma_*(1_*)$  belongs to  $\mathcal{A}(*, *)$ , let us call  $\psi(\gamma_*(1_*)) = h \in G$ . Thus, there exists  $h^{-1} \in G$ .
- As  $\psi$  is onto, there exists  $a \in \mathcal{A}(*, *)$  such that  $\psi(a) = h^{-1}$ .

# The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .
- Since  $\gamma_*(1_*)$  belongs to  $\mathcal{A}(*, *)$ , let us call  $\psi(\gamma_*(1_*)) = h \in G$ . Thus, there exists  $h^{-1} \in G$ .
- As  $\psi$  is onto, there exists  $a \in \mathcal{A}(*, *)$  such that  $\psi(a) = h^{-1}$ .
- We know that any natural transformation  $\alpha$  is defined in terms of  $\alpha_*(1_*) \in \mathcal{A}(*, *)$ . Thus, we define  $\delta : H_* \rightarrow H_*$  with  $\delta_*(1_*) = a$ . Giving us that  $h^{-1} = \psi(\delta_*(1_*))$ . And as  $\psi$  is a group isomorphism,

$$1_* = \psi^{-1}(h.h^{-1}) = \psi^{-1}(h).\psi^{-1}(h^{-1}) = (\gamma_*(1_*)).(\delta_*(1_*)).$$

## The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .
- Since  $\gamma_*(1_*)$  belongs to  $\mathcal{A}(*, *)$ , let us call  $\psi(\gamma_*(1_*)) = h \in G$ . Thus, there exists  $h^{-1} \in G$ .
- As  $\psi$  is onto, there exists  $a \in \mathcal{A}(*, *)$  such that  $\psi(a) = h^{-1}$ .
- We know that any natural transformation  $\alpha$  is defined in terms of  $\alpha_*(1_*) \in \mathcal{A}(*, *)$ . Thus, we define  $\delta : H_* \rightarrow H_*$  with  $\delta_*(1_*) = a$ . Giving us that  $h^{-1} = \psi(\delta_*(1_*))$ . And as  $\psi$  is a group isomorphism,

$$1_* = \psi^{-1}(h.h^{-1}) = \psi^{-1}(h).\psi^{-1}(h^{-1}) = (\gamma_*(1_*)).(\delta_*(1_*)).$$

- This gives us that  $\delta$  and  $\gamma$  are inverses, because for any  $k \in \mathcal{A}(*, *)$ ,

$$(\gamma \circ \delta)_*(k) = \gamma_*(\delta_*(k)) = \gamma_*(\delta_*(1_*) \cdot k) = (\gamma_*(1_*)).(\delta_*(1_*) \cdot k) = 1_* \cdot k = k$$



# The collection $[\mathcal{A}^{op}, Set](H_*, H_*)$ is a group.

- We will show that the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group with respect to composition in the category  $[\mathcal{A}^{op}, Set]$ .
- As  $[\mathcal{A}^{op}, Set]$  is a category, we have that the composition is associative. Also, because this collection contains morphisms with the same source and destination, it is closed under composition. Identity of  $[\mathcal{A}^{op}, Set](H_*, H_*)$  will act as the identity for its group structure.
- We will now show closure under inverses. Fix any  $\gamma : H_* \rightarrow H_*$ .
- Since  $\gamma_*(1_*)$  belongs to  $\mathcal{A}(*, *)$ , let us call  $\psi(\gamma_*(1_*)) = h \in G$ . Thus, there exists  $h^{-1} \in G$ .
- As  $\psi$  is onto, there exists  $a \in \mathcal{A}(*, *)$  such that  $\psi(a) = h^{-1}$ .
- We know that any natural transformation  $\alpha$  is defined in terms of  $\alpha_*(1_*) \in \mathcal{A}(*, *)$ . Thus, we define  $\delta : H_* \rightarrow H_*$  with  $\delta_*(1_*) = a$ . Giving us that  $h^{-1} = \psi(\delta_*(1_*))$ . And as  $\psi$  is a group isomorphism,

$$1_* = \psi^{-1}(h.h^{-1}) = \psi^{-1}(h).\psi^{-1}(h^{-1}) = (\gamma_*(1_*)).(\delta_*(1_*)).$$

- This gives us that  $\delta$  and  $\gamma$  are inverses, because for any  $k \in \mathcal{A}(*, *)$ ,

$$(\gamma \circ \delta)_*(k) = \gamma_*(\delta_*(k)) = \gamma_*(\delta_*(1_*) \cdot k) = (\gamma_*(1_*)).(\delta_*(1_*) \cdot k) = 1_* \cdot k = k$$

- Thus, the collection  $[\mathcal{A}^{op}, Set](H_*, H_*)$  is a group.

## Applying Yoneda Lemma, and showing group isomorphism

- As the collection of elements of  $G$  form a set,  $\mathcal{A}(\star, \star)$  is also a set. Hence,  $\mathcal{A}$  is a locally small category. Because  $\mathcal{A}^{op}$  has the same number of morphisms as  $\mathcal{A}$ , it is also a locally small category, and we may apply Yoneda Lemma to it.

## Applying Yoneda Lemma, and showing group isomorphism

- As the collection of elements of  $G$  form a set,  $\mathcal{A}(\star, \star)$  is also a set. Hence,  $\mathcal{A}$  is a locally small category. Because  $\mathcal{A}^{op}$  has the same number of morphisms as  $\mathcal{A}$ , it is also a locally small category, and we may apply Yoneda Lemma to it.
- Taking  $A = \star$  and  $X = H_\star$ , we get:

$$[\mathcal{A}^{op}, Set](H_\star, H_\star) \hat{=} H_\star(\star),$$

where the isomorphism  $\hat{=}$  is between sets.

## Applying Yoneda Lemma, and showing group isomorphism

- As the collection of elements of  $G$  form a set,  $\mathcal{A}(\star, \star)$  is also a set. Hence,  $\mathcal{A}$  is a locally small category. Because  $\mathcal{A}^{op}$  has the same number of morphisms as  $\mathcal{A}$ , it is also a locally small category, and we may apply Yoneda Lemma to it.
- Taking  $A = \star$  and  $X = H_\star$ , we get:

$$[\mathcal{A}^{op}, Set](H_\star, H_\star) \hat{=} H_\star(\star),$$

where the isomorphism  $\hat{=}$  is between sets.

- From the proof of Yoneda Lemma, we know that the map  $\hat{=}$  acts as  $\alpha \mapsto \alpha_\star(1_\star)$ . Hence, for any  $\alpha, \beta : H_\star \rightarrow H_\star$ ,

$$\widehat{\alpha \circ \beta} = (\alpha \circ \beta)_\star(1_\star) = (\alpha)_\star((\beta)_\star(1_\star)) = ((\alpha)_\star(1_\star)) \cdot ((\beta)_\star(1_\star)) = \hat{\alpha} \cdot \hat{\beta},$$

# Applying Yoneda Lemma, and showing group isomorphism

- As the collection of elements of  $G$  form a set,  $\mathcal{A}(\star, \star)$  is also a set. Hence,  $\mathcal{A}$  is a locally small category. Because  $\mathcal{A}^{op}$  has the same number of morphisms as  $\mathcal{A}$ , it is also a locally small category, and we may apply Yoneda Lemma to it.
- Taking  $A = \star$  and  $X = H_\star$ , we get:

$$[\mathcal{A}^{op}, Set](H_\star, H_\star) \hat{=} H_\star(\star),$$

where the isomorphism  $\hat{=}$  is between sets.

- From the proof of Yoneda Lemma, we know that the map  $\hat{=}$  acts as  $\alpha \mapsto \alpha_\star(1_\star)$ . Hence, for any  $\alpha, \beta : H_\star \rightarrow H_\star$ ,

$$\widehat{\alpha \circ \beta} = (\alpha \circ \beta)_\star(1_\star) = (\alpha)_\star((\beta)_\star(1_\star)) = ((\alpha)_\star(1_\star)) \cdot ((\beta)_\star(1_\star)) = \hat{\alpha} \cdot \hat{\beta},$$

- Using I and II, we get that  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  is a group with all of it's elements being bijections on  $G$ . Thus, it is a subgroup of the symmetric group on  $G$ .

# Applying Yoneda Lemma, and showing group isomorphism

- As the collection of elements of  $G$  form a set,  $\mathcal{A}(\star, \star)$  is also a set. Hence,  $\mathcal{A}$  is a locally small category. Because  $\mathcal{A}^{op}$  has the same number of morphisms as  $\mathcal{A}$ , it is also a locally small category, and we may apply Yoneda Lemma to it.
- Taking  $A = \star$  and  $X = H_\star$ , we get:

$$[\mathcal{A}^{op}, Set](H_\star, H_\star) \hat{\cong} H_\star(\star),$$

where the isomorphism  $\hat{\phantom{x}}$  is between sets.

- From the proof of Yoneda Lemma, we know that the map  $\hat{\phantom{x}}$  acts as  $\alpha \mapsto \alpha_\star(1_\star)$ . Hence, for any  $\alpha, \beta : H_\star \rightarrow H_\star$ ,

$$\widehat{\alpha \circ \beta} = (\alpha \circ \beta)_\star(1_\star) = (\alpha)_\star((\beta)_\star(1_\star)) = ((\alpha)_\star(1_\star)) \cdot ((\beta)_\star(1_\star)) = \hat{\alpha} \cdot \hat{\beta},$$

- Using I and II, we get that  $[\mathcal{A}^{op}, Set](H_\star, H_\star)$  is a group with all of it's elements being bijections on  $G$ . Thus, it is a subgroup of the symmetric group on  $G$ .
- Finally, the isomorphism  $\hat{\phantom{x}}$  is between groups, with the *LHS* being the above mentioned subgroup. And *RHS* being  $\mathcal{A}(\star, \star)$ , which is further isomorphic to group  $G$ :

$$G \stackrel{\psi}{\cong} \mathcal{A}(\star, \star) \hat{\cong} [\mathcal{A}^{op}, Set](H_\star, H_\star) \leq Sym(G).$$

This is precisely the statement of Cayley's theorem.

## Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, Set](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .

## Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, Set](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .
- Thus, we need to show that their  $K$ -components are equal for every  $K \in \mathcal{A}$ . Using the definition of  $H_\bullet(f)$ , this amounts to showing that

for any morphism  $k \in H_X(K)$ ,  $\left(H_\bullet(f)\right)_K(k) = \alpha_K(k)$ , that is  $f \circ k = \alpha_K(k)$ .



## Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, Set](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .
- Thus, we need to show that their  $K$ -components are equal for every  $K \in \mathcal{A}$ . Using the definition of  $H_\bullet(f)$ , this amounts to showing that

$$\text{for any morphism } k \in H_X(K), \left( H_\bullet(f) \right)_K(k) = \alpha_K(k), \text{ that is } f \circ k = \alpha_K(k).$$

- Because  $\alpha_X$  goes from  $H_X(X)$  to  $H_Y(X)$ ,  $\alpha_X(1_X)$  is a morphism in  $\mathcal{A}(X, Y)$ .

# Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, Set](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .
- Thus, we need to show that their  $K$ -components are equal for every  $K \in \mathcal{A}$ . Using the definition of  $H_\bullet(f)$ , this amounts to showing that

$$\text{for any morphism } k \in H_X(K), \left(H_\bullet(f)\right)_K(k) = \alpha_K(k), \text{ that is } f \circ k = \alpha_K(k).$$

- Because  $\alpha_X$  goes from  $H_X(X)$  to  $H_Y(X)$ ,  $\alpha_X(1_X)$  is a morphism in  $\mathcal{A}(X, Y)$ .
- We will show that choosing this morphism to be  $f$  will give us the required result, that is  $(\alpha_X(1_X)) \circ k = \alpha_K(k)$ .

# Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, \text{Set}](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .
- Thus, we need to show that their  $K$ -components are equal for every  $K \in \mathcal{A}$ . Using the definition of  $H_\bullet(f)$ , this amounts to showing that

$$\text{for any morphism } k \in H_X(K), \left(H_\bullet(f)\right)_K(k) = \alpha_K(k), \text{ that is } f \circ k = \alpha_K(k).$$

- Because  $\alpha_X$  goes from  $H_X(X)$  to  $H_Y(X)$ ,  $\alpha_X(1_X)$  is a morphism in  $\mathcal{A}(X, Y)$ .
- We will show that choosing this morphism to be  $f$  will give us the required result, that is  $(\alpha_X(1_X)) \circ k = \alpha_K(k)$ .

- Using the naturality of  $\alpha$ , we get that

$$\begin{array}{ccc} \mathcal{A}(X, X) & \xrightarrow{- \circ k} & \mathcal{A}(K, X) \\ \alpha_X \downarrow & & \downarrow \alpha_K \\ \mathcal{A}(X, Y) & \xrightarrow{- \circ k} & \mathcal{A}(K, Y) \end{array} \quad \text{commutes.}$$

# Functor $H_\bullet$ is full

We will show that the functor  $H_\bullet$  is full and faithful. Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ .

- Fix any objects  $X, Y$  in a locally small category  $\mathcal{A}$ . To show that  $H_\bullet$  is a full, we need to show that for every  $\alpha \in [\mathcal{A}^{op}, Set](H_X, H_Y)$ , there exists a morphism  $f \in \mathcal{A}(X, Y)$  such that  $H_\bullet(f) = \alpha$ .
- Thus, we need to show that their  $K$ -components are equal for every  $K \in \mathcal{A}$ . Using the definition of  $H_\bullet(f)$ , this amounts to showing that

$$\text{for any morphism } k \in H_X(K), \left(H_\bullet(f)\right)_K(k) = \alpha_K(k), \text{ that is } f \circ k = \alpha_K(k).$$

- Because  $\alpha_X$  goes from  $H_X(X)$  to  $H_Y(X)$ ,  $\alpha_X(1_X)$  is a morphism in  $\mathcal{A}(X, Y)$ .
- We will show that choosing this morphism to be  $f$  will give us the required result, that is  $(\alpha_X(1_X)) \circ k = \alpha_K(k)$ .

- Using the naturality of  $\alpha$ , we get that
 

|                       |                           |                       |           |
|-----------------------|---------------------------|-----------------------|-----------|
| $\mathcal{A}(X, X)$   | $\xrightarrow{- \circ k}$ | $\mathcal{A}(K, X)$   |           |
| $\alpha_X \downarrow$ |                           | $\downarrow \alpha_K$ | commutes. |
| $\mathcal{A}(X, Y)$   | $\xrightarrow{- \circ k}$ | $\mathcal{A}(K, Y)$   |           |

- Thus, for the identity morphism  $1_X \in \mathcal{A}(X, X)$ , we get the following:

$$(H_Y(k) \circ \alpha_X)(1_X) = (\alpha_K \circ H_X(k))(1_X) \implies \alpha_K(1_X) \circ k = \alpha_K(k).$$

- Thus, we have that  $H_\bullet$  is a full functor.

## Functor $H_\bullet$ is faithful

- Fix any morphisms  $f, g$  in  $\mathcal{A}(X, Y)$  and suppose  $H_\bullet(f) = H_\bullet(g)$ . In order to show  $H_\bullet$  is faithful, we need to show that  $f = g$ .

## Functor $H_\bullet$ is faithful

- Fix any morphisms  $f, g$  in  $\mathcal{A}(X, Y)$  and suppose  $H_\bullet(f) = H_\bullet(g)$ . In order to show  $H_\bullet$  is faithful, we need to show that  $f = g$ .
- As  $H_\bullet(f)$  and  $H_\bullet(g)$  are equal natural transformations, we have that the action of their  $X$ -components is equal.

## Functor $H_\bullet$ is faithful

- Fix any morphisms  $f, g$  in  $\mathcal{A}(X, Y)$  and suppose  $H_\bullet(f) = H_\bullet(g)$ . In order to show  $H_\bullet$  is faithful, we need to show that  $f = g$ .
- As  $H_\bullet(f)$  and  $H_\bullet(g)$  are equal natural transformations, we have that the action of their  $X$ -components is equal.
- Thus, in particular, for the identity of  $X$ ,  $(H_\bullet(f))_X(1_X) = (H_\bullet(g))_X(1_X)$ .

## Functor $H_\bullet$ is faithful

- Fix any morphisms  $f, g$  in  $\mathcal{A}(X, Y)$  and suppose  $H_\bullet(f) = H_\bullet(g)$ . In order to show  $H_\bullet$  is faithful, we need to show that  $f = g$ .
- As  $H_\bullet(f)$  and  $H_\bullet(g)$  are equal natural transformations, we have that the action of their  $X$ -components is equal.
- Thus, in particular, for the identity of  $X$ ,  $(H_\bullet(f))_X(1_X) = (H_\bullet(g))_X(1_X)$ .
- Using the definition of  $H_\bullet$ , we get that  $f \circ 1_X = g \circ 1_X$ . And as both  $g$  and  $f$  are morphisms from  $X$ , we get that  $f = g$ .



# Proof: Promod is a 2-category I

In order to show that ProMod is a 2-category, need the following:

- ① (1-Identities) For each quasi-uniform space  $(X, A)$ ,  $A : (X, A) \rightarrow (X, A)$  a promodule because  $A.A = A$  by Lemma 4.5.
- ② (1-Composition) Need composition of promodules to be a promodule. Let  $\phi : (X, A) \rightarrow (Y, B)$  and  $\psi : (Y, B) \rightarrow (Z, C)$  be promodules. To show that  $\psi.\phi : (X, A) \rightarrow (Z, C)$  is a promodule, need it to be a prorelation that satisfies the two conditions required to be a promodule:
  - ① By Lemma 3.4, prorelations are closed under composition. Hence,  $\psi.\phi$  is a prorelation
  - ② Need to show that  $\psi.\phi.A \leq \psi.\phi$ . So, Fix  $p \in \psi$  and  $q \in \phi$ . As  $\phi$  is a promodule,  $\phi.A \leq \phi$  gives that there exists  $q' \in \phi$  and  $a \in A$  such that  $q' a \subseteq q$ . Thus,  $p q' a \subseteq p q$ .
  - ③ Need to show that  $C.\psi.\phi \leq \psi.\phi$ . Fix  $p \in \psi$  and  $q \in \phi$ . Because  $\psi$  is a promodule,  $C.\psi \leq \psi$  gives that there exists  $c \in C$  and  $p' \in \psi$  such that  $c p' \subseteq p$ . Thus,  $c p' q \subseteq p q$
- ③ (2-Identities) As every promodule is contained in itself, always have  $\psi \leq \psi$ . Define this comparison to be the identity 2-cell for  $\psi$  and denote it by  $\leq_\psi$
- ④ (Vertical 2-composition) For promodules  $\psi, \phi, \delta : (X, A) \rightarrow (Y, B)$ , if there is a 2-cell from  $\psi$  to  $\phi$  and another one from  $\phi$  to  $\delta$  i.e.  $\psi \leq \phi \leq \delta$ , then by transitivity of the partial order,  $\psi \leq \delta$  i.e. there's a 2-cell from  $\psi$  to  $\delta$ .
- ⑤ (Horizontal 2-composition) If there are promodules  $\psi, \psi' : (X, A) \rightarrow (Y, B)$  and  $\phi, \phi' : (Y, B) \rightarrow (Z, C)$  such that  $\psi \leq \psi'$  and  $\phi \leq \phi'$ , need to show that  $\psi.\phi \leq \psi'.\phi'$ . Fix  $p' \in \psi'$  and  $q' \in \phi'$ . As  $\psi \leq \psi'$ ,  $\exists p \in \psi : p \subseteq p'$  and as  $\psi \leq \psi'$ ,  $\exists q \in \phi : q \subseteq q'$ . Thus,  $p q \subseteq p' q'$

# Proof: Promod is a 2-category II

- ⑥ (1-Identity) Need to show that for any promodule  $\phi : (X, A) \rightarrow (Y, B)$ ,  $\phi.A = \phi = B.\phi$ . By quasi-uniformity of  $A$ , every  $a \in A$ , is reflexive. Thus, for any  $p \in \phi$  and  $a \in A$ ,  $p = p.\Delta_X \subseteq p a$  giving that  $\phi \leq \phi.A$ . And as  $\phi$  is a promodule,  $\phi \geq \phi.A$ . Hence, by anti-symmetry of the partial order,  $\phi = \phi.A$ . Similarly, By quasi-uniformity of  $B$ , every  $b \in B$ , is reflexive. Thus, for any  $p \in \phi$  and  $b \in B$ ,  $p = \Delta_Y.p \subseteq b p$  giving that  $\phi \leq B.\phi$ . And as  $\phi$  is a promodule,  $\phi \geq B.\phi$ . Hence,  $\phi = B.\phi$ .
- ⑦ (1-Associativity) As composition of relations is associative, so too is the composition of preorders directly giving that composition of promodules i.e. 1-cells is associative.
- ⑧ (Vertical 2-Identity) Let  $\leq : \psi \rightarrow \phi$  be a 2-cell i.e.  $\psi \leq \phi$ . By our definition of identity 2-cell,  $\leq_\psi \cdot \leq_1$  means precisely that  $\psi \leq \psi \leq \phi$ , and by transitivity, this is equivalent to  $\psi \leq \phi$ . Similarly,  $\leq_1 \cdot \leq_\phi$  means exactly that  $\psi \leq \phi \leq \phi$ , and this is equivalent to  $\psi \leq \phi$ .
- ⑨ (Vertical 2-Associativity) Associativity of the partial order on promodules directly gives the associativity of composition of 2-cells in ProMod.
- ⑩ (Horizontal 2-Identity) Let  $\psi, \phi : (X, A) \rightarrow (Y, B)$  be promodules. For any 2-cell  $\leq : \psi \rightarrow \phi$ , need to show that the 2-cell given by the horizontal composition,  $\leq * \leq_A$  is equal to  $\leq$ , as well as equal to  $\leq_B * \leq$ . So, it's required that  $\psi.A \leq \phi.A \iff \psi \leq \phi \iff B.\psi \leq B.\phi$ . And this holds as a direct consequence of (f).

# Proof: Promod is a 2-category III

- 11 (Horizontal 2-Associativity) As there's a unique 2-cell between any two promodules, and composition of promodules is associative, horizontal composition of 2-cells is associative.
- 12 (2-Identity) For promodules  $\psi : (X, A) \rightarrow (Y, B)$  and  $\phi : (Y, B) \rightarrow (Z, C)$  need  $(\leq_\psi * \leq_\phi) = \leq_{\psi \cdot \phi}$ . Both sides of the required equality are 2-cells  $\leq : \psi \cdot \phi \rightarrow \psi \cdot \phi$ . Thus, they are equal by the uniqueness of 2-cells between any two 1-cells.
- 13 (2-Interchange) Let  $\psi, \phi, \delta : (X, A) \rightarrow (Y, B)$  and  $\psi', \phi', \delta' : (Y, B) \rightarrow (Z, C)$  be promodules. For 2-cells  $\leq_1 : \psi \rightarrow \phi, \leq_2 : \phi \rightarrow \delta, \leq_a : \psi' \rightarrow \phi'$  and  $\leq_b : \phi' \rightarrow \delta'$ , need to show  $(\leq_b \cdot \leq_a) * (\leq_2 \cdot \leq_1) = (\leq_b * \leq_2) \cdot (\leq_a * \leq_1)$ . Both RHS and LHS are 2-cells from  $\psi \cdot \psi'$  to  $\delta \cdot \delta'$  and are hence equal.

# Proof: QUnif is a category I

- ① (Associativity) The composition of functions is associative by definition.
- ② (Identity) For each object  $(X, A)$ , the identity function  $\Delta_X : (X, A) \rightarrow (X, A)$  is uniformly continuous as  $\Delta_X.A = A \leq A = A.\Delta_X$ .

# Proof: Covariant Functor I

- (Partial-Order) Inclusion of relations acts as the partial order.
- (Down-Directed) Fix any  $k, k'$  belonging to  $B.f$ . Thus, there exist  $b, b'$  in  $B$  such that  $k = b f$  and  $k' = b' f$ . Using down-directedness of  $B$ , there exists  $c \in B$  such that  $c \subseteq b, b'$ . Hence, by Lemma 3.10,  $c f \subseteq k, k'$ .
- (Up-set) Let  $k$  belong to  $B.f$  and  $l : (X, A) \rightarrow (Y, B)$  be a uniformly continuous function such that  $l \supseteq k$ . Define a relation  $b' = \{(f(d), l(d)) : d \in \text{Dom}(l)\}$ . By definition, for any  $x \in X$  and  $z \in Y$  such that  $(x, z) \in l$ , we get that  $(f(x), z) \in b'$ . And  $l \supseteq k = b f$  implies  $\text{Dom}(l) \supseteq \text{Dom}(f)$ , giving  $(x, f(x)) \in f$ . Therefore, by definition of composition,  $(x, z) \in b'.f$ . Conversely, suppose  $(x, z) \in b'.f$ . By definition of composition, there exists  $f(x) \in Y$  such that  $(f(x), z) \in b'$ . Again using the definition of  $b'$ , we get that  $z = l(x)$  i.e.  $(x, z) \in l$ . Hence,  $l = b' f$ . Now we will show that  $b' \supseteq b$ . Because  $b' f = l \supseteq k = b f$ , for any  $x \in X$  we have that  $b'(f(x)) \supseteq b(f(x))$ . Thus,  $b'|_{f(x)} \supseteq b|_{f(x)}$ . By down-directedness of  $B$ ,  $b|_{f(x)} \subseteq b$  implies  $b(x)|_{f(x)} \in B$ . Finally,  $b' \supseteq b'|_{f(x)} \supseteq b|_{f(x)}$  gives  $b' \in B$ . Hence,  $b'.f \in B.f$ .
- Need to show that  $(B.f).A \leq B.f$ . So, fix any  $b \in B$ , we will find  $b' \in B$  and  $a \in A$  such that  $b' f a \subseteq b f$ . By quasi-uniformity of  $B$ , there exists  $b' \in B$  such that  $b' b' \subseteq b$ . Using Lemma 3.10, we get that  $b' b' f \subseteq b f$ . As  $f$  is uniformly continuous,  $f.A \leq B.f$  gives that there is some  $a \in A$  such that  $f a \subseteq b' f$ . Using this in the previous inequality, we get  $b' f a \subseteq b' b' f \subseteq b f$ .

## Proof: Covariant Functor II

- ⑤ Need to show that  $B.B.f \leq B.f$ . Fix any  $b \in B$ , we will find  $b' \in B$  such that  $b' b' f \subseteq b f$ . By quasi-uniformity of  $B$ , there exists  $b \in B$  such that  $b' b' \subseteq b$ . Using Lemma 3.10, we get  $b' b' f \subseteq b f$ .

Thus,  $B.f$  is a promodule. We now proceed to show that  $(\_)_{*}$  defines a functor.

- ① (Composition) Need to show that  $(g \circ f)_{*} = g_{*} f_{*}$  i.e.  $C.g.f = C.g.B.f$ .  
 In order to show  $C.g.f \leq C.g.B.f$ , fix any  $b \in B, c \in C$ . We will show that  $c g f \subseteq c g b f$ . As  $f$  is uniformly continuous,  $f.A \leq B.f$  gives that there exists  $a \in A$  such that  $f a \subseteq b f$ . Using Lemma 3.9, we get  $(c g) f a \subseteq (c g) b f$ . Now, using reflexivity of  $a$ , we get  $c g f \subseteq c g b f$ .  
 Now, to show that  $C.g.f \geq C.g.B.f$ . Fix any  $c \in C$ , we will find  $c' \in C$  and  $b \in B$  such that  $c g f \supseteq c g b f$ . By quasi-uniformity of  $C$ , there exists  $c' \in C$  such that  $c \supseteq c' c'$ . Using Lemma 3.10 gives that  $c(g f) \supseteq c' c' (g f)$ . Because  $g$  is uniformly continuous,  $C.g \geq g.B$  gives us  $b \in B$  such that  $c' g \supseteq b g$ . Using this in the previous inequality gives that  $c g f \supseteq c' g b f$ .
- ② (Identity) Let  $(X, A)$  be an object of QUnif and  $1_{(X,A)} : (X, A) \rightarrow (X, A)$  be the identity of  $(X, A)$ . That is,  $1_{(X,A)}$  is defined as  $x \mapsto x$ . Need to show that  $(1_{(X,A)})_{*} = 1_{(X,A)*}$ . Using functor's definition,  
 $LHS = (1_{(X,A)})_{*} = A.(1_{(X,A)}) = A.1_{(X,A)} = A$  and  $RHS = 1_{(X,A)*} = 1_{(X,A)}$   
 Using Proposition 4.8 (f), we get that  $A = 1_{(X,A)} = RHS$ .

# Proof: Contravariant Functor I

Showing that  $f^\circ.B : (Y, B) \rightarrow (X, A)$  is a promodule.

So, need to show  $f^\circ.B$  a prorelation  $Y \rightarrow X$  and that  $(f^\circ.B).B \leq f^\circ.B$  and  $A.(f^\circ.B) \leq f^\circ.B$

To show prorelation,

- ① (Partial-order) Inclusion of relations is the partial order.
- ② (Down directed) for  $k, k' \in f^\circ.B$ , need that  $\exists l \in f^\circ.B$  such that  $l \subseteq k, k'$   
 Fix  $k, k' \in f^\circ.B \implies \exists b, b' \in B : k = f^\circ \circ b$  and  $k' = f^\circ \circ b'$   
 By down-directedness of  $B$ , there exists  $c \in B$  such that  $c \subseteq b, b'$ , define  $l = f^\circ \circ c$ . Now, using Lemma 3.9 gives  $l = f^\circ \circ c \subseteq k, k'$ .
- ③ (Up-set) for a relation  $l : Y \rightarrow X$  and  $k \in f^\circ.B$  such that  $l \supseteq k$ , need  $l \in f^\circ.B$   
 Let  $b \in B$  be such that  $k = f^\circ \circ b$  and define  
 $b' := \{(y, y') : y \in \text{Dom}(l) \text{ and } y' \in (f^\circ)^{-1}(l(y))\}$   
 As  $l \supseteq k = f^\circ \circ b$ ,  $\text{Dom}(b') = \text{Dom}(l) \supseteq \text{Dom}(b)$   
 and  $\text{Ran}(l) \supseteq \text{Ran}(f^\circ \circ b) \implies \forall y \in \text{Dom}(b), \text{Ran}(b') = (f^\circ)^{-1}(l(y)) \supseteq (f^\circ)^{-1}(f^\circ \circ b) = \text{Ran}(b)$ .  
 Now, by definition of  $b'$ ,  $f^\circ \circ b' \supseteq l$ . To show  $f^\circ \circ b' \subseteq l$ ,  
 $(x, y) \in f^\circ \circ b' \implies \exists z \in Y : (x, z) \in b' \text{ and } (z, y) \in f^\circ \implies x \in \text{Dom}(l) \text{ and } z \in l(x) \text{ i.e. } (x, z) \in l$ .

# Proof: Contravariant Functor II

- ④ To show  $(f^\circ.B).B \leq f^\circ.B$ , need that  $\forall b \in B, \exists b' \in B : f^\circ \circ b' \circ b' \subseteq f^\circ \circ b$ ,  
 Fix any  $b \in B$ , as  $B$  is a quasi-uniformity,  
 $\exists b' \in B : b' \circ b' \subseteq b \implies f^\circ \circ b' \circ b' \subseteq f^\circ \circ b$ .  
 To show  $A.(f^\circ.B) \leq f^\circ.B$ , need that  $\forall b \in B$ ,  
 $\exists b' \in B, a \in A : a \circ f^\circ \circ b' \subseteq f^\circ \circ b$ .  
 As  $f$  is uniformly continuous,  $f.A \leq B.f$  i.e.  
 $\forall b \in B, \exists a \in A : f \circ a \subseteq b \circ f \implies a = f^\circ \circ f \circ a \subseteq f^\circ \circ b \circ f$ .  
 Fix any  $b \in B$ , so,  $\exists b' \in B : b' b' \subseteq b$ . And, for this  
 $b', \exists a : a \subseteq f^\circ b' f \implies a f^\circ b' \subseteq f^\circ b' f f^\circ b' \subseteq f^\circ b' b' \subseteq f^\circ b \implies a f^\circ b' \subseteq f^\circ b$ .

Now, need to show that  $(.)^*$  respects composition and identity.

- ① (Composition) let  $f, g$  be uniformly continuous,  $(X, A) \xrightarrow{f} (Y, B) \xrightarrow{g} (Z, C)$  need that  $(g \circ f)^* = f^*.g^*$   
 LHS  $= (g \circ f)^* = (g \circ f)^\circ.C = (f^\circ \circ g^\circ).C$  and RHS  $= f^*.g^* = (f^\circ.B).(g^\circ.C)$   
 For equality, showing that LHS  $\geq$  RHS and LHS  $\leq$  RHS:  
 To show  $(f^\circ \circ g^\circ).C \geq (f^\circ.B).(g^\circ.C)$ , need that  
 $\forall c \in C, \exists b \in B, c' \in C : f^\circ g^\circ c \supseteq f^\circ b g c'$   
 Fix any  $c \in C$ , so,  $\exists c' \in C : c' \circ c' \subseteq c \implies f^\circ g^\circ c \supseteq f^\circ g^\circ (c' c') = f^\circ g^\circ (c' \Delta_Z c') \supseteq f^\circ g^\circ c' (g g^\circ) c'$   
 By uniform continuity of  $g$ , for  $c' \in C, \exists b \in B : g b \subseteq c' g$   
 Thus,  $f^\circ g^\circ c \supseteq f^\circ g^\circ (c' g) g^\circ c' \supseteq f^\circ (g^\circ g) b g^\circ c' = f^\circ b g^\circ c'$ .  
 To show  $(f^\circ \circ g^\circ).C \leq (f^\circ.B).(g^\circ.C)$ , need that  
 $\forall b \in B, c \in C, \exists c' \in C : f^\circ g^\circ c \subseteq f^\circ b g^\circ c$



# Proof: Contravariant Functor III

Fix any  $c \in C, b \in B$  will show that  $c' := c$  works:

As  $B$  is a quasi-uniformity,

$$\Delta_Y \subseteq b \implies f^\circ \Delta_Y g^\circ c = f^\circ g^\circ c \subseteq f^\circ b g^\circ c = f^\circ b g^\circ c'$$

- ② (Identity) let  $(X, A) \in \mathbf{QUnif}^{op}$ , and  $1_{(X,A)} : (X, A) \rightarrow (X, A)$  as  $x \mapsto x$  need that  $(1_{(X,A)})^* = 1_{(X,A)}^*$  LHS  $= (1_{(X,A)})^* = (1_{(X,A)})^\circ \cdot A = 1_{(X,A)} \cdot A = A$ .  
And as  $RHS = 1_{(X,A)}^* = 1_{(X,A)}$  Using Proposition 3.2(f), we get that  $A = 1_{(X,A)} = RHS$ .

# Proof: Composition of Prorelations is a prorelation I

For prorelations  $P : X \rightarrow Y$  and  $Q : Y \rightarrow Z$ , need to show that  $Q.P$  is a prorelation.

- ① (Partial Order) Inclusion of relations gives a partial order.
- ② (Down-Directed) If  $k, k' \in Q.P$ , then  $k = q.p$  and  $k' = q'.p'$  for some  $q, q' \in Q$  and  $p, p' \in P$ . Because  $Q$  and  $P$  are prorelations, and hence down-directed sets there exists,  $a \in Q$  such that  $a \subseteq q, q'$  and  $b \in P$  such that  $b \subseteq p, p'$ . Thus, giving an element,  $a \circ b$  of  $Q.P$  such that  $a \circ b \subseteq k, k'$ .
- ③ (Up-Set) Let  $I : X \rightarrow Z$  be a relation, and  $k \in Q.P$  such that  $I \supseteq k$ . Define relations  $p : X \rightarrow Y$  and  $q : Y \rightarrow Z$  as,  $p = \{(x, y) : x \in \text{Dom}(I) \text{ and } y \in Y\}$  and  $q = \{(y, z) : y \in Y \text{ and } z \in \text{Ran}(I)\}$ . Because  $k \in Q.P$ , there exist  $q' \in Q$  and  $p' \in P$  such that  $k = q' \circ p'$ . Thus by definition of  $p$  and  $q$ , we get that  $p \supseteq p'$  and  $q \supseteq q'$ . Hence  $p \in P$  and  $q \in Q$  because  $P$  and  $Q$  are up-sets, which gives us that  $q \circ p \in Q.P$ . For any  $(x, z) \in I$ , by definition of  $p$  and  $q$ , we get that for every  $y \in Y$ ,  $(x, y) \in p$  and  $(y, z) \in q$ . By definition of composition, this gives that  $(x, z) \in q \circ p$ , giving that  $I \subseteq q \circ p$ . And, by definition of  $q \circ p$  we get that  $I \supseteq q \circ p$ . Finally giving that  $I = q \circ p \in Q.P$ .

## Topology induced by a quasi-uniformity

A subfamily  $\mathbb{B}$  of quasi-uniformity  $A$  is called a base for  $A$  if each relation in  $A$  contains a relation in  $\mathbb{B}$ .

### Proposition

*Let  $\mathbb{B}$  be the base for quasi-uniformity  $A$  on  $X$ . For  $x \in X$ , define  $\mathbb{B}(x) = \{B(x) | B \in \mathbb{B}\}$ . Then there is a unique topology on  $X$  such that for each  $x \in X$ ,  $\mathbb{B}(x)$  is a base for the neighborhood of  $x$  in this topology.*

We skip the proof as we have no requirement of it. But refer the interested reader to [2] for similar results.