

Q16 | Correct option: C

Source: Meeusen et al., ECSS/ACSM consensus statement on overtraining syndrome — PDF page 1; PubMed abstract (AB field) for PMID 23247672

PMID— 23247672

OWN — NLM

STAT— MEDLINE

DCOM— 20130517

LR — 20260128

IS — 1530-0315 (Electronic)

IS — 0195-9131 (Linking)

VI — 45

IP — 1

DP — 2013 Jan

TI — Prevention, diagnosis, and treatment of the overtraining syndrome: joint consensus statement of the European College of Sport Science and the American College of Sports Medicine.

PG — 186-205

LID — 10.1249/MSS.0b013e318279a10a [doi]

AB — Successful training not only must involve overload but also must avoid combination of excessive overload plus inadequate recovery. Athletes can experience short-term performance decrement without severe psychological lasting other negative symptoms. This functional overreaching will even lead to an improvement in performance after recovery. When athletes do sufficiently respect the balance between training and recovery, nonfunctional overreaching (NFOR) can occur. The distinction between NFOR and overtraining syndrome (OTS) is very difficult and will depend on the clinical outcome exclusion diagnosis. The athlete will often show the same clinical, hormonal other signs and symptoms. A keyword in the recognition of OTS might be "maladaptation" not only of the athlete but also of several biological, neurochemical, and hormonal regulation mechanisms. It is generally thought symptoms of OTS, such as fatigue, performance decline, and mood disturbance more severe than those of NFOR. However, there is no scientific evidence either confirm or refute this suggestion. One approach to understanding etiology of OTS involves the exclusion of organic diseases or infectious factors such as dietary caloric restriction (negative energy balance) and insufficient carbohydrate and/or protein intake, iron deficiency, magnesium deficiency, allergies, and others together with identification of initiating events or triggers. In this article, we provide the recent status of potential markers for the detection of OTS. Currently, several markers (hormones, performance tests, psychological tests, and biochemical and immune markers) are used, but none of them meet all the criteria to make their use generally accepted.

FAU — Meeusen, Romain

AU — Meeusen R

FAU — Duclos, Martine

AU — Duclos M

FAU — Foster, Carl

AU — Foster C

FAU — Fry, Andrew

AU — Fry A

FAU — Gleeson, Michael

AU — Gleeson M

FAU — Nieman, David