

Indian Institute of Technology Patna

Physics Department

PH 201: Tutorial (Lasers)

1. Find phase and group velocities of an electron whose de-Broglie's wavelength is 14 nm.
2. A silicon photo diode has quantum efficiency of 65% with photon energy 1.5×10^{-19} J. Its bandgap energy is 0.67 eV. Calculate
 - (a) long wavelength cut off (λ_c),
 - (b) responsivity (R), and
 - (c) incident power required to obtain a photo current $2.5 \mu A$?
3. Find how many photons are required per second to produce laser beam of 3 mW power, when wavelength of laser is 694.3 nm.
4. Calculate the frequency in Hertz and wavenumbers (cm^{-1}) and the energy in electronvolts of a photon of wavelength $\lambda = 1 \mu\text{m}$ in vacuum.
5. Calculate the wavenumbers corresponding to an energy spacing of kT , where k is the Boltzmann constant and T is the absolute temperature. Assume $T = 300$ K.
6. The brightness of probably the brightest lamp so far available (PEK Labs type 107/109TM, excited by 100 W of electrical power) is about $95 \text{ W/cm}^2 \text{ sr}$ in its most intense green line ($\lambda = 546 \text{ nm}$). Compare this brightness with that of a 1 W Argon laser ($\lambda = 514.5 \text{ nm}$), which can be assumed to be diffraction limited.
7. An interference filter with a pass band of 10 nm centered at 500 nm is used to obtain approximately monochromatic light from a white source. Calculate the coherence time of the filtered light and its coherence length.
8. When 3×10^{11} photons each with wavelength of $0.85 \mu\text{m}$ are incident on a photo diode, on average 1.2×10^{11} electrons are generated. Determine quantum efficiency and responsivity.

9. Determine the ABCD matrix for a beam translated a distance d_1 , focused through a thin lens of focal length f_1 , and then translated a distance d_2 .
10. Determine the saturation energy of Nd:YAG laser operating at wavelength 1.06 μm . Given that cross-section $\sigma^H = 1.14 \times 10^{-22} \text{ m}^2$.
11. At the surface of the earth, the intensity of the sun is approximately 1 kW m^{-2} . Calculate the intensity at the retina that results when looking directly at the sun. Assume that: (i) the pupil of a bright-adapted eye is 2 mm in diameter; (ii) the focal length of the eye is 22.5 mm; (iii) the Sun subtends an angle of 0.5° . Compare this intensity with that resulting when looking into a 1 mW He-Ne laser ($\lambda = 632.8 \text{ nm}$) with a 2 mm diameter.
12. Light travelling in air strikes a glass plate at a glancing angle 33° . While striking the glass plate, part of the beam is reflected and part is refracted. If the refracted and reflected beams make an angle 90° , with each other, then
 - (a) what is the refractive index of the glass and
 - (b) what is the critical angle for that glass?
13. The numerical aperture of an optical fiber is 0.5 and core refractive index 1.54.
 - (a) Find refractive index of cladding.
 - (b) Calculate the change in core cladding refractive index per unit refractive index of core and
 - (c) Critical angle.
14. Relative refractive index of a fiber is 5.5% when core refractive index is 1.48. Find numerical aperture, cladding refractive index, acceptance angle, and velocity of light in core and cladding.
