

MA 201, Mathematics III, July-November 2016, Fourier Transform (Contd.)

Lecture 17

Fourier Transform

For a given function f , the improper integral

$$\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} f(\tau) e^{-i\sigma\tau} d\tau \right] d\sigma = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g(\sigma) d\sigma$$

is called Fourier integral of f .

If f is continuous at t , we have

$$f(t) = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g(\sigma) d\sigma.$$

The function g is called *Fourier transform* of a function f and it is denoted by $\mathcal{F}\{f(t)\}$

$$\mathcal{F}\{f(t)\} = g(\sigma) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma\tau} f(\tau) d\tau. \quad (1)$$

The *inverse Fourier transform* is defined as

$$f(t) = \mathcal{F}^{-1}\{g(\sigma)\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g(\sigma) e^{i\sigma t} d\sigma. \quad (2)$$

Fourier cosine transform

For an even function f defined in $(-\infty, \infty)$, the Fourier integral

$$\begin{aligned}f(t) &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} f(\tau) e^{-i\sigma\tau} d\tau \right] d\sigma \quad (\text{Basic definition}) \\&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g_c(\sigma) d\sigma \\&= \frac{2}{\pi} \int_0^{\infty} \int_0^{\infty} f(\tau) \cos \sigma\tau \cos \sigma t d\tau d\sigma \quad (\text{Special case}) \\&= \int_0^{\infty} \sqrt{\frac{2}{\pi}} \cos \sigma t \left[\int_0^{\infty} \sqrt{\frac{2}{\pi}} f(\tau) \cos \sigma\tau d\tau \right] d\sigma \\&= \int_0^{\infty} \sqrt{\frac{2}{\pi}} \cos \sigma t g_c(\sigma) d\sigma\end{aligned}$$

We can define the *Fourier cosine transform* of f as

$$\begin{aligned}\mathcal{F}_c\{f(t)\} = g_c(\sigma) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma\tau} f(\tau) d\tau \quad (\text{From Basic definition}) \\&= \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(\tau) \cos \sigma\tau d\tau. \quad (\text{From Special case})\end{aligned}$$

The *inverse Fourier cosine transform* is defined as

$$\begin{aligned}\mathcal{F}_c^{-1}\{g_c(\sigma)\} = f(t) &= \sqrt{\frac{2}{\pi}} \int_0^{\infty} g_c(\sigma) \cos \sigma t d\sigma. \quad (\text{From Special case}) \\&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g_c(\sigma) d\sigma. \quad (\text{From Basic definition})\end{aligned}$$

Fourier sine transform

For an odd function f defined in $(-\infty, \infty)$, the Fourier integral

$$\begin{aligned}f(t) &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} \left[\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} f(\tau) e^{-i\sigma\tau} d\tau \right] d\sigma \quad (\text{Basic definition}) \\&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g_s(\sigma) d\sigma \\&= \frac{2}{\pi} \int_0^{\infty} \int_0^{\infty} f(\tau) \sin \sigma\tau \sin \sigma t d\tau d\sigma \quad (\text{Special case}) \\&= \int_0^{\infty} \sqrt{\frac{2}{\pi}} \sin \sigma t \left[\int_0^{\infty} \sqrt{\frac{2}{\pi}} f(\tau) \sin \sigma\tau d\tau \right] d\sigma \\&= \int_0^{\infty} \sqrt{\frac{2}{\pi}} \sin \sigma t g_s(\sigma) d\sigma\end{aligned}$$

We can define the *Fourier sine transform* of f as

$$\begin{aligned}\mathcal{F}_s\{f(t)\} = g_s(\sigma) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma\tau} f(\tau) d\tau \quad (\text{From Basic definition}) \\&= \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(\tau) \sin \sigma\tau d\tau. \quad (\text{From Special case})\end{aligned}$$

The *inverse Fourier sine transform* is defined as

$$\begin{aligned}\mathcal{F}_s^{-1}\{g_s(\sigma)\} = f(t) &= \sqrt{\frac{2}{\pi}} \int_0^{\infty} g_s(\sigma) \sin \sigma t d\sigma. \quad (\text{From Special case}) \\&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{i\sigma t} g_s(\sigma) d\sigma. \quad (\text{From Basic definition})\end{aligned}$$

Example

Consider $f(t) = e^{-t}$, $t > 0$

Justify that

- (a) f has Fourier sine and cosine transformations.
- (b) Find Fourier cosine transformation of f .
- (c) Use part (b) to evaluate following improper integral

$$\int_0^{\infty} \frac{\cos \sigma t}{\sigma^2 + 1} d\sigma, \quad t > 0.$$

(a) Use even and odd extensions of f in $(-\infty, \infty)$.

(b) Suppose f_e is the even extension of f in $(-\infty, \infty)$. Then

$$\begin{aligned} \mathcal{F}_c\{f_e(t)\} = g_c(\sigma) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma\tau} f_e(\tau) d\tau \quad (\text{From Basic definition}) \\ &= \frac{1}{\sqrt{2\pi}} \frac{2}{1 + \sigma^2} \end{aligned}$$

Example (Contd.)

(c) Finally, use inverse Fourier cosine transform to have

$$\begin{aligned} f_e(t) = \mathcal{F}_c^{-1}\{g_c(\sigma)\} &= \sqrt{\frac{2}{\pi}} \int_0^\infty g_c(\sigma) \cos \sigma t \, d\sigma. \quad (\text{From Special case}) \\ &= \int_0^\infty \frac{2 \cos \sigma t}{\pi \sigma^2 + 1} d\sigma. \quad (\text{From part (b)}) \end{aligned}$$

What about following examples?

$$(a) \int_0^\infty \frac{\sigma^3 \sin \sigma t}{\sigma^4 + 4} d\sigma, \quad t > 0.$$

$$(b) \int_0^\infty \frac{\sin \sigma t}{\sigma} d\sigma, \quad t > 0.$$

Here, f is not given explicitly.

Convolution

Convolution Theorem

Theorem

If $h(t)$ is the response obtained by convolving the function $f_1(t)$ with $f_2(t)$, then

$$\mathcal{F}\{h(t)\} = \mathcal{F}\{f_1 * f_2\} = \mathcal{F}\{f_1(t)\}\mathcal{F}\{f_2(t)\}.$$

In other words,

we can say that the Fourier transform of the convolution of $f_1(t)$ and $f_2(t)$ is the product of the Fourier transforms of $f_1(t)$ and $f_2(t)$.

Recall: The convolution of two functions $f_1(t)$ and $f_2(t)$, $-\infty < t < \infty$, is defined as

$$f_1(t) * f_2(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau.$$

Convolution

Proof: By definition

$$\begin{aligned}\mathcal{F}\{h(t)\} = \mathcal{F}\{f_1 * f_2\} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (f_1(t) * f_2(t)) e^{-i\sigma t} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau \right] e^{-i\sigma t} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} f_1(\tau) \left[\int_{-\infty}^{\infty} f_2(t - \tau) e^{-i\sigma t} dt \right] d\tau.\end{aligned}$$

Putting $t - \tau = \omega$ in the inner integral

$$\begin{aligned}\mathcal{F}\{h(t)\} = \mathcal{F}\{f_1 * f_2\} &= \frac{1}{2\pi} \int_{-\infty}^{\infty} f_1(\tau) \left[\int_{-\infty}^{\infty} f_2(\omega) e^{-i\sigma(\omega + \tau)} d\omega \right] d\tau \\ &= \left(\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_1(\tau) e^{-i\sigma\tau} d\tau \right) \left(\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_2(\omega) e^{-i\sigma\omega} d\omega \right) \\ &= \mathcal{F}\{f_1\} \mathcal{F}\{f_2\}.\end{aligned}$$

This is known as the *convolution integral*.

Convolution

Example:

A system has the impulse response given by $f_1(t)$ and is subjected to the rectangular pulse given by $f_2(t)$ as follows:

$$f_1(t) = \begin{cases} 0, & -\infty < t \leq 0, \\ ae^{-at}, & 0 < t < \infty, \quad a > 0 \end{cases}$$
$$f_2(t) = \begin{cases} 0, & -\infty < t \leq -1, \\ 1, & -1 < t \leq 1, \\ 0, & 1 < t < \infty. \end{cases}$$

Find the output time function $h(t)$ by convolving $f_1(t)$ with $f_2(t)$.

Convolution

Solution:

The Fourier transform of $f_1(t)$:

$$\begin{aligned}\mathcal{F}\{f_1(t)\} &= g_1(\sigma) \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_1(t) e^{-i\sigma t} dt \\&= \frac{1}{\sqrt{2\pi}} \int_0^{\infty} a e^{-at} e^{-i\sigma t} dt \\&= \frac{a}{\sqrt{2\pi}} \left| \frac{e^{-(a+i\sigma)t}}{-(a+i\sigma)} \right|_0^{\infty} \\&= \frac{a}{\sqrt{2\pi}(a+i\sigma)}.\end{aligned}$$

The Fourier transform of $f_2(t)$:

$$\begin{aligned}\mathcal{F}\{f_2(t)\} &= g_2(\sigma) \\&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f_2(t) e^{-i\sigma t} dt \\&= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 e^{-i\sigma t} dt \\&= \frac{1}{\sqrt{2\pi}} \left| \frac{e^{-i\sigma t}}{-i\sigma} \right|_{-1}^1 \\&= \frac{1}{\sqrt{2\pi}} \frac{e^{i\sigma} - e^{-i\sigma}}{i\sigma} \\&= \sqrt{\frac{2}{\pi}} \frac{\sin \sigma}{\sigma}.\end{aligned}$$

Convolution

Therefore

$$\mathcal{F}\{f_1(t)\}\mathcal{F}\{f_2(t)\} = \frac{1}{\pi} \frac{a \sin \sigma}{\sigma(a + i\sigma)}$$

According to convolution theorem:

$$\mathcal{F}\{h(t)\} = \mathcal{F}\{f_1 * f_2\} = \mathcal{F}\{f_1(t)\}\mathcal{F}\{f_2(t)\}.$$

Hence

$$h(t) = \mathcal{F}^{-1}[\mathcal{F}\{f_1(t)\}\mathcal{F}\{f_2(t)\}] = \mathcal{F}^{-1}\left[\frac{1}{\pi} \frac{a \sin \sigma}{\sigma(a + i\sigma)}\right]$$

which will give us

$$h(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{\pi} \frac{a \sin \sigma}{\sigma(a + i\sigma)} e^{i\sigma t} d\sigma$$

which is the output time function.

Application of Fourier Transform to PDEs

As of now we have defined all variants of Fourier transform for a function of one variable only.

It is obvious that if we want to use Fourier transforms to solve PDEs, we need to define Fourier transform for a function of at least two variables.

Based on the definitions discussed earlier for a function of one variable, let us define Fourier transform of a function of two variables x and t and those of their partial derivatives

so that we can at least solve one-dimensional transient IBVPs governed by a PDE.

Application of Fourier Transform to PDEs

Fourier transform:

The Fourier transform of a function $U(x, t)$ with respect to x is defined as

$$\mathcal{F}\{U(x, t)\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma x} U(x, t) dx = \hat{U}(\sigma, t). \quad (3)$$

Inverse Fourier transform:

The inverse Fourier transform $U(x, t)$ of $\hat{U}(\sigma, t)$ is defined as

$$U(x, t) = \mathcal{F}^{-1}\{\hat{U}(\sigma, t)\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i\sigma x} \hat{U}(\sigma, t) d\sigma.$$

Under the assumption that U and $\frac{\partial U}{\partial x}$ vanish as $x \rightarrow \pm\infty$, we obtain the following results of transforms of the partial derivatives:

$$\begin{aligned} \mathcal{F}\left\{\frac{\partial}{\partial x}U(x, t)\right\} &= i\sigma\hat{U}(\sigma, t), \\ \mathcal{F}\left\{\frac{\partial^2}{\partial x^2}U(x, t)\right\} &= (i\sigma)^2\hat{U}(\sigma, t), \\ \mathcal{F}\left\{\frac{\partial}{\partial t}U(x, t)\right\} &= \frac{d}{dt}\hat{U}(\sigma, t), \\ \mathcal{F}\left\{\frac{\partial^2}{\partial t^2}U(x, t)\right\} &= \frac{d^2}{dt^2}\hat{U}(\sigma, t). \end{aligned}$$

Application of Fourier Transform to PDEs

Result I:

$$\mathcal{F}\left\{\frac{\partial}{\partial x}U(x,t)\right\} = i\sigma\hat{U}(\sigma,t). \quad (4)$$

Proof:

$$\begin{aligned}\mathcal{F}\left\{\frac{\partial}{\partial x}U(x,t)\right\} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma x} \frac{\partial U}{\partial x} dx \\ &= \frac{1}{\sqrt{2\pi}} \left\{ [e^{-i\sigma x} U(x,t)]_{-\infty}^{\infty} + i\sigma \int_{-\infty}^{\infty} e^{-i\sigma x} U(x,t) dx \right\} \\ &= i\sigma\hat{U}(\sigma,t).\end{aligned}$$

□

Result II:

$$\mathcal{F}\left\{\frac{\partial^2}{\partial x^2}U(x,t)\right\} = (i\sigma)^2\hat{U}(\sigma,t). \quad (5)$$

Proof:

$$\begin{aligned}\mathcal{F}\left\{\frac{\partial^2}{\partial x^2}U(x,t)\right\} &= \frac{1}{\sqrt{2\pi}} \left\{ \int_{-\infty}^{\infty} e^{-i\sigma x} \frac{\partial^2 U}{\partial x^2} dx \right\} \\ &= \frac{1}{\sqrt{2\pi}} \left\{ [e^{-i\sigma x} \frac{\partial U}{\partial x}]_{-\infty}^{\infty} + i\sigma \int_{-\infty}^{\infty} e^{-i\sigma x} \frac{\partial U}{\partial x} dx \right\} \\ &= \frac{1}{\sqrt{2\pi}} \left\{ i\sigma ([e^{-i\sigma x} U(x,t)]_{-\infty}^{\infty} + i\sigma \int_{-\infty}^{\infty} e^{-i\sigma x} U(x,t) dx) \right\} \\ &= (i\sigma)^2\hat{U}(\sigma,t).\end{aligned}$$

□

Application of Fourier Transform to PDEs

Result III:

$$\mathcal{F}\left\{\frac{\partial}{\partial t}U(x,t)\right\} = \frac{d}{dt}\hat{U}(\sigma,t). \quad (6)$$

Proof:

$$\begin{aligned}\mathcal{F}\left\{\frac{\partial}{\partial t}U(x,t)\right\} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-i\sigma x} \frac{\partial U}{\partial t} dx \\ &= \frac{1}{\sqrt{2\pi}} \frac{\partial}{\partial t} \int_{-\infty}^{\infty} e^{-i\sigma x} U(x,t) dx \\ &= \frac{1}{\sqrt{2\pi}} \frac{d}{dt} \int_{-\infty}^{\infty} e^{-i\sigma x} U(x,t) dx \\ &= \frac{d}{dt} \hat{U}(\sigma,t).\end{aligned}$$

□

Proceeding in a similar manner as above, Result 4 as given below can be obtained:

$$\mathcal{F}\left\{\frac{\partial^2}{\partial t^2}U(x,t)\right\} = \frac{d^2}{dt^2}\hat{U}(\sigma,t). \quad (7)$$

□

The use of Fourier transform to solve partial differential equations is best described by examples.

Application of Fourier Transform to PDEs

Example A:

An infinitely long string extending in $-\infty < x < \infty$ under uniform tension is displaced into the curve $y = f(x)$ and let go from rest with velocity $g(x)$. To find the displacement $U(x, t)$ at any point at any subsequent time.

Solution:

The boundary value problem is

$$\frac{\partial^2 U}{\partial t^2} = c^2 \frac{\partial^2 U}{\partial x^2}, \quad -\infty < x < \infty, \quad t > 0, \quad (8)$$

$$U(x, 0) = f(x) \quad (\text{initial displacement}), \quad (9)$$

$$\frac{\partial U}{\partial t}(x, 0) = g(x) \quad (\text{initial velocity}). \quad (10)$$

Taking Fourier transform on both sides of PDE (8),

$$\frac{d^2}{dt^2} \hat{U}(\sigma, t) = -c^2 \sigma^2 \hat{U}(\sigma, t), \quad (11)$$

Application of Fourier Transform to PDEs

(11) can be written in standard form as

$$\frac{d^2}{dt^2}\hat{U}(\sigma, t) + c^2\sigma^2\hat{U}(\sigma, t) = 0. \quad (12)$$

On solving, we get

$$\hat{U}(\sigma, t) = A(\sigma)\cos(c\sigma t) + B(\sigma)\sin(c\sigma t). \quad (13)$$

Taking Fourier transforms on the initial conditions (9) and (10),

$$\hat{U}(\sigma, 0) = \hat{f}(\sigma), \quad (14)$$

$$\frac{d}{dt}\hat{U}(\sigma, 0) = \hat{g}(\sigma), \quad (15)$$

where $\hat{f}(\sigma)$ and $\hat{g}(\sigma)$, are, respectively the Fourier transform of $f(x)$ and $g(x)$.

Using the initial conditions, $A(\sigma)$ and $B(\sigma)$ can be obtained as:

$$\begin{aligned} \hat{U}(\sigma, 0) &= A(\sigma) = \hat{f}(\sigma), \\ \frac{d}{dt}\hat{U}(\sigma, 0) &= c\sigma B(\sigma) = \hat{g}(\sigma). \end{aligned}$$

Application of Fourier Transform to PDEs

Now

$$\hat{U}(\sigma, t) = \hat{f}(\sigma) \cos(c\sigma t) + \frac{1}{c\sigma} \hat{g}(\sigma) \sin(c\sigma t). \quad (16)$$

To get the solution, we use the inverse Fourier transform to obtain

$$U(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \left[\hat{f}(\sigma) \cos(c\sigma t) + \frac{1}{c\sigma} \hat{g}(\sigma) \sin(c\sigma t) \right] e^{i\sigma x} d\sigma. \quad (17)$$

Formula (17) gives us the solution of the wave boundary value problem in the form of an integral involving the Fourier transform of the initial displacement and velocity.

Application of Fourier Transform to PDEs

We summarize the Fourier transform method as follows:

Step 1: Take Fourier transform on the given boundary value problem in $U(x, t)$ and get an ordinary differential equation in $\hat{U}(\sigma, t)$ in the variable t .

Step 2: Solve the ordinary differential equation and find $\hat{U}(\sigma, t)$.

Step 3: Take inverse Fourier transform to get $U(x, t)$

Application of Fourier Transform to PDEs

Example B:

Consider the heat conduction in an infinite rod with thermal diffusivity α with initial temperature distribution $f(x)$. To find the temperature distribution $U(x, t)$ at any point at any subsequent time.

Solution:

The boundary value problem is

$$\frac{\partial U}{\partial t} = \alpha \frac{\partial^2 U}{\partial x^2}, \quad -\infty < x < \infty, \quad t > 0, \quad (18)$$

$$U(x, 0) = f(x) \quad (\text{initial temperature distribution}). \quad (19)$$

Taking Fourier transform on both sides of PDE (18), we observe

$$\hat{U}(\sigma, t) = \hat{f}(\sigma)e^{-\alpha\sigma^2 t}, \quad (20)$$

where $\hat{f}(\sigma)$ is the Fourier transform of $f(x)$.

We use the inverse Fourier transform to obtain

$$U(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(\sigma)e^{-\alpha\sigma^2 t} e^{i\sigma x} d\sigma. \quad (21)$$

Application of Fourier Transform to PDEs

Example C: Semi-infinite rod with Dirichlet condition

Mathematical Model

The boundary value problem is the following:

$$\frac{\partial U}{\partial t} = \alpha \frac{\partial^2 U}{\partial x^2}, \quad x > 0, \quad t > 0, \quad (22)$$

$$U(x, 0) = f(x), \quad t > 0 \quad U(0, t) = U_0, \quad x > 0. \quad (23)$$

Example D: Semi-infinite rod with Neumann condition

Consider the same equation as in the previous example subject to the boundary conditions

$$\frac{\partial U}{\partial x}(0, t) = 0, \quad U(x, 0) = f(x)$$

The Fourier sine and cosine transforms can be employed to solve a partial differential equation when the range of the spatial variable extends from 0 to ∞ .

If the boundary condition is in terms of some value of $U(0, t)$ (i.e. Dirichlet boundary condition), then sine transform is to be used.

When the boundary condition is in terms of some value of $\frac{\partial U}{\partial x}(0, t)$ (i.e. Neumann boundary condition), then cosine transform is to be used.

Application of Fourier Transform to PDEs

The *Fourier sine transform* of a function $U(x, t)$ with respect to x is defined as

$$\mathcal{F}_s\{U(x, t)\} = \sqrt{\frac{2}{\pi}} \int_0^\infty U(x, t) \sin \sigma x \, dx = \hat{U}_s(\sigma, t). \quad (24)$$

Under the assumption that U and $\frac{\partial U}{\partial x}$ vanish as $x \rightarrow \infty$,

$$\mathcal{F}_s\left\{\frac{\partial U(x, t)}{\partial t}\right\} = \frac{d}{dt}\hat{U}_s(\sigma, t), \quad (25)$$

$$\mathcal{F}_s\left\{\frac{\partial^2 U(x, t)}{\partial t^2}\right\} = \frac{d^2}{dt^2}\hat{U}_s(\sigma, t), \quad (26)$$

$$\mathcal{F}_s\left\{\frac{\partial^2 U(x, t)}{\partial x^2}\right\} = \sqrt{\frac{2}{\pi}}\sigma U(0, t) - \sigma^2 \hat{U}_s(\sigma, t). \quad (27)$$

Application of Fourier Transform to PDEs

The *Fourier cosine transform* of a function $U(x, t)$ with respect to x is defined as

$$\mathcal{F}_c\{U(x, t)\} = \sqrt{\frac{2}{\pi}} \int_0^\infty U(x, t) \cos \sigma x \, dx = \hat{U}_c(\sigma, t). \quad (28)$$

Under the assumption that U and $\frac{\partial U}{\partial x}$ vanish as $x \rightarrow \infty$,

$$\mathcal{F}_c\left\{\frac{\partial U(x, t)}{\partial t}\right\} = \frac{d}{dt} \hat{U}_c(\sigma, t), \quad (29)$$

$$\mathcal{F}_c\left\{\frac{\partial^2 U(x, t)}{\partial t^2}\right\} = \frac{d^2}{dt^2} \hat{U}_c(\sigma, t), \quad (30)$$

$$\mathcal{F}_c\left\{\frac{\partial^2 U(x, t)}{\partial x^2}\right\} = -\sqrt{\frac{2}{\pi}} U_x(0, t) - \sigma^2 \hat{U}_c(\sigma, t). \quad (31)$$

Application of Fourier Transform to PDEs

Example C:

If $U(x, t)$ is the temperature at time t and α the thermal diffusivity of a semi-infinite metal bar, find the temperature distribution in the bar at any point at any subsequent time if the initial temperature distribution is given as $f(x)$ and the boundary is kept at U_0 degrees.

Solution:

The boundary value problem is the following:

$$\frac{\partial U}{\partial t} = \alpha \frac{\partial^2 U}{\partial x^2}, \quad x > 0, \quad t > 0, \quad (32)$$

$$U(x, 0) = f(x), \quad t > 0 \quad U(0, t) = U_0, \quad x > 0. \quad (33)$$

The boundary condition suggests that we need to use Fourier sine transform. Taking the transform on (32),

$$\frac{d}{dt} \hat{U}_s(\sigma, t) = \alpha \left[\sqrt{\frac{2}{\pi}} \sigma U(0, t) - \sigma^2 \hat{U}_s(\sigma, t) \right]. \quad (34)$$

Using the boundary condition,

$$\frac{d}{dt} \hat{U}_s(\sigma, t) + \alpha \sigma^2 \hat{U}_s(\sigma, t) = \alpha \sqrt{\frac{2}{\pi}} \sigma U_0. \quad (35)$$

Application of Fourier Transform to PDEs

On solving

$$\hat{U}_s(\sigma, t) = A(\sigma)e^{-\alpha\sigma^2 t} + \frac{1}{\sigma}\sqrt{\frac{2}{\pi}}U_0. \quad (36)$$

Using the initial condition,

$$\hat{f}_s(\sigma) = A(\sigma) + \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma},$$

where $\hat{f}_s(\sigma)$ is the Fourier sine transform of $f(x)$. that is,

$$A(\sigma) = \hat{f}_s(\sigma) - \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma}$$

Now $\hat{U}_s(\sigma, t)$ is

$$\begin{aligned} \hat{U}_s(\sigma, t) &= (\hat{f}_s(\sigma) - \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma})e^{-\alpha\sigma^2 t} + \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma} \\ &= \hat{f}_s e^{-\alpha\sigma^2 t} + \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma}(1 - e^{-\alpha\sigma^2 t}) \end{aligned} \quad (37)$$

The inversion gives

$$U(x, t) = \sqrt{\frac{2}{\pi}} \int_0^\infty \left[\hat{f}_s e^{-\alpha\sigma^2 t} + \sqrt{\frac{2}{\pi}}\frac{U_0}{\sigma}(1 - e^{-\alpha\sigma^2 t}) \right] \sin \sigma x \, d\sigma. \quad (38)$$

Application of Fourier Transform to PDEs

Similarly, we can try the vibration problem in a semi-infinite string, $0 < x < \infty$.

In this case, there will be a second-order ordinary differential equation due to the presence of the term $\frac{\partial^2 U}{\partial t^2}$.

The two initial conditions $U(x, 0)$ and $\frac{\partial U}{\partial t}(x, 0)$ will help us in determining the coefficients of $\hat{U}_c(\sigma, t)$ or $\hat{U}_s(\sigma, t)$ depending on the boundary condition at $x = 0$.