



first order in α if $\alpha << 1$.

INDIAN INSTITUTE OF TECHNOLOGY PATNA

[2+3]

MSE, PH 101 (PHYSICS I), Sept. 2012

Time: 2 hrs, Full Marks: 50

(Answer all questions. You should answer according to the marks allotted for each question)

	1. a) If the cosine of the angles between a vector and the x, y, z axes are α , β , γ , respectively. Prove that $\alpha^2 + \beta^2 + \gamma^2 = 1$. [2] b) Write down the three most important properties of a conservative force. [3] c) A rocket of mass M moves in space with a velocity \mathbf{v} . Consider the rocket at time t . Between $t \& t + \Delta t$ a mass of fuel Δm is burned and expelled as gas with velocity \mathbf{u} relative to the rocket. \mathbf{u} is independent of the velocity of the rocket. Derive the expression of the external force. [4]
	2. a) For a fixed axis rotation of a rigid body, show that the velocity is always perpendicular to the radius vector.b) Find the position of center of mass of a uniform solid hemisphere of radius R and mass M.
	c) An automobile enters a turn whose radius is R . The road is banked at angle θ , and the coefficient of friction between wheels and road is μ . Find the maximum and minimum speeds for the car to stay on the road without skidding sideways. [4]
	3. a) Prove that the force is everywhere perpendicular to the constant energy surfaces and points from higher to lower potential energy. [2]
	b) For a potential energy function: $U = \epsilon \left[\left(\frac{r_0}{r} \right)^{12} - 2 \left(\frac{r_0}{r} \right)^6 \right]$
	 i) What are the radius at the potential minimum and the depth of the potential well? [2] ii) Find the frequency of small oscillations about equilibrium for two identical atoms of mass m bound to each other inside the potential. [2]
	 4. a) Prove that, motion of a body under central force field always lies in a plane. Also show that the area swept out by the body per unit time is constant. [2+3] b) Derive the angular momentum of a system of particle involving both translation and rotation. [2] c) For a general rotation with an angular velocity w, around an arbitrary axis, prove that the velocity, v=w × r, where r is the radius vector from the origin of the coordinate system. [2]
	5. a) Explain the theory & stability of a Gyroscope? [3] b) A gyroscope wheel is at one end of an axle of length l . The other end of the axle is suspended from a string of length L . The wheel is set into motion so that it executes uniform precession in the horizontal plane. The wheel has mass M and moment of inertia about its center of mass I_0 . Its spin angular velocity is w_s . Neglect the mass of the shaft and of the string. Find the angle that the string makes with the vertical. Assume the angle is small. [4]
j	 6. a) Write down the expression of angular momentum for the general rotation of a rigid body and define inertia tensor. What do you mean by principal moment of inertia? [3+2] b) A particle of mass m is located at x = 2, y = 0, z = 3. i) Find the inertia tensor relative to the origin. ii) The particle undergoes pure rotation about the z axis through a small analysis.

about the z axis through a small angle α . Show that its moments of inertia are unchanged to