
Covariance Sensitivity Proofs

March 21, 2020

1 PRELIMINARIES

Lemma 1. $\forall i$,

$$\sum_{j=1}^n (x_{ji} - \bar{X}_i) = 0.$$

Proof.

$$\begin{aligned} \sum_{i=1}^n (x_{ji} - \bar{X}_i) &= \sum_{i=1}^n x_{ji} - n\bar{X}_i \\ &= \sum_{i=1}^n x_{ji} - n \left(\frac{1}{n} \sum_{i=1}^n x_{ji} \right) \\ &= 0. \end{aligned}$$

□

Lemma 2. *Let*

$$f_{ij} = \sum_{k=1}^n (x_{ki} - \bar{X}_i)(x_{kj} - \bar{X}_j),$$

and let $X'_i = X_i \cup \{y_i\}$, and say X_i has size n . Let \bar{X}'_i and \bar{X}'_j be the sample means of X'_i and X'_j respectively. Then,

$$f_{ij}(X') = f_{ij}(X) + n(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j) + (y_i - \bar{X}_i)(y_j - \bar{X}_j).$$

Proof. Note that

$$\begin{aligned}
f_{ij}(X') &= \sum_{k=1}^{n+1} (x'_{ki} - \bar{X}'_i)(x'_{kj} - \bar{X}'_j), \\
&= \sum_{k=1}^n (x_{ki} - \bar{X}'_i)(x_{kj} - \bar{X}'_j) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j), \\
&= \sum_{k=1}^n ((x_{ki} - \bar{X}_i) + (\bar{X}_i - \bar{X}'_i)) ((x_{kj} - \bar{X}_j) + (\bar{X}_j - \bar{X}'_j)) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j), \\
&= \sum_{k=1}^n (x_{ki} - \bar{X}_i)(x_{kj} - \bar{X}_j) + (\bar{X}_j - \bar{X}'_j) \sum_{k=1}^n (x_{ki} - \bar{X}_i) + (\bar{X}_i - \bar{X}'_i) \sum_{k=1}^n (x_{kj} - \bar{X}_j), \\
&\quad + \sum_{k=1}^n (\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j), \\
&= f_{ij}(X) + n(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j),
\end{aligned}$$

where the cancellation of the second and third terms in the second-to-last line is due to Lemma 1. \square

Lemma 3. Let X_i have size n and say $X'_i = X_i \cup \{y_i\}$ where $y_i \in \mathcal{X}_i$. Say that the space of datapoints \mathcal{X}_i is bounded above by M_i and bounded below by m_i , and let $y_i \in \mathcal{X}_i$. Let \bar{X}_i , \bar{X}_j , \bar{X}'_i , and \bar{X}'_j be the sample means of X_i , X_j , X'_i and X'_j respectively. Then,

$$n |(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j)| \leq \frac{n}{(n+1)^2} (M_i - m_i)(M_j - m_j).$$

Proof. Note that

$$\begin{aligned}
n |(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j)| &= n \left| \left(\frac{1}{n} \sum_{k=1}^n x_{ki} - \frac{1}{n+1} \sum_{k=1}^{n+1} x'_{ki} \right) \left(\frac{1}{n} \sum_{k=1}^n x_{kj} - \frac{1}{n+1} \sum_{k=1}^{n+1} x'_{kj} \right) \right|, \\
&= n \left| \left(\left(\frac{1}{n} - \frac{1}{n+1} \right) \sum_{k=1}^n x_{ki} - \frac{y_i}{n+1} \right) \left(\left(\frac{1}{n} - \frac{1}{n+1} \right) \sum_{k=1}^n x_{kj} - \frac{y_j}{n+1} \right) \right|, \\
&= n \left| \left(\frac{1}{n(n+1)} \sum_{k=1}^n x_{ki} - \frac{y_i}{n+1} \right) \left(\frac{1}{n(n+1)} \sum_{k=1}^n x_{kj} - \frac{y_j}{n+1} \right) \right|, \\
&= \frac{n}{(n+1)^2} \left| \left(\frac{1}{n} \sum_{k=1}^n x_{ki} - \frac{y_i}{n+1} \right) \left(\frac{1}{n} \sum_{k=1}^n x_{kj} - \frac{y_j}{n+1} \right) \right|, \\
&\leq \frac{n}{(n+1)^2} (M_i - m_i)(M_j - m_j).
\end{aligned}$$

\square

Lemma 4. Let X_i have size n and say $X'_i = X_i \cup \{y_i\}$ where $y_i \in \mathcal{X}_i$. Say that the space of datapoints \mathcal{X}_i is bounded above by M_i and bounded below by m_i , and let $y_i \in \mathcal{X}_i$. Let \bar{X}_i , \bar{X}_j , \bar{X}'_i , and \bar{X}'_j be the sample means of X_i , X_j , X'_i and X'_j respectively. Then,

$$|(y_i - \bar{X}'_i)(y_j - \bar{X}'_j)| \leq \frac{n^2}{(n+1)^2} (M_i - m_i)(M_j - m_j).$$

Proof. Note that

$$\begin{aligned}
|(y_i - \bar{X}'_i)(y_j - \bar{X}'_j)| &= \left| \left(y_i - \frac{y_i + n\bar{X}_i}{n+1} \right) \left(y_j - \frac{y_j + n\bar{X}_j}{n+1} \right) \right|, \\
&= \frac{1}{(n+1)^2} |((n+1)y_i - y_i - n\bar{X}_i) ((n+1)y_j - y_j - n\bar{X}_j)|, \\
&= \frac{n^2}{(n+1)^2} |(y_i - \bar{X}_i)(y_j - \bar{X}_j)|, \\
&\leq \frac{n^2}{(n+1)^2} (M_i - m_i)(M_j - m_j).
\end{aligned}$$

□

2 NEIGHBORING DEFINITION: ADD/DROP ONE

2.1 ℓ_1 -sensitivity

Theorem 1. *Let*

$$f_{ij}(X) = \sum_{k=1}^n (x_{ki} - \bar{X}_i)(x_{kj} - \bar{X}_j).$$

Say that the space of datapoints \mathcal{X}_i is bounded above by M_i and bounded below by m_i . Then the ℓ_1 -sensitivity of $f(\cdot)$ is bounded by

$$\frac{n}{(n+1)} (M_i - m_i)(M_j - m_j).$$

Proof. Adding a point:

$$\begin{aligned}
|f_{ij}(X') - f_{ij}(X)| &= |n(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j) + (y_i - \bar{X}_i)(y_j - \bar{X}_j)| \quad (\text{By Lemma 2}) \\
&\leq n |(\bar{X}_i - \bar{X}'_i)(\bar{X}_j - \bar{X}'_j)| + |(y_i - \bar{X}_i)(y_j - \bar{X}_j)| \\
&\leq \frac{n}{(n+1)^2} (M_i - m_i)(M_j - m_j) + \frac{n^2}{(n+1)^2} (M_i - m_i)(M_j - m_j) \\
&\quad (\text{By Lemmas 3 and 4}) \\
&= \frac{n}{(n+1)} (M_i - m_i)(M_j - m_j). \tag{2.1}
\end{aligned}$$

Removing a point:

Note that Lemma 2 can be rewritten to this setting by parametrizing n as $n-1$ and swapping X and X' in its expression:

$$f_{ij}(X) = f_{ij}(X') + (n-1)(\bar{X}'_i - \bar{X}_i)(\bar{X}'_j - \bar{X}_j) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j).$$

Note that Lemmas 3 and 4 may be rewritten with the same reparametrization:

$$(n-1) |(\bar{X}'_i - \bar{X}_i)(\bar{X}'_j - \bar{X}_j)| \leq \frac{n-1}{n^2} (M_i - m_i)(M_j - m_j),$$

and

$$|(y_i - \bar{X}'_i)(y_j - \bar{X}'_j)| \leq \frac{(n-1)^2}{n^2} (M_i - m_i)(M_j - m_j).$$

Then,

$$\begin{aligned}
|f_{ij}(X) - f_{ij}(X')| &= |(n-1)(\bar{X}'_i - \bar{X}_i)(\bar{X}'_j - \bar{X}_j) + (y_i - \bar{X}'_i)(y_j - \bar{X}'_j)|, \\
&\leq |(n-1)(\bar{X}'_i - \bar{X}_i)(\bar{X}'_j - \bar{X}_j)| + |(y_i - \bar{X}'_i)(y_j - \bar{X}'_j)|, \\
&\leq \frac{n-1}{n^2}(M_i - m_i)(M_j - m_j) + \frac{(n-1)^2}{n^2}(M_i - m_i)(M_j - m_j), \\
&= \frac{n-1}{n}(M_i - m_i)(M_j - m_j).
\end{aligned} \tag{2.2}$$

Note that for any $n \geq 1$,

$$\frac{n}{n+1} > \frac{n-1}{n}. \tag{2.3}$$

So, the worst-case bound always occurs in the “add-one” case, and in general the ℓ_1 sensitivity of $f(\cdot)$ is bounded by

$$\frac{n}{(n+1)}(M_i - m_i)(M_j - m_j).$$

□

Corollary 1. *The ℓ_1 -sensitivity of sample covariance is bounded by*

$$\frac{n}{(n^2 - 1)}(M_i - m_i)(M_j - m_j).$$

2.2 ℓ_2 -sensitivity

Theorem 2. *Sample covariance has ℓ_2 sensitivity bounded above by*

$$\left(\frac{n}{(n^2 - 1)}(M_i - m_i)(M_j - m_j) \right)^2.$$

Proof.

□

3 NEIGHBORING DEFINITION: CHANGE ONE

3.1 ℓ_1 -sensitivity

3.2 ℓ_2 -sensitivity