3Rs: Replace, Refine, Reduce

Matthew J. Salganik Department of Sociology Princeton University

Summer Institute in Computational Social Science June 23, 2018

The Summer Institute in Computational Social Science is supported by grants from the Russell Sage Foundation and the Alfred P. Sloan Foundation.





The Principles of Humane Experimental Technique by Russell and Burch (1959)

- Replace
- Refine
- ▶ Reduce

Experimental evidence of massive-scale emotional contagion through social networks

Adam D. I. Kramer^{a,1}, Jamie E. Guillory^{b,2}, and Jeffrey T. Hancock^{b,c}

^aCore Data Science Team, Facebook, Inc., Menlo Park, CA 94025; and Departments of ^bCommunication and ^cInformation Science, Cornell University, Ithaca, NY 14853

http://dx.doi.org/10.1073/pnas.1320040111




Fig. 1. Mean number of positive (*Upper*) and negative (*Lower*) emotion words (percent) generated people, by condition. Bars represent standard errors.

PNAS

Editorial Expression of Concern and Correction

PSYCHOLOGICAL AND COGNITIVE SCIENCES

PNAS is publishing an Editorial Expression of Concern regarding the following article: "Experimental evidence of massive-scale emotional contagion through social networks," by Adam D. I. Kramer, Jamie E. Guillory, and Jeffrey T. Hancock, which appeared in issue 24, June 17, 2014, of *Proc Natl Acad Sci USA* (111:8788–8790; first published June 2, 2014; 10.1073/pnas.1320040111). This paper represents an important and emerging area of social science research that needs to be approached with sensitivity and with vigilance regarding personal privacy issues.

Replace experiments with less invasive methods

Replace experiments with less invasive methods

Detecting Emotional Contagion in Massive Social Networks

Lorenzo Coviello¹, Yunkyu Sohn², Adam D. I. Kramer³, Cameron Marlow³, Massimo Franceschetti¹, Nicholas A. Christakis^{4,5}, James H. Fowler^{2,6}*

Refine treatments to make them less harmful

Refine treatments to make them less harmful

Rather than blocking posts, they could have boosted posts

Reduce the number of participants

Reduce the number of participants

▶ Difference-in-difference estimator rather than a difference-of-means estimator.

Reduce the number of participants

- ▶ Difference-in-difference estimator rather than a difference-of-means estimator.
- ▶ Would have cut the required sample size, perhaps by half (based on Deng et al. (2013) & Xie and Aurisset (2016)).

When should we care about reducing the number of participants?

When should we care about reducing the number of participants?

- 1. uncertainty about whether the experiment will cause harm
- 2. participation was not voluntary



AND A LEAN, SILENT FIGURE SLOWLY FADES INTO THE GATHERING DARKNESS, AWARE AT LAST THAT IN THIS WORLD, WITH GREAT POWER THERE MUST ALSO COME - GREAT RESPONSIBILITY!



AND SO A LEGEND IS BORN AND A NEW NAME IS ADDED TO THE ROSTER OF THOSE WHO MAKE THE WORLD OF FANTASY THE MOST EXCITING REALM OF ALL!



With great power there must also come great responsibility

The 3 Rs shows that humane methods can be an opportunity:

The 3 Rs shows that humane methods can be an opportunity:

potentially more efficient than standard methods

The 3 Rs shows that humane methods can be an opportunity:

- potentially more efficient than standard methods
- stimulates interesting research (e.g., differential privacy)

Questions?

SICSS

June 17, 2018 - June 30, 2018 | Duke University

Sponsored by the Russell Sage Foundation & The Alfred P. Sloan Foundation