

## 10. Exercise: The mean-squared error

### Exercise: The mean-squared error

1/1 point (graded)

In this exercise we want to understand a little better the formula

$$\frac{1}{\sum_{i=0}^n \frac{1}{\sigma_i^2}}$$

for the mean squared error by considering two alternative scenarios.

In the first scenario,  $\Theta \sim N(0, 1)$  and we observe  $X = \Theta + W$ , where  $W \sim N(0, 1)$  is independent of  $\Theta$ .

In the second scenario, the prior information on  $\Theta$  is extremely inaccurate:  $\Theta \sim N(0, \sigma_0^2)$ , where  $\sigma_0^2$  is so large that it can be treated as infinite. But in this second scenario we obtain two observations of the form  $X_i = \Theta + W_i$ , where the  $W_i$  are standard normals, independent of each other and of  $\Theta$ .

The mean squared error is

☐ smaller in the first scenario.

☐ smaller in the second scenario.

☒ the same in both scenarios. ✓

### Solution:

We use the formula for the mean squared error. For the second scenario, we set  $\sigma_0^2 = \infty$ . In the first scenario, we obtain

$$\frac{1}{\frac{1}{1} + \frac{1}{1}} = \frac{1}{2},$$

and in the second scenario, we obtain the same mean squared error:

$$\frac{1}{\frac{1}{\infty} + \frac{1}{1} + \frac{1}{1}} = \frac{1}{2}.$$

This suggests the following interpretation: the prior information on  $\Theta$  in the first scenario is, in a loose sense, exactly as informative as having no useful prior information but one more observation, as in the second scenario.

提交

You have used 1 of 1 attempt

**i** Answers are displayed within the problem

讨论

显示讨论

**Topic:** Unit 7 / Lec. 15 / 10. Exercise: The mean-squared error