


3. Introduction to the K-Medoids Algorithm

Introduction to the K-Medoids Algorithm



because we explicitly are going to compute the distances between these points and the rest.

So doing this algorithm, we for sure can solve the two problems which were limiting

for us in the case of K-means.

We can work with any distance functions as you can compute it.

And we also are guaranteed to get points from our regional set.



[End of transcript. Skip to the start.](#)

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)



K-Medoids Algorithm as a Variation of K-Means

1/1 point (graded)

As explained in the lecture video, the K-Medoids algorithm is a variation of the K-Means algorithm that addresses some of the K-Means algorithm's limitations. The K-Medoids algorithm is given by

1. Randomly select $\{z_1, \dots, z_K\} \subseteq \{x_1, \dots, x_n\}$

2. Iterate

1. Given z_1, \dots, z_K , assign each $x^{(i)}$ to the closest z_j , so that

$$\text{Cost}(z_1, \dots, z_K) = \sum_{i=1}^n \min_{j=1, \dots, K} \text{dist}(x^{(i)}, z_j)$$

2. Given $C_j \in \{C_1, \dots, C_K\}$ find the best representative $z_j \in \{x_1, \dots, x_n\}$ such that

$$\sum_{x^{(i)} \in C_j} \text{dist}(x^{(i)}, z_j)$$

is minimal.

Which part of the K-Medoids algorithm is **different** from its equivalent counterpart in the K-Means algorithm?

☐ Part 2.1

☒ Part 2.2 

Solution:

As mentioned in the lecture, the k-medoids algorithm is another version of the k-means algorithm with line 2.2 changed so that

1. It is guaranteed that the K representatives $z_1, \dots, z_K \in \{x_1, \dots, x_n\}$
2. Line 2.2 finds cost-minimizing representatives z_1, \dots, z_K with any kind of cost measure

Submit

You have used 1 of 1 attempt


 Answers are displayed within the problem

Fact Check on the K-Medoids Algorithm

1/1 point (graded)

Which of the following is true about the K-Medoids algorithm? Choose all those apply.

☒ It is always guaranteed that the K representatives $z_1, \dots, z_K \in \{x_1, \dots, x_n\}$ 

☒ Line 2.2 of the algorithm(Given $C_j \in \{C_1, \dots, C_K\}$ find the best representative $z_j \in \{x_1, \dots, x_n\}$ such that...) finds the cost-minimizing representatives z_1, \dots, z_K for any distance measure 



Solution:

The K-Medoids algorithm is designed so that the two limitations of the K-Means algorithm are resolved.

Submit

You have used 1 of 2 attempts

 Answers are displayed within the problem

Discussion

Show Discussion

Topic: Unit 4 Unsupervised Learning (2 weeks) :Lecture 14. Clustering 2 / 3. Introduction to the K-Medoids Algorithm