7. LLMS estimation

Problem 6. LLMS estimation

2.5/5.0 points (graded)

Let X and W be independent and uniformly distributed on [-1,1]. We have given the following facts:

$$egin{aligned} \mathbf{E}[X] &= \mathbf{E}[X^3] = \mathbf{E}[X^5] = 0 \ & \mathbf{E}[X^2] = 1/3 \ & \mathbf{E}[X^4] = 1/5 \end{aligned}$$

Suppose that

$$Y = X^3 + W$$

1. Find the LMS estimate of Y, given that X = x. (Notice we are trying to estimate Y from X, not the opposite direction.) (Your answer should be a function of x.)

2. Find the LLMS estimate for $m{Y}$, given that $m{X}=m{x}$. (Your answer should be a function of $m{x}$.)

$$\widehat{Y}_{\mathsf{LLMS}}(x) = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$$
 Answer: 0.6*x

STANDARD NOTATION

Solution:

1.
$$\widehat{Y}_{\mathsf{LMS}}(x) = \mathbf{E}[Y|X=x] = \mathbf{E}[X^3 + W|X=x] = \mathbf{E}[x^3 + W] = x^3$$
 .

2. Since X,Y are both zero mean, we have $\mathrm{Cov}(X,Y)=\mathbf{E}[XY]$, and

$$egin{align} \hat{Y}_{\mathsf{LLMS}}(x) &= \mathbf{E}[Y] + rac{\mathbf{E}[XY]}{\mathbf{E}[X^2]}(x - \mathbf{E}[X]) \ &= 0 + rac{\mathbf{E}[X(X^3 + W)]}{\mathbf{E}[X^2]}x \ &= rac{\mathbf{E}[X^4]}{\mathbf{E}[X^2]}x \ &= rac{3}{5}x \ \end{split}$$

提交

你已经尝试了2次(总共可以尝试2次)

Answers are displayed within the problem