



课程 > Unit 1: ... > Proble... > 2. Set ...

2. Set operations and probabilities

Problem 2. Set operations and probabilities

3/3 points (graded)

Find the value of $\mathbf{P}(A \cup (B^c \cup C^c)^c)$ for each of the following cases:

1. The events A, B, C are disjoint events and $\mathbf{P}(A) = 2/5$.

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = \boxed{2/5} \quad \checkmark \text{ Answer: 0.4}$$

2. The events A and C are disjoint, and $\mathbf{P}(A) = 1/2$ and $\mathbf{P}(B \cap C) = 1/4$.

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = \boxed{3/4} \quad \checkmark \text{ Answer: 0.75}$$

3. $\mathbf{P}(A^c \cap (B^c \cup C^c)) = 0.7$.

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = \boxed{0.3} \quad \checkmark \text{ Answer: 0.3}$$

Solution:

1. Using de Morgan's law, we have $(B^c \cup C^c)^c = B \cap C = \emptyset$ so that

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = \mathbf{P}(A \cup \emptyset) = \mathbf{P}(A) = \boxed{2/5}.$$

2. Note that A and $B \cap C$ are disjoint. Therefore, using de Morgan's law again, together with the additivity axiom for two disjoint events, we have

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = \mathbf{P}(A \cup (B \cap C)) = \mathbf{P}(A) + \mathbf{P}(B \cap C) = \boxed{3/4}.$$



3. De Morgan's law implies that $(A^c \cap (B^c \cup C^c))^c = A \cup (B^c \cup C^c)^c$, which is the event of interest. Therefore,

$$\mathbf{P}(A \cup (B^c \cup C^c)^c) = 1 - \mathbf{P}(A^c \cap (B^c \cup C^c)) = \boxed{0.3}.$$

提交

You have used 1 of 3 attempts

i Answers are displayed within the problem

讨论

显示讨论

Topic: Unit 1 / Problem Set / 2. Set operations and probabilities

Learn About Verified Certificates

© All Rights Reserved