课程 > Unit 1 Introduction to statistics > Lecture 1: What is statistics > 8. Let's do some statistics

8. Let's do some statistics The first example

Start of transcript. Skip to the end.

(Caption will be displayed when you start playing the video.)

This is it for the philosophy part of the class. And we'll start doing some really serious

statistics

in good French tradition.

Let's talk about kissing.

All right.

So everybody's familiar with this sculpture by Rodin.

So this is The Kiss.

视频

下载视频文件

字幕

下载 SubRip (.srt) file 下载 Text (.txt) file

Review: probability question

1/1 point (graded)

Assume that we observe three draws, X_1, X_2, X_3 from a Bernoulli distribution with parameter $p = \frac{1}{2}$. For example, imagine that in the model for the preferred head direction for kissing, either direction were actually equally likely and we observed three kissing couples.

What is the probability of observing at least two ones, i.e., what is $\mathbf{P}(\sum_{i=1}^3 X_i \geq 2)$?

1/2

✓ Answer: 0.5

Solution:

 $\sum_{i=1}^3 X_i$ follows a Binomial distribution with parameters n=3 and $p=rac{1}{2}$, hence the probability in question is

$$\mathbf{P}\left(\sum_{i=1}^3 X_i \geq 2
ight) = {3 \choose 2} {\left(rac{1}{2}
ight)}^3 + {3 \choose 3} {\left(rac{1}{2}
ight)}^3 = rac{4}{8} = rac{1}{2}.$$

提交

你已经尝试了1次(总共可以尝试1次)

Answers are displayed within the problem

Confidence, continued

1/1 point (graded)

direction for kiss	sing. Now, 10 stude	e we decided to consider two or more right-turns as significant evidence for a predisposition of this ents go out and each observe three different couples kissing. How many of them would on average eaning is more common than left-leaning when kissing?
5	✓ A	nswer: 5
Solution:		
We just comput	ed the chance for o	ne of these events to occur to be $rac{1}{2}$, so if we perform 10 repeats, we expect it to happen $f 5$ times.
提交 你已经	圣尝试了1次(总共可以	人尝试1次)
• Answers a	re displayed within	the problem
Friendships		
	people indexed $oldsymbol{1}$ th	arough n , each pair (i,j) (there are $\binom{n}{2}$ of them) are either friends, or not friends. To model this able to each pair. Which one of the probability distributions below is the most appropriate model?
A Poisson	random variable.	
O A Gaussiar	n random variable.	
A Bernoull	i random variable. •	✓
O An expone	ntial random variab	ole.
Bernoulli distrib		takes only two values (for convenience, ${f 1}$ for being friends; and ${f 0}$ for strangers), and this is precisely a 从尝试3次)
• Answers a	re displayed within	the problem
1/2 points (graded We say a group Independent of	of four people is "in	nteresting", if there are at most five pairs who are friends. Assume that each pair of people are friends, ith probability $1/2$. Let N be the number of pairs that are friends in this group. w?
O Poisson	n	
O Bernou	ılli	
Binomi	ial 🗸	
What is the i	aronability that a ra	andomly chosen group of four people is "interesting"?

Solution:

- There are, in total, $\binom{4}{2}=6$ different pairs. Notice that N is a random sum of 6 Bernoulli trials; this is a binomial random variable: $N\sim \mathrm{Bin}\,(6,1/2)$.
- $\mathbb{P}(N \le 5) = 1 \mathbb{P}(N = 6) = 1 (1/2)^6$.

提交

你已经尝试了3次(总共可以尝试3次)

1 Answers are displayed within the problem

How many interesting groups?

0/1 point (graded)

Following the model above, if 128 different people observe randomly chosen groups of four people, how many times on average do these observations lead to the conclusion that the society is interesting?

0.984

X Answer: 126

Solution:

We just computed the chance for one of these events to occur to be 1-1/64, so if we perform 128 repeated experiments, we expect it to happen $128\left(1-\frac{1}{64}\right)=126$ times.

提交

你已经尝试了3次(总共可以尝试3次)

• Answers are displayed within the problem

讨论

显示讨论

主题: Unit 1 Introduction to statistics:Lecture 1: What is statistics / 8. Let's do some statistics

认证证书是什么?

© 保留所有权利