

## 2. Review of Parametric Hypothesis Testing

### Review: Goal of Hypothesis Testing

1/1 point (graded)

You have access to samples  $\mathbf{X}_1, \dots, \mathbf{X}_n \stackrel{iid}{\sim} \mathbf{P}_\theta$  for some unknown parameter  $\theta \in \mathbb{R}$ . You would like to hypothesis test between some null hypothesis  $H_0$  and an alternative hypothesis  $H_1$ .

What is the purpose of hypothesis testing?

- ☐ To solve exactly for the true parameter  $\theta$ .
- ☐ To provide a consistent estimator for the true parameter  $\theta$ .
- ☐ To develop an estimator that is close to the true parameter  $\theta$ .
- ☒ To decide with quantified error whether or not  $\theta$  lies in a certain region of the parameter space. ✓

#### Solution:

We provide the correct response and then discuss the incorrect ones.

"To decide with quantified error whether or not  $\theta$  lies in a certain region of the parameter space." is the goal of hypothesis testing. The null and alternative hypothesis both describe subsets of the parameter space, so to decide whether or not to reject the null (which is the goal of a hypothesis test), we must assess how likely it is, based on given samples, that the parameter would fall within the region defined by  $H_0$ .

- "To solve exactly for the true parameter  $\theta$ ." is incorrect. In general in statistics, since we only have samples from the distribution, it will not be possible to solve for  $\theta$  exactly.
- The second and third choices "To provide a consistent estimator for the true parameter  $\theta$ ." and "To develop an estimator that is close to the true parameter  $\theta$ ." are incorrect. These are some of the goals of parameter estimation.

提交

你已经尝试了1次（总共可以尝试2次）

❗ Answers are displayed within the problem

### Review: Rejection Region

0/1 point (graded)

#### Setup:

You have samples  $\mathbf{X}_1, \dots, \mathbf{X}_n \stackrel{iid}{\sim} \text{Ber}(p^*)$  for some true parameter  $p^* \in (0, 1)$ . Let  $(\{0, 1\}, \{\mathbf{P}_p\}_{p \in (0, 1)})$  denote the associated statistical model, where  $\mathbf{P}_p = \text{Ber}(p)$ .

You conduct a hypothesis test between

- a null hypothesis  $H_0 : p^* \in \Theta_0$  and
- an alternative hypothesis  $H_1 : p^* \in \Theta_1$ ,

where  $\Theta_0, \Theta_1 \subset (0, 1)$  and  $\Theta_0$  and  $\Theta_1$  are disjoint.

You construct a statistical **test**

$$\psi : \{0, 1\}^n \rightarrow \{0, 1\}$$

which takes as input the sample  $(X_1, \dots, X_n)$ . If  $\psi(X_1, \dots, X_n) = 1$ , you will **reject** the null  $H_0$  in favor of the alternative  $H_1$ , and otherwise you will **fail to reject** the null.

Recall that the **rejection region**  $R_\psi$  describes which outcomes  $(x_1, \dots, x_n)$  will result in  $\psi(x_1, \dots, x_n) = 1$  and, hence, rejection of the null.


**Questions:**

The rejection region is a subset of ...  
(Choose all that apply.)

☒  $(\Theta_0)^n$  where  $\Theta_0$  defines the null hypothesis  $H_0$  in the parameter space  $\Theta$

☐  $(\Theta_1)^n$  where  $\Theta_1$  defines the alternative hypothesis  $H_1$  in the parameter space  $\Theta$

☒  $(\Theta)^n$  where  $\Theta$  is the parameter space

☐  $E^n$  where  $E$  is the sample space of  $X_i$  



**Solution:**

The rejection region is by definition the set of all observed outcomes for which  $H_0$  will be rejected by the test  $\psi = \mathbf{1}((X_1, \dots, X_n) \in R_\psi)$ . It is a subset of  $E^n$ , where  $E$  is the sample space of  $X_i$ .

提交

你已经尝试了2次（总共可以尝试2次）

 Answers are displayed within the problem

### Review : What Type of Objects are These?

5/5 points (graded)

**Setup as above:**

You have samples  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p^*)$  for some true parameter  $p^* \in (0, 1)$ . Let  $(\{0, 1\}, \{\mathbf{P}_p\}_{p \in (0, 1)})$  denote the associated statistical model, where  $\mathbf{P}_p = \text{Ber}(p)$ .

You conduct a hypothesis test between

- a null hypothesis  $H_0 : p^* \in \Theta_0$  and
- an alternative hypothesis  $H_1 : p^* \in \Theta_1$ ,

where  $\Theta_0, \Theta_1 \subset (0, 1)$  and  $\Theta_0$  and  $\Theta_1$  are disjoint.

You construct a statistical **test**

$$\psi : \{0, 1\}^n \rightarrow \{0, 1\}$$

which takes as input the sample  $(X_1, \dots, X_n)$ . If  $\psi(X_1, \dots, X_n) = 1$ , you will **reject** the null  $H_0$  in favor of the alternative  $H_1$ , and otherwise you will **fail to reject** the null.

Recall that the **rejection region**  $R_\psi$  describes which samples  $(X_1, \dots, X_n)$  will result in  $\psi(X_1, \dots, X_n) = 1$  and, hence, rejection of the null.

Let  $\alpha_\psi$  and  $\beta_\psi$  denote the **type 1 error** and **type 2 error**, respectively.

**Questions:**  
Determine which type of mathematical object each of the following is.(You are encouraged to review the definitions from the slides of the last lecture.)  
(Choose one for each column.)

Rejection Region:	Type 1 Error:	Level:	Type 2 Error:	Power:
<input type="radio"/> A number.	<input type="radio"/> A number.	<input checked="" type="radio"/> A number. ✓	<input type="radio"/> A number.	<input checked="" type="radio"/> A number. ✓
<input checked="" type="radio"/> A set. ✓	<input type="radio"/> A set.	<input type="radio"/> A set.	<input type="radio"/> A set.	<input type="radio"/> A set.
<input type="radio"/> A function.	<input checked="" type="radio"/> A function. ✓	<input type="radio"/> A function.	<input checked="" type="radio"/> A function. ✓	<input type="radio"/> A function.

**Solution:**

We recall the definitions of each object in the context of the statistical model  $(\{0, 1\}, \{\mathbf{P}_p\}_{p \in (0,1)})$ .

The rejection region is defined to be

$$R_\psi := \{\mathbf{x} \in \{0, 1\}^n : \psi(\mathbf{x}) = 1\},$$

so this is a **set**.

The type 1 error is defined to be

$$\begin{aligned} \alpha_\psi : \Theta_0 &\rightarrow [0, 1] \\ p &\mapsto P_p(\psi = 1), \end{aligned}$$

so this is a **function** of  $p$ .

A level of a test is defined to be a **number**  $\alpha$  such that

$$\alpha \geq \alpha_\psi(p) \quad \text{for all } p \in \Theta_0. \text{ or equivalently } \alpha \geq \sup_{p \in \Theta_0} \alpha_\psi(p)$$

The type 2 error is defined to be

$$\begin{aligned} \beta_\psi : \Theta_1 &\rightarrow [0, 1] \\ p &\mapsto P_p(\psi = 0), \end{aligned}$$

so this is a **function** of  $p$ .

The power  $\pi_\psi$  is defined as

$$\pi_\psi := \inf_{p \in \Theta_1} (1 - \beta_\psi(p)).$$

This greatest lower bound of  $(1 - \beta_\psi(p))$  over a set of  $p$  values is a **number**.

提交

你已经尝试了1次（总共可以尝试2次）

**i** Answers are displayed within the problem

2/2 points (graded)

Setup as above:

let  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p^*)$  for some true parameter  $p^* \in (0, 1)$ , and let  $(\{0, 1\}, \{P_p\}_{p \in (0, 1)})$  denote the associated statistical model where  $P_p = \text{Ber}(p)$ .

Hypotheses for this problem:

You would like to hypothesis test between two simple hypotheses:

$$H_0 : p^* \in \Theta_0 = \{1/2\}$$

$$H_1 : p^* \in \Theta_1 = \{3/4\}.$$

That is, the regions defined by the null and alternative hypotheses each consists of a single point in the parameter space  $\Theta = [0, 1]$ .

You constructed a statistical test  $\psi$ , and let  $\alpha_\psi$  and  $\beta_\psi$  denote the **type 1 error** and **type 2 error**, respectively, associated to this test.

Questions:

What does  $\alpha_\psi(1/2)$  represent?

- ☒ The probability that we **reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  even though **in fact**  $p^* = 1/2$  ✓
- ☐ The probability that we **fail to reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  given **in fact**  $p^* = 1/2$
- ☐ The probability that we **reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  given **in fact**  $p^* = 3/4$
- ☐ The probability that we **fail to reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  even though **in fact**  $p^* = 3/4$

What does  $\beta_\psi(3/4)$  represent?

- ☐ The probability that we **reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  even though **in fact** that  $p^* = 1/2$
- ☐ The probability that we **fail to reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  given that **in fact**  $p^* = 1/2$
- ☐ The probability that we **reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  given that **in fact**  $p^* = 3/4$
- ☒ The probability that we **fail to reject**  $p^* = 1/2$  in favor of  $p^* = 3/4$  even though **in fact**  $p^* = 3/4$  ✓

Solution:

Let's consider the first question. If  $\psi = 1$ , then we would reject the null-hypothesis  $p^* \in \Theta_0 = \{1/2\}$ . Therefore  $\alpha_\psi(1/2) = \mathbf{P}_{1/2}(\psi = 1)$  is the probability of **rejecting**  $p^* \in \Theta_0 = \{1/2\}$  in favor of  $p^* \in \Theta_1 = \{3/4\}$  even when in fact  $p^* \in \Theta_0 = \{1/2\}$ .

Now let's consider the second question. If  $\psi = 0$ , then we would fail to reject the null hypothesis  $p^* \in \Theta_0 = \{1/2\}$  in favor of the alternative hypothesis  $p^* \in \Theta_1 = \{3/4\}$ . Therefore  $\beta_\psi(3/4) = \mathbf{P}_{3/4}(\psi = 0)$  is the probability of not rejecting  $H_0 : p^* \in \Theta_0 = \{1/2\}$  even when  $p^* \in \Theta_1 = \{3/4\}$ .

The other two choices are probabilities when the correct conclusions are made, not errors.

提交

你已经尝试了1次（总共可以尝试2次）

Answers are displayed within the problem

讨论

显示讨论