

### 3. Method of moments estimators

For each of the following distributions, give the method of moments estimator in terms of the sample averages  $\overline{X}_n$  and  $\overline{X}_n^2$ , assuming we have access to  $n$  i.i.d. observations  $X_1, \dots, X_n$ . In other words, express the parameters as functions of  $\mathbb{E}[X_1]$  and  $\mathbb{E}[X_1^2]$  and then apply these functions to  $\overline{X}_n$  and  $\overline{X}_n^2$ .

(a)

1/1 point (graded)

$$X_i \sim \text{Ber}(p), \quad p \in (0, 1)$$

(If applicable, write **barX\_n** for  $\overline{X}_n$ .)

Method of moments estimator  $\hat{p} =$   ☐

你已经尝试了1次（总共可以尝试3次）

☐ 正确 (1/1 分)

(b)

1/1 point (graded)

$$X_i \sim \text{Poiss}(\lambda), \quad \lambda > 0,$$

which means that each  $X_1$  has density

$$\mathbf{P}_\lambda(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k \in \mathbb{N}.$$

Method of moments estimator  $\hat{\lambda} =$   ☐

你已经尝试了1次（总共可以尝试3次）

☐ 正确 (1/1 分)

(c)

1/1 point (graded)

$$X_i \sim \text{Exp}(\lambda), \quad \lambda > 0,$$

which means that each  $X_1$  has density

$$f_{\lambda}(x) = \lambda e^{-\lambda x}, \quad x > 0.$$

Method of moments estimator  $\hat{\lambda} =$ 

1/ barX\_n

提交

你已经尝试了1次（总共可以尝试3次）

正确 (1/1 分)

(d)

2/2 points (graded)

$$X_i \sim \mathcal{N}(\mu, \sigma^2), \quad \mu \in \mathbb{R}, \sigma^2 > 0,$$

which means that each  $X_1$  has density

$$f_{\mu, \sigma^2}(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

(If applicable, enter **barX\_n** for  $\overline{X_n}$  and **bar(X\_n^2)** for  $\overline{X_n^2}$ .)

Method of moments estimator  $\hat{\mu} =$ 

barX\_n

Method of moments estimator  $\widehat{\sigma^2} =$ 

bar(X\_n^2) - barX\_n^2

STANDARD NOTATION

提交

你已经尝试了1次（总共可以尝试3次）

正确 (2/2 分)

(e)

2 points possible (graded)

$X_i$  follows a shifted exponential distribution with parameters  $a \in \mathbb{R}$  and  $\lambda > 0$ . That means each  $X_i$  has density

$$f_{a,\lambda}(x) = \lambda e^{-\lambda(x-a)} \mathbf{1}\{x \geq a\}, \quad x \in \mathbb{R}.$$

(If applicable, enter **barX\_n** for  $\overline{X_n}$  and **bar(X\_n^2)** for  $\overline{X_n^2}$ .)

Method of moments estimator  $\hat{a} =$

Method of moments estimator  $\hat{\lambda} =$

STANDARD NOTATION

提交

你已经尝试了0次（总共可以尝试3次）

Add a Post

显示所有帖子 ▼	近期活动 ▼
<div><div></div><div><a href="#">Algorithm of Method of Moments [edited]</a></div></div>	7 ▼
<div><div></div><div><a href="#">[STAFF] Part (e) 2nd question</a></div><div><a href="#">Hi, Can you please check if my answer to the 2nd question of part (e) is actually wrong. I was getting a math processing error but was able to submit and got 1s...</a></div></div>	1 ▼
<div><div></div><div><a href="#">[Staff] Estimator of a in (e).</a></div></div>	13 ▼
<div><div></div><div><a href="#">[Staff] e) Strange behavior of grader rendering.</a></div><div><a href="#">After inputting `sqrt` in one of the fields, the first rendering window stop working and the second one shows `math rendering error`. Page refresh doesn't help....</a></div></div>	5 ▼
<div><div></div><div><a href="#">e) Just to make sure</a></div><div><a href="#">1) Does <math>\bar{X}_n</math> denote the sample mean of the shifted exponential distribution? 2) Does <math>\bar{(X_n^2)}</math> denote the sample mean of each realization of the shifted ...</a></div></div>	2 ▼
<div><div></div><div><a href="#">a), b), c).</a></div><div><a href="#">There is a problem with <b><math>\bar{X}_n</math></b> rendering.</a></div></div>	2 ▼
<div><div></div><div><a href="#">[staff] <math>\bar{(X_n^2)}</math> is not accepted.</a></div><div><a href="#"><math>\bar{(X_n^2)}</math> is not accepted . Message is "Invalid Input: <math>X_n</math> not permitted in answer as a variable".</a></div></div>	3 ▼