

5. Parameter Tuning

Extension Note: Project 5 due date has been extended by 1 **more** day to **September 6 23:59UTC** .

Effects of adjusting epsilon

0 points possible (ungraded)

Ungrading Note: The problem is now ungraded because there has been a lot of confusion.

In this question, you will investigate the impact of ϵ on the convergence of Q-learning algorithm. Which of the below do you observe from running the algorithm?

☒ For very large ϵ (say $\epsilon = 1$), the algorithm converges slower compared to $\epsilon = 0.5$ ✓

☐ For very large ϵ (say $\epsilon = 1$), the algorithm converges faster compared to $\epsilon = 0.5$

☒ For very small ϵ (say $\epsilon = 0.00001$), the algorithm converges slower compared to $\epsilon = 0.5$

☐ For very small ϵ (say $\epsilon = 0.00001$), the algorithm converges faster compared to $\epsilon = 0.5$ ✓

✗

Solution:

A large value of ϵ means exploring more (randomly), not using much of what we have learned. A small ϵ , on the other hand, will generate experience consistent with the current estimates of Q-values, but will explore less. For this toy task, however, the state space is small enough that random initialization is enough to induce diversity in the experience collected.

Submit

You have used 0 of 3 attempts

i Answers are displayed within the problem

Effects of alpha

0 points possible (ungraded)

In this question, you will investigate the impact of α on the convergence of Q-learning algorithm. Fix the exploration parameter $\epsilon = 0.5$ and do the experiments with different values of the training $\alpha \in [10^{-6}, 1]$. What you have observed?

☐ The algorithm converges for all values of α in less than 200 epochs

☒ The algorithm does not converge for all values of α in less than 200 epochs

☐ The smaller α , the slower the convergence

☐ The smaller α , the faster the convergence

✓

Solution:

For large values of α , learning is too instable. For small values of α , learning is too slow.

 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 5 Reinforcement Learning (2 weeks) :Project 5: Text-Based Game / 5. Parameter Tuning

Add a Post

◀ All Posts

[staff, please look again] Effects of adjusting epsilon: what I see on the plots is not accepted by the grader

question posted 27 days ago by anonymous

Tried to mark also the opposite choice to what I see, and this answer also got rejected.

This post is visible to everyone.



Soumya_Ram (Staff)

14 days ago - marked as answer 14 days ago by **Soumya_Ram** (Staff)



Sorry, this is an error on our behalf. We updated the problem one too many times. (now the solutions don't match the edited problem description). The problem is ungraded now.

Show Comment (1) ▼

Add a Response

15 other responses

Cool7 (Community TA)

27 days ago



I think normally a very small ϵ will also lead to slower convergence. Agent will most likely stick to first action will yield positive reward even it's not the optimal action.

But in our case, all most all Q values are negative, first positive Q value discovered is most likely within optimal policy.

Maybe this is just a bad example for this question, but we really need to put in something we didn't observe?



I also believe this example does not work well for the question asked. To get my answer accepted, I ended up selecting one option that was contradicting the plot.

posted 27 days ago by **OlgaSkv**



Anonymous author, please mark this topic with [Staff] tag. Dear staff, please take a look, as results we obtain by looking at the plots are not accepted by grader.

posted 26 days ago by **disguiser**



Yes... Same here...

posted 26 days ago by **LiWeic1c9**



Same here

posted 24 days ago by **Tantrum83**



Same