

RESOLUÇÃO • LÍSTA • REGRA DA CADEIA

01) Em cada caso, verifique se a função real u , satisfaz à equação à derivadas parciais indicada:

a) $u = f(x, y)$; $x(\lambda, s) = \lambda^3 - s^3$; $y(\lambda, s) = s^3 - \lambda^3$; $s^2 \frac{\partial u}{\partial \lambda} + \lambda^2 \frac{\partial u}{\partial s} = 0$;

Solução: usamos o quadro 01:

$$\frac{\partial u}{\partial \lambda} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \lambda} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \lambda} \therefore \frac{\partial u}{\partial \lambda} = \frac{\partial u}{\partial x} (3\lambda^2) + \frac{\partial u}{\partial y} (-3\lambda^2) \therefore s^2 \frac{\partial u}{\partial \lambda} = 3s^2 \lambda^2 \frac{\partial u}{\partial x} - 3s^2 \lambda^2 \frac{\partial u}{\partial y};$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial s} \therefore \frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} (-3s^2) + \frac{\partial u}{\partial y} (3s^2) \therefore \lambda^2 \frac{\partial u}{\partial s} = -3\lambda^2 s^2 \frac{\partial u}{\partial x} + 3\lambda^2 s^2 \frac{\partial u}{\partial y};$$

$$\text{Logo, } s^2 \frac{\partial u}{\partial \lambda} + \lambda^2 \frac{\partial u}{\partial s} = 3s^2 \lambda^2 \frac{\partial u}{\partial x} - 3s^2 \lambda^2 \frac{\partial u}{\partial y} - 3\lambda^2 s^2 \frac{\partial u}{\partial x} + 3\lambda^2 s^2 \frac{\partial u}{\partial y} = 0;$$

Sim, satisfaz;

b) $u = f(x, y)$; $x(\lambda, s) = \lambda + s$; $y(\lambda, s) = \lambda - s$; $\left(\frac{\partial u}{\partial x}\right)^2 - \left(\frac{\partial u}{\partial y}\right)^2 = \frac{\partial u}{\partial \lambda} \frac{\partial u}{\partial s}$;

Solução: novamente usaremos o quadro 01:

$$\frac{\partial u}{\partial \lambda} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \lambda} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \lambda} \therefore \frac{\partial u}{\partial \lambda} = \frac{\partial u}{\partial x} \cdot (1) + \frac{\partial u}{\partial y} \cdot (1) = \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y};$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial s} \therefore \frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \cdot (1) + \frac{\partial u}{\partial y} \cdot (-1) = \frac{\partial u}{\partial x} - \frac{\partial u}{\partial y};$$

$$\text{Logo, } \frac{\partial u}{\partial \lambda} \frac{\partial u}{\partial s} = \left(\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y}\right) \left(\frac{\partial u}{\partial x} - \frac{\partial u}{\partial y}\right) = \left(\frac{\partial u}{\partial x}\right)^2 - \left(\frac{\partial u}{\partial y}\right)^2; \text{ Sim, satisfaz;}$$

c) $u = f(x, y, z)$; $x(\lambda, s, t) = \lambda - s$; $y(\lambda, s, t) = s - t$; $z(\lambda, s, t) = t - \lambda$; $\frac{\partial u}{\partial \lambda} + \frac{\partial u}{\partial s} + \frac{\partial u}{\partial t} = 0$;

Solução: usaremos o quadro 05:

$$\frac{\partial u}{\partial \lambda} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \lambda} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \lambda} + \frac{\partial u}{\partial z} \frac{\partial z}{\partial \lambda} = \frac{\partial u}{\partial x} \cdot (1) + \frac{\partial u}{\partial y} \cdot (0) + \frac{\partial u}{\partial z} \cdot (-1) = \frac{\partial u}{\partial x} - \frac{\partial u}{\partial z};$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial u}{\partial z} \frac{\partial z}{\partial s} = \frac{\partial u}{\partial x} \cdot (-1) + \frac{\partial u}{\partial y} \cdot (1) + \frac{\partial u}{\partial z} \cdot (0) = -\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y};$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial t} + \frac{\partial u}{\partial z} \frac{\partial z}{\partial t} = \frac{\partial u}{\partial x} \cdot (0) + \frac{\partial u}{\partial y} \cdot (-1) + \frac{\partial u}{\partial z} \cdot (1) = -\frac{\partial u}{\partial y} + \frac{\partial u}{\partial z};$$

$$\text{Logo, } \frac{\partial u}{\partial \lambda} + \frac{\partial u}{\partial s} + \frac{\partial u}{\partial t} = \frac{\partial u}{\partial x} - \frac{\partial u}{\partial z} - \frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} - \frac{\partial u}{\partial y} + \frac{\partial u}{\partial z} = 0;$$

Sim, satisfaz;

(d) $u = F(x, y)$; $x(n, s) = \frac{s-n}{ns}$; $y(s, t) = \frac{t-s}{st}$; $n, s, t \neq 0$; $n^2 \frac{\partial u}{\partial n} + s^2 \frac{\partial u}{\partial s} + t^2 \frac{\partial u}{\partial t} = 0$;

Solução: em primeiro lugar, como n, s e t são não-nulos, escrevamos:

$$x(n, s) = \frac{s}{ns} - \frac{n}{ns} = \frac{1}{n} - \frac{1}{s}; \quad y(s, t) = \frac{t-s}{st} = \frac{t}{st} - \frac{s}{st} = \frac{1}{s} - \frac{1}{t};$$

Em segundo lugar, observemos que nenhum dos quadros da teoria se encaixa, exatamente, em nossa situação.

Vamos então criar nosso quadro:

$$\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial n} = \frac{\partial u}{\partial x} \left(-\frac{1}{n^2}\right);$$

$$\frac{\partial u}{\partial s} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial s} = \frac{\partial u}{\partial x} \left(\frac{1}{s^2}\right) + \frac{\partial u}{\partial y} \left(-\frac{1}{s^2}\right);$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial y} \frac{\partial y}{\partial t} = \frac{\partial u}{\partial y} \cdot \left(-\frac{1}{t^2}\right);$$

$$\begin{aligned} \text{Logo, } n^2 \frac{\partial u}{\partial n} + s^2 \frac{\partial u}{\partial s} + t^2 \frac{\partial u}{\partial t} &= n^2 \left(\frac{\partial u}{\partial x} \cdot \left(-\frac{1}{n^2}\right) \right) + s^2 \left(\frac{\partial u}{\partial x} \cdot \left(\frac{1}{s^2}\right) - \frac{\partial u}{\partial y} \left(\frac{1}{s^2}\right) \right) + t^2 \left(\frac{\partial u}{\partial y} \cdot \left(-\frac{1}{t^2}\right) \right); \\ &= -\frac{\partial u}{\partial x} + \frac{\partial u}{\partial x} - \frac{\partial u}{\partial y} + \frac{\partial u}{\partial y} = 0; \text{ Sim, satisfaz;} \end{aligned}$$

(e) $u = F(an+bs)$; $a, b \in \mathbb{R}$; $b \frac{\partial u}{\partial n} - a \frac{\partial u}{\partial s} = 0$;

Solução: começamos introduzindo a variável $x(n, s) = an+bs$;

Assim, ficamos com $u = F(x)$; $x(n, s) = an+bs$; Usaremos o quadro (03);

$$\frac{\partial u}{\partial n} = \frac{du}{dx} \frac{\partial x}{\partial n} = \frac{du}{dx} \cdot a; \quad \text{Logo, } b \frac{\partial u}{\partial n} - a \frac{\partial u}{\partial s} = b a \frac{du}{dx} - a b \frac{du}{dx} = 0;$$

$$\frac{\partial u}{\partial s} = \frac{du}{dx} \frac{\partial x}{\partial s} = \frac{du}{dx} \cdot b; \quad \text{Sim, satisfaz;}$$

(f) $u(n, s) = s + F(n^2 - s^2)$; $s \frac{\partial u}{\partial n} + n \frac{\partial u}{\partial s} = n$;

Solução: vamos introduzir, $v = F(x)$ com $x(n, s) = n^2 - s^2$;

$$\text{Então, } \frac{\partial u}{\partial n} = \frac{\partial(s)}{\partial n} + \frac{\partial v}{\partial n} = \frac{\partial v}{\partial n}; \quad \text{e } \frac{\partial u}{\partial s} = \frac{\partial(s)}{\partial s} + \frac{\partial v}{\partial s} = 1 + \frac{\partial v}{\partial s};$$

Falta descobirmos $\frac{\partial v}{\partial n}$ e $\frac{\partial v}{\partial s}$; Usaremos, novamente, o quadro (03) para $v = F(x)$;

$$\begin{aligned} \frac{\partial v}{\partial n} &= \frac{dv}{dx} \cdot \frac{\partial x}{\partial n} = 2n \frac{dv}{dx}; \\ \frac{\partial v}{\partial s} &= \frac{dv}{dx} \cdot \frac{\partial x}{\partial s} = -2s \frac{dv}{dx}; \end{aligned} \quad \begin{aligned} s \frac{\partial u}{\partial n} + n \frac{\partial u}{\partial s} &= s \frac{\partial v}{\partial n} + n \left(1 + \frac{\partial v}{\partial s} \right) \\ &= s(2n \frac{dv}{dx}) + n + n(-2s \frac{dv}{dx}) \\ &= 2ns \frac{dv}{dx} + n - 2ns \frac{dv}{dx} = n; \end{aligned}$$

Sim, satisfaz;

⑧ $u(r,s) = rs + f(r^2 + s^2)$; $s \frac{\partial u}{\partial r} - r \frac{\partial u}{\partial s} = s^2 - r^2$;

Solução: vamos introduzir, $v = f(x)$ com $x(r,s) = r^2 + s^2$;

Então, $\frac{\partial u}{\partial r} = \frac{\partial(rs)}{\partial r} + \frac{\partial v}{\partial r} = s + \frac{\partial v}{\partial r}$; e $\frac{\partial u}{\partial s} = \frac{\partial(rs)}{\partial s} + \frac{\partial v}{\partial s} = r + \frac{\partial v}{\partial s}$;

E, para encontrarmos $\frac{\partial v}{\partial r}$ e $\frac{\partial v}{\partial s}$ vamos usar o quadro ③:

$$\begin{aligned} \frac{\partial v}{\partial r} &= \frac{dv}{dx} \frac{\partial x}{\partial r} = 2r \frac{dv}{dx}; \\ \frac{\partial v}{\partial s} &= \frac{dv}{dx} \frac{\partial x}{\partial s} = 2s \frac{dv}{dx}; \end{aligned} \quad \left. \begin{array}{l} \\ \end{array} \right\} \begin{aligned} s \frac{\partial u}{\partial r} - r \frac{\partial u}{\partial s} &= s(s + \frac{\partial v}{\partial r}) - r(r + \frac{\partial v}{\partial s}) \\ &= s^2 + 2rs \frac{dv}{dx} - r^2 - 2rs \frac{dv}{dx} = s^2 - r^2; \end{aligned}$$

Sim, satisfaz;

⑨ Sejam $u = f(x,y)$; $x(r,\theta) = r \cos \theta$; $y(r,\theta) = r \sin \theta$; Mostre que:

$$\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 = \left(\frac{\partial u}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial u}{\partial \theta}\right)^2;$$

Solução: usaremos o quadro ①:

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta;$$

$$\frac{\partial u}{\partial \theta} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial u}{\partial x} (-r \sin \theta) + \frac{\partial u}{\partial y} (r \cos \theta) = r \left[\frac{\partial u}{\partial y} \cos \theta - \frac{\partial u}{\partial x} \sin \theta \right];$$

Então:

$$\begin{aligned} \left(\frac{\partial u}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial u}{\partial \theta}\right)^2 &= \left(\frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta\right)^2 + \frac{1}{r^2} \left[r^2 \left(\frac{\partial u}{\partial y} \cos \theta - \frac{\partial u}{\partial x} \sin \theta\right)^2\right] \\ &= \left(\frac{\partial u}{\partial x}\right)^2 \cos^2 \theta + 2 \frac{\partial u}{\partial x} \frac{\partial u}{\partial y} \cos \theta \sin \theta + \left(\frac{\partial u}{\partial y}\right)^2 \sin^2 \theta + \left(\frac{\partial u}{\partial y}\right)^2 \cos^2 \theta - 2 \frac{\partial u}{\partial x} \frac{\partial u}{\partial y} \cos \theta \sin \theta + \left(\frac{\partial u}{\partial x}\right)^2 \sin^2 \theta \\ &= \left(\frac{\partial u}{\partial x}\right)^2 (\cos^2 \theta + \sin^2 \theta) + \left(\frac{\partial u}{\partial y}\right)^2 (\sin^2 \theta + \cos^2 \theta) \\ &= \left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2; \end{aligned}$$

03) [Equações de Cauchy-Riemann em coordenadas polares;]

Sejam $u(x,y)$ e $v(x,y)$ Funções reais satisfazendo às equações de Cauchy (1789-1857) - Riemann (1826-1866). Mostre que se $x(r,\theta) = r \cos \theta$ e $y(r,\theta) = r \sin \theta$, então: $\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$ e $\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$;

Solução:

Pelo que acabamos de ver na questão 02 a Regra da Cadeia nos diz, que:

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta; \quad \text{e Cauchy-Riemann nos diz, que: } \begin{cases} \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \end{cases}$$

$$\frac{\partial v}{\partial \theta} = r \left(\frac{\partial v}{\partial x} \cos \theta - \frac{\partial v}{\partial y} \sin \theta \right);$$

$$\text{Logo, } \frac{1}{r} \frac{\partial v}{\partial \theta} = \frac{\partial v}{\partial x} \cos \theta - \frac{\partial v}{\partial y} \sin \theta = \frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta = \frac{\partial u}{\partial r};$$

Beim como,

$$\frac{\partial v}{\partial r} = \frac{\partial v}{\partial x} \cos \theta + \frac{\partial v}{\partial y} \sin \theta;$$

$$\frac{\partial u}{\partial \theta} = r \left(\frac{\partial u}{\partial x} \cos \theta - \frac{\partial u}{\partial y} \sin \theta \right);$$

$$\text{Logo, } -\frac{1}{r} \frac{\partial u}{\partial \theta} = -\frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta = \frac{\partial v}{\partial x} \cos \theta + \frac{\partial v}{\partial y} \sin \theta = \frac{\partial v}{\partial r};$$

04) [Fórmula de Leibniz]

Sejam $u, v: F(u,v) = \int_v^u p(t) dt$; $u = g(x)$; $v = h(x)$; Mostre a seguinte Fórmula devida à Leibniz (1646-1716): $\frac{dw}{dx} = p(g(x))g'(x) - p(h(x))h'(x)$;

Solução: Usaremos o quadro 07 e o Teorema Fundamental do Cálculo:

$$\frac{dw}{dx} = \frac{\partial w}{\partial u} \frac{du}{dx} + \frac{\partial w}{\partial v} \frac{dv}{dx} = \frac{\partial \left(\int_v^u p(t) dt \right)}{\partial u} \cdot g'(x) + \frac{\partial \left(\int_v^u p(t) dt \right)}{\partial v} \cdot h'(x)$$

$$= p(u) \cdot g'(x) + \frac{\partial \left(-\int_u^v p(t) dt \right)}{\partial v} \cdot h'(x)$$

$$= p(g(x))g'(x) - p(v)h'(x)$$

$$= p(g(x))g'(x) - p(h(x))h'(x);$$

05 [Teorema de Euler para Funções homogêneas;]

Diz-se que $F(x,y)$ é homogênea de grau n , $n \geq 0$ um inteiro, quando $F(tx,ty) = t^n F(x,y)$. Mostre a seguinte igualdade, devida à Euler (1707-1783);

$$x \frac{\partial F}{\partial x} + y \frac{\partial F}{\partial y} = n F(x,y);$$

Solução: Vamos denotar $w = F(tx,ty) = t^n F(x,y)$;

• Então, de $w = t^n F(x,y)$ vemos que $\frac{\partial w}{\partial t} = n t^{n-1} F(x,y)$; (I)

• Agora para encontrarmos $\frac{\partial w}{\partial t}$ na expressão $w = F(tx,ty)$, introduzamos as variáveis $u(t,x) = tx$ e $v(t,y) = ty$. Então:

$$\frac{\partial w}{\partial t} = \frac{\partial w}{\partial u} \frac{\partial u}{\partial t} + \frac{\partial w}{\partial v} \frac{\partial v}{\partial t} = \frac{\partial w}{\partial u} x + \frac{\partial w}{\partial v} y; \text{ (II)}$$

Logo, concluímos de (I) e (II) que para todos t, x e y reais, temos:

$$x \frac{\partial w}{\partial u} + y \frac{\partial w}{\partial v} = n t^{n-1} F(x,y);$$

O que Euler, genialmente, percebeu é que, em particular, esta igualdade vale para $t=1$ e para todos x e y reais, e que nesse caso $u=x$ e $v=y$. Logo ela se torna:

$$x \frac{\partial w}{\partial x} + y \frac{\partial w}{\partial y} = n F(x,y);$$

06 Em cada caso, mostre que se a equação:

Ⓐ $F(x,y) = 0$, definir y como uma função diferenciável de x e $F_y \neq 0$, então $\frac{dy}{dx} = -\frac{F_x}{F_y}$;

Solução: Como y é função derivável de x , o quadro 07 nos informa:

$$\text{De } F(x,y) = 0, \text{ temos: } \frac{dF(x,y)}{dx} = 0; \therefore \frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{dy}{dx} = 0; \therefore F_x + F_y \frac{dy}{dx} = 0;$$

$$\text{E como } F_y \neq 0, \frac{dy}{dx} = -\frac{F_x}{F_y};$$

Ⓑ $F(x,y,z) = 0$, definir z como uma função diferenciável de x e y e $F_z \neq 0$, então: $\frac{\partial z}{\partial x} = -\frac{F_x}{F_z}$; e $\frac{\partial z}{\partial y} = -\frac{F_y}{F_z}$;

Solução: como z é função diferenciável de x e y , o quadro 02 nos informa:

$$\text{• De } F(x,y,z) = 0: \frac{\partial (F(x,y,z))}{\partial x} = 0: \therefore \frac{\partial F}{\partial x} \frac{\partial x}{\partial x} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} = 0; \therefore F_x + F_z \frac{\partial z}{\partial x} = 0;$$

$$\text{E como, } F_z \neq 0, \text{ então: } \frac{\partial z}{\partial x} = -\frac{F_x}{F_z};$$

$$\text{• De } F(x,y,z) = 0: \frac{\partial (F(x,y,z))}{\partial y} = 0: \therefore \frac{\partial F}{\partial x} \frac{\partial x}{\partial y} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial y} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial y} = 0; \therefore F_y + F_z \frac{\partial z}{\partial y} = 0;$$

$$\text{E como, } F_z \neq 0, \text{ temos: } \frac{\partial z}{\partial y} = -\frac{F_y}{F_z};$$