

# Introduction to Induction

## Recognizing a pattern

Consider the following pattern

$$\begin{aligned}1 + 3 &= 4 \\1 + 3 + 5 &= 9 \\1 + 3 + 5 + 7 &= 16\end{aligned}$$

The pattern at this point is hopefully becoming clear. Let us do one more

$$1 + 3 + 5 + 7 + 9 = 25$$

Looks like the sum of odd numbers is equal to some perfect square. It is time to formalize this. Every odd number is of the form  $2k + 1$  for some integer  $k$ . That should lead us to this

$$\sum_{i=1}^n 2i - 1 = n^2$$

where  $n \in \mathbb{N}$ .

Whenever you have a result that follows a pattern like this, in order to prove it you have to show the formula/pattern holds for every single value of  $n$ . Since  $n$  takes values in ‘discrete’ steps (1,2,3 etc), the powerful method called induction can be applied to prove it.

## Idea behind induction

Induction is used to prove that a theorem holds for all natural numbers (sometimes we will include 0).

If we are able to show the following.

- the theorem holds for 1 (or some small number)
- If the theorem holds for  $n - 1$ , then it will hold for  $n$ .

Now let’s say we want to show the theorem is going to hold for 5. Here’s how we could do it

- it holds for 1

- Since it holds for 1, it holds for 2.
- Since it holds for 2, it holds for 3.
- Since it holds for 3, it holds for 4.
- Since it holds for 4, it holds for 5.

This idea of using the previous to prove the current is the crux of induction. If you have seen recursion in programming, this methodology of solving sub problems and using their solution to solve the big problem should look familiar. It is closely related to recursion.

## Proof algorithm for induction

1. Write the statement of the theorem in terms of a predicate that has an input of a single number.
2. Prove the theorem for the smallest possible  $n$ . Generally this will be something like 0 or 1 but it depends on the theorem. This is called the base case.
3. Assume the theorem is true for  $n - 1$ . That is assume  $P(n-1)$  is true.
4. Show that  $P(n - 1) \implies P(n)$ .

## Proof of summations

Let us attempt an inductive proof of the first result.

$$\sum_{i=1}^n 2i - 1 = n^2$$

Define the predicate  $P(n)$  as a function which returns true or false depending

Base case: When  $n = 1$ , there is only term in the summation which is  $2 - 1 = 1$  and the right side is also 1. So it is true for 1.

Assume the result is true for  $n - 1$  So

$$\sum_{i=1}^{n-1} 2i - 1 = (n - 1)^2$$

To show if  $P(n - 1)$  then  $P(n)$ .

Consider the sum

$$\begin{aligned}
\sum_{i=1}^n 2i - 1 &= \sum_{i=1}^{n-1} 2i - 1 + 2n - 1 \\
&= (n-1)^2 + 2n - 1 && \text{(using P(n-1) true)} \\
&= n^2 - 2n + 1 + 2n - 1 \\
&= n^2
\end{aligned}$$

which shows that the predicate is true for  $n$ .

## Proof of inequalities

Which is greater  $n!$  or  $2^n$ ?

$3!$  is 6 and  $2^3$  is 8.  $4!$  is 24 and  $2^4$  is 16.  $5!$  is 120 and  $2^5 = 32$ . So looks like the inequality  $n! > 2^n$  holds for  $n \geq 4$

To prove this claim by induction.

Let the predicate be defined as  $n! > 2^n, n \geq 4$ .

The base case is  $n = 4$  which we have already shown to be true.

Let the statement hold true for  $n - 1$ . Meaning

$$(n-1)! > 2^{n-1} \tag{1}$$

To prove this inductively we need to use this to show  $n! > 2^n$ .

Multiply both sides of (1) by  $n$  since  $n! = (n-1)!n$ .

So we get  $n! > 2^{n-1} \cdot n$ . Since  $n > 4$ , we know that

$$2^n \cdot n > 2^{n-1} \cdot 2$$

$$\text{But } 2^{n-1} \cdot 2 = 2^n$$

Hence proved

## Example involving sets

Generalized De-Morgan's law.

We will do this in class and the steps are all in the Zybook.

## Induction puzzle

In Josephine's Kingdom every woman has to pass a logic exam before being allowed to marry. Every married woman knows about the fidelity of every man in the Kingdom except for her own husband, and etiquette demands that no woman should tell another about the fidelity of her husband. Also, a gunshot fired in any house in the Kingdom will be heard in any other house. Queen Josephine announced that unfaithful men had been discovered in the

Kingdom, and that any woman knowing her husband to be unfaithful was required to shoot him at midnight following the day after she discovered his infidelity. How did the wives manage this?

This is a classic puzzle and the solution can actually be found on wikipedia. We will discuss in class or recitation.

## Divisibility proof

$2^{n+2} + 3^{2n+1}$  is divisible by 7 for all positive integers

Proof by induction

Let  $P(n)$  be the predicate that returns true when  $2^{n+2} + 3^{2n+1}$  is divisible by 7.

Base case:  $P(1)$  says  $2^3 + 3^3 = 8 + 27 = 35$  and 35 is  $5 \times 7$ .

To show  $P(n-1) \rightarrow P(n)$ .

We are given that  $2^{(n-1)+2} + 3^{2(n-1)+1}$  is divisible by 7. That is  $2^{n+1} + 3^{2n-1}$  is divisible by 7.

We now consider  $2^{n+2} + 3^{2n+1}$  and simplify it to the point where we can use our induction hypothesis and prove the theorem.

$$\begin{aligned} 2^{n+2} + 3^{2n+1} &= 2 \cdot (2^{n+1} + 3^{2n-1}) + 3^{2n+1} - 2 \cdot 3^{2n-1} \\ &= 2 \cdot (2^{n+1} + 3^{2n-1}) + 3^{2n-1}(3^2 - 2) \\ &= 2 \cdot (2^{n+1} + 3^{2n-1}) + 3^{2n-1} \cdot 7 \end{aligned}$$

The first term is divisible by 7 because of the induction hypothesis. The second term is clearly divisible by 7.

Hence we have shown that  $P(n-1) \rightarrow P(n)$ , which along with the base case completes a proof by induction.