

Where the DEGs sit on the genome - scheme (B) per strain

All 11 contigs, stacked one row per strain (pME008 sits at the right of the Odo_St23 row).
Upper track: DESeq2 log2 fold change, KO in the numerator and WT in the denominator, apeglm-shrunk (lfcShrink, coef = group_KO_vs_WT). No pseudocount is added: it is NOT log2((KO+1)/(WT+1)), it is the coefficient of the negative-binomial GLM fitted to size-factor-normalized counts.
Lower track: log10(x+1) of the group mean size-factor-normalized count, WT drawn upward and KO downward. Here the +1 IS added, only so that zeros fit on a log scale.
Red = DEG (padj < 0.05). 9,665 genes drawn (every gene with an LFC; 27 of them have padj = NA, almost all Cook-distance outliers, and are drawn grey).
Bands = loci where 3+ consecutive DEGs all move the same way (red = up, blue = down): 298 loci / 1344 genes. Purple = eptA genotype markers (triangle = off the y axis).

Bth_VPI




Bth_St8



Bun_8492



Odo_St23



Scheme (B) fits every strain on its own, so the pME008 plasmid is not in any model and has no LFC: its panel is drawn but left empty on purpose, and that emptiness is itself the result. x in Mb. px/Mb is identical across strain rows and between the two schemes (each row padded on the right to 6.88 Mb). Contigs shorter than 250 kb are drawn at that minimum width, so their internal scale is expanded (7 of 11 panels). The y axis (+/-7 and +/-7) is shared by every panel and by both schemes, so the two figures can be compared directly.