

Math Notes

Chang Si Yuan

May 29, 2016

(ver 0.0.1a)

This compilation of notes are to be used as a reference for the GCE "A"-level Mathematics paper, both as a refresher in theories as well as for general descriptions of presentation form. These notes are meant for free, public use, but at the reader's own risk.
Good luck with your exams.

1 Assumed Knowledge

1.1 Algebra

1.1.1 Completing the Square

$$x^2 + bx + c = \left(x + \frac{b}{2}\right)^2 + c - \left(\frac{b}{2}\right)^2$$

1.1.2 Polynomial Expansions

$$\begin{aligned}(a \pm b)^2 &= a^2 \pm 2ab + b^2 \\ a^2 - b^2 &= (a + b)(a - b) \\ a^3 \pm b^3 &= (a \pm b)(a^2 \mp 2ab + b^2)\end{aligned}$$

1.1.3 Partial Fractions

$$\begin{aligned}&\frac{f(x)}{(ax + b)(cx + d)} \\&= g(x) + \frac{A}{ax + b} + \frac{B}{cx + d} \\&\frac{f(x)}{(ax + b)(cx + d)^2} \\&= g(x) + \frac{A}{ax + b} + \frac{B}{cx + d} + \frac{C}{(cx + d)^2} \\&\frac{f(x)}{(ax + b)(x^2 + c)} \\&= g(x) + \frac{A}{ax + b} + \frac{Bx + C}{x^2 + c}\end{aligned}$$

1.1.4 Exponent and Logarithm

$$e^n = \underbrace{e \times e \times e \times \dots \times e}_{n \text{ times}}$$

$$e^{\frac{1}{2}} = \sqrt{e}$$


$$\log_e(x) = \ln(x)$$

= how many times e is multiplied by itself to get x

$$\log_{10}(x) = \lg(x)$$

$$x = e^{\ln(x)}$$

$$\log_x(y) = \frac{\log_{\text{base}}(y)}{\log_{\text{base}}(x)}$$



1.2 Trigonometry

1.2.1 Sine and Cosine Rule

For any triangle with length of sides a , b and c and with opposite angles A , B and C :

$$\begin{aligned}\frac{a}{\sin(A)} &= \frac{b}{\sin(B)} = \frac{c}{\sin(C)} \\ a^2 &= b^2 + c^2 - 2bccos(A) \\ cos(A) &= \frac{b^2 + c^2 - a^2}{2bc}\end{aligned}$$

1.2.2 Sum of Angles

$$\begin{aligned}\sin(A \pm B) &= \sin(A)\cos(B) \pm \cos(A)\sin(B) \\ \sin(2A) &= 2\sin(A)\cos(A) \\ \cos(A \pm B) &= \cos(A)\cos(B) \mp \sin(A)\sin(B) \\ \cos(2A) &= \cos^2(A) - \sin^2(A) \\ &= 2\cos^2(A) - 1 \\ &= 1 - 2\sin^2(A) \\ \tan(A \pm B) &= \frac{\tan(A) \pm \tan(B)}{1 \mp \tan(A)\tan(B)} \\ \tan(2A) &= \frac{2\tan(A)}{1 - \tan^2(A)} \\ \text{Area of Triangle} &= \frac{1}{2}absin(C)\end{aligned}$$

1.2.3 Factor and Reverse Factor Formula

$$\begin{aligned}\sin(A) + \sin(B) &= 2\sin\left(\frac{A+B}{2}\right)\cos\left(\frac{A-B}{2}\right) \\ \sin(A) - \sin(B) &= 2\cos\left(\frac{A+B}{2}\right)\sin\left(\frac{A-B}{2}\right) \\ \cos(A) + \cos(B) &= 2\cos\left(\frac{A+B}{2}\right)\cos\left(\frac{A-B}{2}\right) \\ \cos(A) - \cos(B) &= -2\sin\left(\frac{A+B}{2}\right)\sin\left(\frac{A-B}{2}\right) \\ \sin(A)\cos(B) &= \frac{1}{2}[\sin(A+B) + \sin(A-B)] \\ \cos(A)\sin(B) &= \frac{1}{2}[\sin(A+B) - \sin(A-B)] \\ \cos(A)\cos(B) &= \frac{1}{2}[\cos(A+B) + \cos(A-B)] \\ \sin(A)\sin(B) &= -\frac{1}{2}[\cos(A+B) - \cos(A-B)]\end{aligned}$$

2 Inequalities

2.1 Inequalities

2.1.1 Properties of Inequalities

$$a > b, c > 0 \implies ac > bc$$

$$a > b, c < 0 \implies ac < bc$$

$$\frac{a}{b} > 0 \implies ab > 0$$

$$\frac{a}{b} < 0 \implies ab < 0$$

Positive sides of inequalities suggest that both terms share similar positive or negative signs, negative sides of inequalities suggest that both terms have opposite positive or negative signs

2.1.2 Quadratic Inequalities

Find where $f(x) = 0$ by completing square or quadratic formula and sketch graph

2.1.3 Inequality Reduction

For any inequality $\frac{f(x)}{g(x)} > \text{or} < 0$ where $f(x)$ or $g(x)$ is strictly positive or negative, reduce inequality to non-strictly positive/negative function and change sign accordingly

2.1.4 Modulus Inequalities

$$|x| < a \iff -a < x < a$$

$$|x| > a \iff x < -a \text{ or } a < x$$

$$|x - a| < b \iff a - b < x < a + b$$

$$|x - a| > b \iff x < a - b \text{ or } a + b < x$$

To solve inequalities, sketch and find intercept, then deduce suitable range of x

3 Vectors

3.1 Representation

3.1.1 Point Representation

O is always defined as the origin

Written: \underline{r} or \overrightarrow{OR}

Column: $\mathbf{r} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$

Unit Vector: $\mathbf{r} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ where \mathbf{i}, \mathbf{j} and \mathbf{k} are unit vectors in the x, y and z dimensions

3.1.2 Line Representation

Vector Equation:

$$l : \mathbf{r} = \mathbf{a} + \lambda \mathbf{b} \quad \lambda \in \mathbb{R}$$

Parametric Equation:

$$l : \begin{cases} x = a_x + \lambda b_x \\ y = a_y + \lambda b_y \\ z = a_z + \lambda b_z \end{cases} \quad \lambda \in \mathbb{R}$$

Cartesian Equation:

$$\frac{x - a_x}{b_x} = \frac{y - a_y}{b_y} = \frac{z - a_z}{b_z}$$

3.1.3 Plane Representation

Vector Equation:

$$\pi : \mathbf{r} = \mathbf{a} + \lambda \mathbf{b} + \mu \mathbf{c}$$

Scalar Product Equation:

$$\pi : \mathbf{r} \cdot \mathbf{n} = d$$

Cartesian Equation:

$$\pi : x n_x + y n_y + z n_z = d$$

3.2 Manipulation

3.2.1 Vector Algebra

Vector Addition: remove same inside or outside terms

$$\overrightarrow{OA} + \overrightarrow{AB} = \overrightarrow{OB}$$

Negative Vectors: reverse the points

$$\overrightarrow{AB} = -\overrightarrow{BA}$$

Vector Subtraction: reverse points, then add

$$\overrightarrow{OB} - \overrightarrow{OA} = \overrightarrow{OB} + \overrightarrow{AO} = \overrightarrow{AB}$$

3.2.2 Vector Properties

Modulus / Magnitude : The total distance of a vector

$$\left| \begin{pmatrix} x \\ y \\ z \end{pmatrix} \right| = \sqrt{x^2 + y^2 + z^2}$$

Unit Vector / Cap : A vector which defines a direction and has modulus of 1

$$\hat{\mathbf{a}} = \frac{1}{|\mathbf{a}|} \mathbf{a}$$

3.2.3 Ratio Theorem

3.2.4 Scalar Product

Scalar/Dot product produces a scalar which is a representation of how inline two vectors are with each other

$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} \cdot \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = x_1 x_2 + y_1 y_2 + z_1 z_2$$

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos(\theta)$$

$$\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$$

$$\mathbf{a} \cdot (\perp \mathbf{a}) = 0$$

$$\mathbf{a} \cdot (\mathbf{b} \pm \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$$

$$(\lambda \mathbf{a}) \cdot \mathbf{b} = \lambda (\mathbf{a} \cdot \mathbf{b})$$

3.2.5 Vector Product

Vector/Cross product produces a vector which has direction perpendicular to its input vectors and has magnitude similar to area subtended by its input vectors

$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} \times \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} y_1 z_2 - z_1 y_2 \\ z_1 x_2 - x_1 z_2 \\ x_1 y_2 - y_1 x_2 \end{pmatrix}$$

$$\mathbf{a} \times \mathbf{b} = \hat{\mathbf{n}} |\mathbf{a}| |\mathbf{b}| \sin(\theta)$$

$$\mathbf{a} \times \mathbf{a} = 0$$

$$\mathbf{a} \times (\perp \mathbf{a}) = \hat{\mathbf{n}} |\mathbf{a}| |\mathbf{b}|$$

$$\mathbf{a} \times (\mathbf{b} \pm \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}$$

$$(\lambda \mathbf{a}) \times \mathbf{b} = \lambda (\mathbf{a} \times \mathbf{b})$$

3.3 Angles Between Vectors

General formula:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$$

Lesser used formula:

$$\sin(\theta) = \frac{|\mathbf{a} \times \mathbf{b}|}{|\mathbf{a}||\mathbf{b}|}$$

3.3.1 Point-Point Angle

Let \mathbf{a} and \mathbf{b} be point vectors

3.3.2 Line-Line Angle

Let \mathbf{a} and \mathbf{b} be direction vectors of lines

3.3.3 Line-Plane Angle

Let \mathbf{a} be direction vector of line and \mathbf{b} be normal vector of plane

Angle between line and plane will be $90^\circ - \theta$

3.3.4 Plane-Plane Angle

Let \mathbf{a} and \mathbf{b} be normal vectors of planes

3.4 Intersection

3.4.1 Point-Line Intersection

Solve series of parametric equations or find λ which lets point equal to point on line

3.4.2 Line-Line Intersection

If direction vectors are scalar multiples of each other, lines are parallel

Find a point which satisfies both lines, solving parametric equations of both line equations

$$\mathbf{a}_1 + \lambda \mathbf{b}_1 = \mathbf{a}_2 + \mu \mathbf{b}_2 \quad \lambda, \mu \in \mathbb{R}$$

If lines are both non-parallel and non-intersecting, lines are skew

3.4.3 Line-Plane Intersection

If dot product of direction vector of line and normal of plane equals to 0, line is parallel to plane

Substitute line equation into plane equation and expand to solve for λ

$$\begin{aligned} p &= (\mathbf{a} + \lambda \mathbf{b}) \cdot \mathbf{n} \\ &= \mathbf{a} \cdot \mathbf{n} + \lambda(\mathbf{b} \cdot \mathbf{n}) \end{aligned}$$

3.4.4 Plane-Plane Intersection

If normal of planes are scalar multiples of each other, planes are parallel

Equating two planes results in a line

Cross product of normal vectors of two planes produces the direction vector of line

Position vector of line can be observed from equations, find a vector which satisfies both plane equations

3.5 Projections

3.5.1 Point-Point Projection

To find distance d of projection of point vector \mathbf{a} on point vector \mathbf{b} :

$$d = \mathbf{a} \cdot \hat{\mathbf{b}}$$

3.5.2 Point-Line Projection

To find distance d of projection of point vector \mathbf{a} on direction vector of line \mathbf{b} , similar to Point-Point Projection

3.5.3 Point-Line Perpendicular

To find perpendicular of point vector \mathbf{a} on line $l : \mathbf{r} = \mathbf{b} + \lambda \mathbf{c}$, let \mathbf{P} be a point on the line such that \overrightarrow{AP} is perpendicular to \mathbf{c} and solve for λ :

$$\begin{aligned} 0 &= \overrightarrow{AP} \cdot \mathbf{c} \\ &= (\mathbf{b} + \lambda \mathbf{c} - \mathbf{a}) \cdot \mathbf{c} \quad \lambda |\mathbf{c}|^2 = (\mathbf{a} - \mathbf{b}) \cdot \mathbf{c} \\ &= \lambda \mathbf{c} \cdot \mathbf{c} + (\mathbf{b} - \mathbf{a}) \cdot \mathbf{c} \end{aligned}$$

To find distance d between point vector \mathbf{a} and its perpendicular on line with direction vector \mathbf{b} , find magnitude of the cross product of \mathbf{a} and unit vector of \mathbf{b} :

$$d = |\mathbf{a} \times \hat{\mathbf{b}}|$$

3.5.4 Point-Plane Perpendicular

To find perpendicular of point vector \mathbf{a} on plane $\pi : \mathbf{r} \cdot \mathbf{n} = d$, consider a line containing \mathbf{a} and with direction vector \mathbf{n} , equate the two equations and then solve for λ

$$l = \mathbf{a} + \lambda \mathbf{n}$$

$$\begin{aligned} d &= (\mathbf{a} + \lambda \mathbf{n}) \cdot \mathbf{n} \quad \lambda |\mathbf{n}|^2 = d - \mathbf{a} \cdot \mathbf{n} \\ &= \mathbf{a} \cdot \mathbf{n} + \lambda \mathbf{n} \cdot \mathbf{n} \end{aligned}$$

To find distance d between point vector \mathbf{a} and plane with normal vector \mathbf{n} , find projection of \mathbf{a} on unit vector of \mathbf{n} :

$$d = |\mathbf{a} \cdot \hat{\mathbf{n}}|$$

4 Complex Numbers

4.1 Imaginary Numbers

$$i = \sqrt{-1}$$

$$i^2 = -1$$

$$i^4 = i^2 \times i^2 = 1$$

4.2 Cartesian Representation

4.2.1 Complex Numbers

Complex numbers are written in the form:

$$z = a + ib \quad a, b \in \mathbb{R}$$

$$\text{Where } \operatorname{Re}(z) = a \quad \operatorname{Im}(z) = b$$

And populate the set \mathbb{C}

4.2.2 Conjugates

$$\text{For } w = a + ib \quad z = c + id$$

$$w^* = a - ib$$

$$ww^* = a^2 + b^2 = |w|^2$$

$$(w + z)^* = w^* + z^*$$

$$(wz)^* = w^*z^*$$

4.2.3 Algebraic Manipulation

$$\text{For } w = a + ib \quad z = c + id$$

$$w = z \implies a = c, b = d \quad \text{IMPT}$$

$$w + z = (a + c) + i(b + d)$$

$$w - z = (a - c) + i(b - d)$$

$$w * z = (ac - bd) + i(ad + bc)$$

$$|w| = \sqrt{a^2 + b^2}$$

$$\sqrt{a^2 + b^2} \quad \text{occurs with a } \pm \text{ sign}$$

For division, remove i from denominator by multiplying numerator and denominator by conjugate and then solve:

$$\begin{aligned} \frac{w}{z} &= \frac{w}{z} \times \frac{z^*}{z^*} \\ &= \frac{wz^*}{zz^*} \\ &= \frac{wz^*}{c^2 + d^2} \end{aligned}$$

4.3 Complex Polynomial Roots

4.3.1 Theorem of Algebra

A polynomial of degree n has n real or complex roots

4.3.2 Finding Complex Roots

If a polynomial has all real coefficients, complex roots occur in conjugate pairs. Write as such in exams:

$$f(x) \quad \text{has real coefficients}$$

$$a + ib \quad \text{is a root} \implies a - ib \quad \text{is a root}$$

For a polynomial with complex coefficients, use quadratic general formula. Note that \pm will still be present somewhere

4.4 Polar Representation

4.4.1 Polar Representation

$$z = re^{i\theta} \quad r \in \mathbb{R}_0^+ \quad -\pi < \theta \leq \pi$$

$$|z| = r \quad \arg(z) = \theta = \tan^{-1}\left(\frac{a}{b}\right)$$

$$\operatorname{Re}(z) = r \cos(\theta)$$

$$\operatorname{Im}(z) = r \sin(\theta)$$

$$re^{i\pi} = -1 \quad re^{i0} = 1$$

$$re^{i\frac{\pi}{2}} = i \quad re^{i\frac{-\pi}{2}} = -i$$

4.4.2 Algebraic Manipulation

$$\text{For } z = r_1 e^{i\theta_1} \quad z = r_2 e^{i\theta_2}$$

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}$$

$$z_1^n = r_1^n e^{in\theta_1}$$


$$\sqrt{z_1} = \sqrt{r_1} e^{i(\frac{\theta_1}{2})}, \sqrt{r_1} e^{i(\frac{\theta_1}{2} + \pi)}$$

4.5 Geometric Representation

4.5.1 Argand Diagrams

Cartesian form describes the x and y coordinate on the real and imaginary axis

Polar form describes the distance between the point and the origin as well as the angle a line from the point to the origin makes with the positive real axis



4.5.2 Geometric Manipulation

Multiplying by $re^{i\theta}$ scales the number by a factor of r and rotates anticlockwise by an angle of θ about the origin
 Multiplying by i rotates a complex number by $\frac{\pi}{2}$, or 90° anticlockwise

The conjugate of a complex number is a reflection of the complex number on the x axis

5 Functions

5.1 Properties of a Function

5.1.1 Individual Properties

A function f is a relation which maps input of a set D_f to outputs of a set R_f using a certain rule
 Multiple elements in the input set can have the same output, but one single element in the input set can only have one output.

5.1.2 Function Presentation

When questions ask for functions in a similar form, be sure to maintain presentation

$$f(x) = x^2 \quad x \in (-\infty, \infty)$$

$$f : x \mapsto x^2 \quad x \in (-\infty, \infty)$$

$$g(x) = \begin{cases} x^2 & x \in \mathbb{R}, x > 0 \\ -x & x \in \mathbb{R}, x < 0 \end{cases}$$

$$D_f = (-\infty, \infty) \quad R_f = [0, \infty)$$

Note: infinity is always written as non-inclusive

5.2 Inverse Functions

Inverse functions map the output of a function to the input of a function. Inverse functions only exist when each element of the output set of the original function is mapped to one and only one element in the input set, i.e. inverse functions only exist if a function is one-one
 Inverse functions are written as f^{-1} and

$$D_{f^{-1}} = R_f \quad R_{f^{-1}} = D_f$$

5.2.1 Proving existence and inexistence

$f(x)$ cuts each line $y = k$, $k \in R_f$ at one and only one point, f is one-one, hence f^{-1} exists
 Replace R_f with the actual set
 The line $y = k$ cuts $f(x)$ at more than one point, f is not one-one, hence f^{-1} does not exist
 Replace k with the actual edge case

5.2.2 Finding Inverse Functions

$$\begin{aligned} f(x) &= x^2 + 1 \quad x \in [0, \infty) \text{ Let } y = f(x) = x^2 + 1 \\ x^2 &= y - 1 \\ x &= \sqrt{y - 1} \\ f^{-1}(y) &= \sqrt{y - 1} \\ f^{-1}(x) &= \sqrt{y - 1} \\ D_{f^{-1}} &= R_f = [1, \infty) \\ R_{f^{-1}} &= D_f = [0, \infty) \end{aligned}$$

5.2.3 Graphical Relationships Between Functions and Inverse Functions

Inverse functions are, in essence, a function reflected along the line $y = x$
 Intersections of functions with their inverse also satisfy the condition $f(x) = x$

5.3 Composite Functions

Considering two functions f and g , the composite function fg is obtained when inputs of g are mapped to their outputs of g , which are then used as inputs to f and mapped to outputs of f :

$$fg(x) = f(g(x))$$

5.3.1 Deriving Composite Functions

For the composite function fg to exist, $R_g \subseteq D_f$
 The domain of function fg follows the domain of function g , i.e. $D_{fg} = D_g$
 The rule of fg is obtained by substituting the rule of g into the rule of f
 The range of fg is a subset of R_f and may be limited due to the fact that R_g may be smaller than D_f , hence R_{fg} must be reevaluated after creating its rule

6 APGP

6.1 Arithmetic Progression

6.1.1 Arithmetic Sequence

An arithmetic progression is a sequence of numbers which have the same difference between consecutive elements. Sequences are defined by their initial term a and their constant difference d

$$\begin{aligned} u_1 &= a + (1 - 1)d = a \\ u_2 &= a + (2 - 1)d = a + d \\ u_3 &= a + (3 - 1)d = a + 2d \\ &\dots \\ u_n &= a + (n - 1)d \end{aligned}$$

6.1.2 Arithmetic Series

Series is defined as the sum of a certain number of consecutive elements in an arithmetic sequence.

$$\begin{aligned}S_1 &= u_1 \\S_2 &= S_1 + u_2 = u_1 + u_2 \\S_3 &= S_2 + u_3 = u_1 + u_2 + u_3 \\&\dots \\S_n &= u_1 + u_2 + \dots + u_{n-1} + u_n\end{aligned}$$

For a known arithmetic sequence, the term S_n can be derived from the equation

$$S_n = \frac{n}{2}(2a + (n-1)d) = \frac{n}{2}(u_1 + u_n)$$

6.1.3 Proving a AP

6.1.4 Exponent and Log

6.2 Geometric Progression

6.2.1 Geometric Sequence

6.2.2 Geometric Series

6.2.3 Proving a GP

6.2.4 Convergence