


DIS Galactic Archaeology minor module


Lecture 4: Stars as probes of Galactic Archaeology II







Dr Anke Ardern-Arentsen

Recap from previous lecture



Recap from previous lecture










1 H	big bang fusion 										cosmic ray fission 										2 He								
3 Li	4 Be	merging neutron stars? 										exploding massive stars 										5 B	6 C	7 N	8 O	9 F	10 Ne		
11 Na	12 Mg	dying low mass stars 										exploding white dwarfs 										13 Al	14 Si	15 P	16 S	17 Cl	18 Ar		
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr												
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe												
55 Cs	56 Ba			72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn											
87 Fr	88 Ra																												
		57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu													
		89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	Very radioactive isotopes; nothing left from stars																					

Graphic created by Jennifer Johnson
<http://www.astronomy.ohio-state.edu/~jaj/nucleo/>

Astronomical Image Credits:
ESA/NASA/AASNova


Recap from previous lecture




1 H	big bang fusion 										cosmic ray fission 										2 He																						
3 Li	4 Be	merging neutron stars? 										exploding massive stars 										5 B	6 C	7 N	8 O	9 F	10 Ne																
11 Na	12 Mg	dying low mass stars 										exploding white dwarfs 										13 Al	14 Si	15 P	16 S	17 Cl	18 Ar																
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr																										
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe																										
55 Cs	56 Ba							72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn																					
87 Fr	88 Ra																																										
																						57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu							
																						89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	Very radioactive isotopes; nothing left from stars															

Graphic created by Jennifer Johnson
<http://www.astronomy.ohio-state.edu/~jaj/nucleo/>



Astronomical Image Credits:
ESA/NASA/AASNova



Recap from previous lecture



- Uploaded a notebook with instructions how to make the apparent, absolute and extinction-corrected colour-magnitude diagrams (CMDs)
- Also one there to reproduce a couple of the APOGEE figures I will show today





This lecture

- How to get metallicity and chemical abundances for stars?
 - Spectroscopy (background & techniques)
 - Some recent examples from the literature
- How to use chemical abundances to trace chemical evolution in galaxies?



Measuring abundances

Spectral lines



- The composition of stars is encoded in **stellar spectra, thanks to absorption line features**
- Spectrum = energy (light) across different frequencies (wavelengths)
- Absorption = atoms and molecules absorb energy at specific frequencies (wavelengths)




Continuum Spectrum



Emission Line Spectrum





Absorption Line Spectrum





Spectral lines



- The composition of stars is encoded in **stellar spectra, thanks to absorption line features**
 - Spectrum = energy (light) across different frequencies (wavelengths)
 - Absorption = atoms and molecules absorb energy at specific frequencies (wavelengths)
- Absorption lines are the result of light generated in the hot inside of a star **moving out through the cooler atmosphere**




Continuum Spectrum




Emission Line Spectrum



Absorption Line Spectrum




Solar spectrum



Solar spectrum



From this spectrum, can you tell which element is most abundant in the Sun?




Stellar spectra: metal-poor & metal-rich

*only 1 Fe atom per
 10^9 atoms of H!


Ultra metal-poor star*: only H lines and a tiny Ca line

Sun (metal-rich): absorption features everywhere (not noise!)




Stellar spectra

- Spectra are the result of a combination of **black body radiation** + **absorption lines**




Stellar spectra

- Spectra are the result of a combination of **black body radiation + absorption lines**




S/N of 120, 50 and 13 (top to bottom)




Stellar spectra

- Spectra are the result of a combination of **black body radiation + absorption lines**
- In practice also:
 - the the instrument profile
 - noise (S/N = signal to noise)
 - spurious features (e.g. cosmic rays or sky)




S/N of 120, 50 and 13 (top to bottom)



Stellar spectra

Not all spectra are equal, important differences are:

- **wavelength coverage**
(affecting what elemental lines can be measured)




Stellar spectra

Not all spectra are equal, important differences are:

- **wavelength coverage**
(affecting what elemental lines can be measured)
- **spectral resolution**
(affecting the *quality* of abundance measurements)

$$R = \frac{\lambda}{\Delta\lambda}$$

(and more of each is more expensive)





Stellar spectra

Not all spectra are equal, important differences are:

- **wavelength coverage**
(affecting what elemental lines can be measured)
- **spectral resolution**
(affecting the *quality* of abundance measurements)

$$R = \frac{\lambda}{\Delta\lambda}$$

(and more of each is more expensive)





Stellar spectra

Not all spectra are equal, important differences are:

- **wavelength coverage**
(affecting what elemental lines can be measured)
- **spectral resolution**
(affecting the *quality* of abundance measurements)

$$R = \frac{\lambda}{\Delta\lambda}$$

(and more of each is more expensive)





Stellar spectra

Not all spectra are equal, important differences are:

- **wavelength coverage**
(affecting what elemental lines can be measured)
- **spectral resolution**
(affecting the *quality* of abundance measurements)

$$R = \frac{\lambda}{\Delta\lambda}$$

(and more of each is more expensive)



Christlieb 2003

How many elements can we derive from spectra?


$$R = \frac{\lambda}{\Delta\lambda}$$

- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**

How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$


- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**



How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$


- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**



How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$


- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**
- Medium-resolution: $R \sim 5000$ - $10\,000$ (e.g. Gaia RVS, 4MOST&WEAVE LR), **~10 elements**



How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$

- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**
- Medium-resolution: $R \sim 5000$ - $10\,000$ (e.g. Gaia RVS, 4MOST&WEAVE LR), **~10 elements**
- Low-resolution: $R \sim 1000$ - 2000 (e.g. LAMOST, SEGUE), **~3-5 elements** (more with ML?)




$R \sim 1300$ (AAT)

How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$


- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**
- Medium-resolution: $R \sim 5000$ - $10\,000$ (e.g. Gaia RVS, 4MOST&WEAVE LR), **~10 elements**
- Low-resolution: $R \sim 1000$ - 2000 (e.g. LAMOST, SEGUE), **~3-5 elements** (more with ML?)
- Very low-resolution: Gaia XP ($R \sim 30$ - 100), **3-4 (?) elements**



How many elements can we derive from spectra?

$$R = \frac{\lambda}{\Delta\lambda}$$

- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**
- Medium-resolution: $R \sim 5000$ - $10\,000$ (e.g. Gaia RVS, 4MOST&WEAVE LR), **~10 elements**
- Low-resolution: $R \sim 1000$ - 2000 (e.g. LAMOST, SEGUE), **~3-5 elements** (more with ML?)
- Very low-resolution: Gaia XP ($R \sim 30$ - 100), **3-4 (?) elements**
- Photometry: combine multiple (narrow) bands, **2-3 (?) elements** (Fe, "alpha", C)



How many elements can we derive from spectra?



$$R = \frac{\lambda}{\Delta\lambda}$$

- Sun: the brightest object = very high S/N, observed at $R > 100\,000$, **67 elements**
- High-resolution: bright stars only ($G < 14$, roughly), $R > 25\,000$ - $30\,000$ (e.g. APOGEE, GALAH surveys, and dedicated individual observations), **~20-30 elements**
- Medium-resolution: $R \sim 5000$ - $10\,000$ (e.g. Gaia RVS, 4MOST&WEAVE LR), **~10 elements**
- Low-resolution: $R \sim 1000$ - 2000 (e.g. LAMOST, SEGUE), **~3-5 elements** (more with ML?)
- Very low-resolution: Gaia XP ($R \sim 30$ - 100), **3-4 (?) elements**
- Photometry: combine multiple (narrow) bands, **2-3 (?) elements** (Fe, "alpha", C)

depends very much on S/N & the effective temperature and metallicity of stars



Analysis of absorption features

- Prior to analysis, spectra are usually **"continuum-fit"** and processed into "1D" spectra, into **normalised flux vs. wavelength**
- Absorption features are well-fit with **Gaussian profiles** (*most of the time*)



Analysis of absorption features




- Prior to analysis, spectra are usually **"continuum-fit"** and processed into "1D" spectra, into **normalised flux vs. wavelength**
- Absorption features are well-fit with **Gaussian profiles** (*most of the time*)
- The **depth of lines relate to the total abundance** of that element
- Often used: **Equivalent Width (EW)**
- Also **full-spectrum fitting**



Shape of absorption lines



Why are these spectral lines not delta functions?





Shape of absorption lines



Why are these spectral lines not delta functions?

- Instrumental resolution





Shape of absorption lines



Why are these spectral lines not delta functions?

- Instrumental resolution
- “Natural broadening”: quantum physics





Shape of absorption lines



Why are these spectral lines not delta functions?

- Instrumental resolution
- “Natural broadening”: quantum physics
- “Pressure broadening”: perturbed energy levels





Shape of absorption lines



Why are these spectral lines not delta functions?

- Instrumental resolution
- “Natural broadening”: quantum physics
- “Pressure broadening”: perturbed energy levels
- “Doppler broadening”: motions of atoms



Abundances and equivalent widths: background

$$\log\left(\frac{W}{\lambda}\right) = \log\left(\frac{\pi e^2}{m_e c^2} \frac{N_k/N_E}{U_k(T)} N_H\right) + \log A + \log(g_{kl} f \lambda) - \chi_{kl} \theta_{exc} + const$$

Abundances and equivalent widths: background

$$\log\left(\frac{W}{\lambda}\right) = \log\left(\frac{\pi e^2}{m_e c^2} \frac{N_k/N_E}{U_k(T)} N_H\right) + \log A + \log(g_{kl} f \lambda) - \chi_{kl} \theta_{exc} + const$$

Astrophysics

Abundances and equivalent widths: background

Equivalent
Widths

Abundance

Effective
Temperature

$$\log\left(\frac{W}{\lambda}\right) = \log\left(\frac{\pi e^2}{m_e c^2} \frac{N_k/N_E}{U_k(T)} N_H\right) + \log A + \log(g_{kl} f \lambda) - \chi_{kl} \theta_{exc} + const$$

Stellar
Atmosphere

Astrophysics

Abundances and equivalent widths: background

Equivalent Widths

Abundance

Effective Temperature

Stellar Atmosphere

$$\log\left(\frac{W}{\lambda}\right) = \log\left(\frac{\pi e^2}{m_e c^2} \frac{N_k/N_E}{U_k(T)} N_H\right) + \log A + \log(g_{kl} f \lambda) - \chi_{kl} \theta_{exc} + const$$

Oscillator strength (transition probability)

Excitation potential (energy required for transition)

Astrophysics

Quantum Chemistry (Line list)

Abundances and equivalent widths: in practice


- To relate EWs (line depth) to abundances in practice, requires:
 - A line list of known ionisation states, transition probabilities and excitation energies
 - A model atmosphere for the star parametrised by **effective temperature** T_{eff} and **surface gravity**, $\log g$ and **metallicity** [Fe/H]

Abundances and equivalent widths: in practice

- To relate EWs (line depth) to abundances in practice, requires:
 - A line list of known ionisation states, transition probabilities and excitation energies
 - A model atmosphere for the star parametrised by **effective temperature** T_{eff} and **surface gravity**, $\log g$ and **metallicity** [Fe/H]

- **Effective temperature:** temperature of a black body with the same radiated power/unit area as the star

$$L = 4\pi R^2 \sigma T_{\text{eff}}^4$$



Abundances and equivalent widths: in practice


- To relate EWs (line depth) to abundances in practice, requires:
 - A line list of known ionisation states, transition probabilities and excitation energies
 - A model atmosphere for the star parametrised by **effective temperature** T_{eff} and **surface gravity**, $\log g$ and **metallicity** [Fe/H]

- **Effective temperature:** temperature of a black body with the same radiated power/unit area as the star


$$L = 4\pi R^2 \sigma T_{\text{eff}}^4$$

- **Log g:** log of the surface gravity of the star, measure of the photospheric pressure of the star


$$g = \frac{GM}{R^2}$$




Spectra changing with temperature




Which star do you think is the hottest?




Spectra changing with temperature




Which star do you think is the hottest?



Spectra changing with surface gravity



Spectra changing with metallicity




Spectra changing with stellar velocity


redshift/blueshift due to movement of a star through space with respect to us

Spectra changing with stellar velocity


redshift/blueshift due to movement of a star through space with respect to us




Kiel diagram (HR diagram analogue)



Kiel diagram (HR diagram analogue)




Historical note: stellar classification based on spectra




Cooler stars

Hotter stars




- Sequence was originally (late 19th century) alphabetical, based on the strength of the clearly visible hydrogen lines in stellar spectra, a system designed by **Annie Jump Cannon**, one of the Harvard "computers" (a team of women processing astronomy data)

Historical note: stellar classification based on spectra




Cooler stars

Hotter stars



- Sequence was originally (late 19th century) alphabetical, based on the strength of the clearly visible hydrogen lines in stellar spectra, a system designed by **Annie Jump Cannon**, one of the Harvard "computers" (a team of women processing astronomy data)
- **Cecilia Payne** (early 20th century) was first to accurately relate these classes to temperatures using ionisation theory (she also was the first to claim stars are predominantly hydrogen)

Historical note: stellar classification based on spectra



Cooler stars

Hotter stars

System still used, e.g. astronomers might refer to FGK stars (which can be giants or dwarfs)

- Sequence was originally (late 19th century) alphabetical, based on the strength of the clearly visible hydrogen lines in stellar spectra, a system designed by **Annie Jump Cannon**, one of the Harvard “computers” (a team of women processing astronomy data)
- **Cecilia Payne** (early 20th century) was first to accurately relate these classes to temperatures using ionisation theory (she also was the first to claim stars are predominantly hydrogen)

Elemental abundances

Elemental abundances

- Abundances are measured by relative number fraction (not mass), compared to the Sun

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_\odot$$

Elemental abundances

- Abundances are measured by relative number fraction (not mass), compared to the Sun
- Usually we use [Fe/H] to denote “metallicity” (because **iron** is relatively easy to measure and generally traces overall metallicity well, and **hydrogen** is mostly constant across stars)

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_\odot$$

Sun: [Fe/H] = 0

metal-poor: [Fe/H] < -1

very metal-poor: [Fe/H] < -2

Elemental abundances

- Abundances are measured by relative number fraction (not mass), compared to the Sun
- Usually we use [Fe/H] to denote “metallicity” (because **iron** is relatively easy to measure and generally traces overall metallicity well, and **hydrogen** is mostly constant across stars)
- Other elemental abundances are typically given as a ratio to iron, e.g. [Mg/Fe] or [C/Fe].

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_{\odot}$$

Sun: [Fe/H] = 0

metal-poor: [Fe/H] < -1

very metal-poor: [Fe/H] < -2

Elemental abundances


- Abundances are measured by relative number fraction (not mass), compared to the Sun
- Usually we use [Fe/H] to denote “metallicity” (because **iron** is relatively easy to measure and generally traces overall metallicity well, and **hydrogen** is mostly constant across stars)
- Other elemental abundances are typically given as a ratio to iron, e.g. [Mg/Fe] or [C/Fe].

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_{\odot}$$

Sun: [Fe/H] = 0

metal-poor: [Fe/H] < -1

very metal-poor: [Fe/H] < -2



APOGEE data
(high- resolution
infrared spectra,
R~22 500)

Elemental abundances



Where is the Sun?


- Abundances are measured by relative number fraction (not mass), compared to the Sun
- Usually we use [Fe/H] to denote “metallicity” (because **iron** is relatively easy to measure and generally traces overall metallicity well, and **hydrogen** is mostly constant across stars)
- Other elemental abundances are typically given as a ratio to iron, e.g. [Mg/Fe] or [C/Fe].

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_{\odot}$$

Sun: [Fe/H] = 0

metal-poor: [Fe/H] < -1

very metal-poor: [Fe/H] < -2



APOGEE data
(high-resolution
infrared spectra,
R~22 500)

Elemental abundances



Where is the Sun?


- Abundances are measured by relative number fraction (not mass), compared to the Sun
- Usually we use [Fe/H] to denote “metallicity” (because **iron** is relatively easy to measure and generally traces overall metallicity well, and **hydrogen** is mostly constant across stars)
- Other elemental abundances are typically given as a ratio to iron, e.g. [Mg/Fe] or [C/Fe].

$$\left[\frac{X}{H} \right] = \log_{10} \left(\frac{N_X}{N_H} \right)_* - \log_{10} \left(\frac{N_X}{N_H} \right)_{\odot}$$

Sun: [Fe/H] = 0

metal-poor: [Fe/H] < -1


very metal-poor: [Fe/H] < -2



APOGEE data
(high-resolution
infrared spectra,
R~22 500)

Elemental abundances


Mg is an “alpha” element, made in type II supernovae (quickly exploding massive stars) and in type Ia supernovae (exploding white dwarfs = long-lived low-mass star remnants)



~270 000 stars from APOGEE
(high-resolution infrared spectra,
 $R \sim 22\,500$)

Elemental abundances

Mg is an “alpha” element, made in type II supernovae (quickly exploding massive stars) and in type Ia supernovae (exploding white dwarfs = long-lived low-mass star remnants)




Solar level
mix of type II & type Ia supernovae

~270 000 stars from APOGEE
(high-resolution infrared spectra,
 $R \sim 22\,500$)

Elemental abundances

Mg is an “alpha” element, made in type II supernovae (quickly exploding massive stars) and in type Ia supernovae (exploding white dwarfs = long-lived low-mass star remnants)



How to derive abundances?

Different types of datasets:

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:

- Line-by-line measurements

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:

- Line-by-line measurements
- Full-spectrum fitting

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:

- Line-by-line measurements
- Full-spectrum fitting
- Label transfer

How to derive abundances?

Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:

- Line-by-line measurements
- Full-spectrum fitting
- Label transfer
- Model-driven vs. data-driven

How to derive abundances?


Different types of datasets:

- Small high-quality, high-resolution datasets
- Medium-sized surveys
- Large surveys at low resolution
- Large surveys at medium/high resolution
- Large datasets with photometry/very low resolution

Some techniques/choices:


- Line-by-line measurements
- Full-spectrum fitting
- Label transfer
- Model-driven vs. data-driven
- Purely spectroscopic or include external data?

Line-by-line measurements




Line-by-line measurements

- Only for **high-resolution spectra** ($R \sim > 10\,000$)



Line-by-line measurements


- Only for **high-resolution spectra** ($R \sim > 10\,000$)
- **Measure EWs** of all individual desired lines



Line-by-line measurements

- Only for **high-resolution spectra** ($R \sim > 10\,000$)
- **Measure EWs** of all individual desired lines
- Guess, assume or derive **stellar parameters** (T_{eff} , $\log g$, $[M/H]$)


Nissen & Schuster (2010)



Line-by-line measurements

- Only for **high-resolution spectra** ($R \sim > 10\,000$)
- **Measure EWs** of all individual desired lines
- Guess, assume or derive **stellar parameters** (T_{eff} , $\log g$, $[M/H]$)
- Choose **stellar atmosphere model and spectral synthesis code** to solve abundance equations, perform radiative transfer & create synthetic spectra


Nissen & Schuster (2010)



Line-by-line measurements

- Only for **high-resolution spectra** ($R \sim > 10\,000$)
- **Measure EWs** of all individual desired lines
- Guess, assume or derive **stellar parameters** (T_{eff} , $\log g$, $[M/H]$)
- Choose **stellar atmosphere model and spectral synthesis code** to solve abundance equations, perform radiative transfer & create synthetic spectra
- Place abundances on a scale (typically Solar)

Nissen & Schuster (2010)





Line-by-line measurements


- Only for **high-resolution spectra** ($R \sim > 10\,000$)
- **Measure EWs** of all individual desired lines
- Guess, assume or derive **stellar parameters** (T_{eff} , $\log g$, $[M/H]$)
- Choose **stellar atmosphere model and spectral synthesis code** to solve abundance equations, perform radiative transfer & create synthetic spectra
- Place abundances on a scale (typically Solar)

At the same metallicity: a group of stars formed in an **efficient star-forming environment** and a group from an **inefficient star-forming environment**

Nissen & Schuster (2010)




Full-spectrum fitting



*Fit of a low-resolution
($R \sim 1300$) spectrum with
an empirical library
(Arentsen+20)*

Full-spectrum fitting


- **Any spectral resolution** (often used at lower R to extract as much information as possible)



*Fit of a low-resolution
($R \sim 1300$) spectrum with
an empirical library
(Arentsen+20)*

Full-spectrum fitting


- **Any spectral resolution** (often used at lower R to extract as much information as possible)
- Need a **spectral grid** (typically pre-computed synthetic spectra with varying stellar parameters and/or abundances, can also use empirical libraries) and an **interpolation tool**



*Fit of a low-resolution
($R \sim 1300$) spectrum with
an empirical library
(Arentsen+20)*

Full-spectrum fitting


- **Any spectral resolution** (often used at lower R to extract as much information as possible)
- Need a **spectral grid** (typically pre-computed synthetic spectra with varying stellar parameters and/or abundances, can also use empirical libraries) and an **interpolation tool**
- Not measuring individual EWs, but trying to **find the best-matching template** to the entire spectrum (or relevant spectral windows) —> many possible ways of doing this



*Fit of a low-resolution
($R \sim 1300$) spectrum with
an empirical library
(Arentsen+20)*

Full-spectrum fitting

- **Any spectral resolution** (often used at lower R to extract as much information as possible)
- Need a **spectral grid** (typically pre-computed synthetic spectra with varying stellar parameters and/or abundances, can also use empirical libraries) and an **interpolation tool**
- Not measuring individual EWs, but trying to **find the best-matching template** to the entire spectrum (or relevant spectral windows) —> many possible ways of doing this
- Tractable for large spectroscopic surveys (e.g. used in APOGEE, GALAH, LAMOST, Gaia RVS)




*Fit of a low-resolution
($R \sim 1300$) spectrum with
an empirical library
(Arentsen+20)*

Label-transfer

- **Any spectral resolution**
- Learn **attributes of spectra (flux)** based on **stellar parameters and abundances (labels)**
- Requires a **reference training set** to calibrate (e.g. overlap with another survey, or a high-quality analysis of a subset of stars)
- Can for example also be used to put labels from different surveys on the same “scale”


Label-transfer

- Any spectral resolution
- Learn **attributes of spectra (flux)** based on **stellar parameters and abundances (labels)**
- Requires a **reference training set** to calibrate (e.g. overlap with another survey, or a high-quality analysis of a subset of stars)
- Can for example also be used to put labels from different surveys on the same “scale”




Label-transfer

- Any spectral resolution
- Learn **attributes of spectra (flux)** based on **stellar parameters and abundances (labels)**
- Requires a **reference training set** to calibrate (e.g. overlap with another survey, or a high-quality analysis of a subset of stars)
- Can for example also be used to put labels from different surveys on the same “scale”




Generative




Label-transfer example

- Method: **XGBoost**, gradient boosted decision trees (similar to random forest)




Label-transfer example

- Method: **XGBoost**, gradient boosted decision trees (similar to random forest)
- Data to be labeled: **Gaia XP spectra** (very low resolution, $R \sim 30-100$)





Label-transfer example

- Method: **XGBoost**, gradient boosted decision trees (similar to random forest)
- Data to be labeled: **Gaia XP spectra** (very low resolution, $R \sim 30-100$)
- Training data: labels from **APOGEE** high-res spectroscopic survey ($R \sim 22\,500$)




Creevey et al. (2022)






Label-transfer example

- Method: **XGBoost**, gradient boosted decision trees (similar to random forest)
- Data to be labeled: **Gaia XP spectra** (very low resolution, $R \sim 30\text{-}100$)
- Training data: labels from **APOGEE** high-res spectroscopic survey ($R \sim 22\,500$)




Creevey et al. (2022)



Using additional data?

For example:



- **Photometry (colours and magnitudes):** constrain spectral energy distribution \rightarrow T_{eff}



Using additional data?

For example:

- **Photometry (colours and magnitudes):** constrain spectral energy distribution \rightarrow T_{eff}
- **Parallax:** constrain the distance and absolute magnitude \rightarrow $\log g$






Using additional data?


For example:

- **Photometry (colours and magnitudes):** constrain spectral energy distribution \rightarrow T_{eff}
- **Parallax:** constrain the distance and absolute magnitude \rightarrow $\log g$
- **Two types of spectra** at the same time: e.g. Gaia XP (R~30-100 but covering the full optical) and Gaia RVS (R~7000 but only a small range in the infrared)

for example Guiglion et al. (2024), who use a convolutional neural network



+



Summary

- Spectra contain a lot of information about the composition of stars
- The analysis of spectra is not straightforward, and there are many methods
- Elemental abundances can be used to trace the history of galaxies


Summary

- Spectra contain a lot of information about the composition of stars
- The analysis of spectra is not straightforward, and there are many methods
- Elemental abundances can be used to trace the history of galaxies

If you want to follow up some of the references, in astronomy we typically use the Astrophysics Data System (ADS) to search for publications:

<https://ui.adsabs.harvard.edu/>

(e.g. searching for author name & year)



The screenshot shows the ADS search interface. At the top, there's a header with the ADS logo and the text "astrophysics data system". Below the header, there are three tabs: "Classic Form", "Modern Form" (which is selected), and "Paper Form". Under the "Modern Form" tab, there's a "QUICK FIELD:" section with a dropdown menu showing "Author", "First Author", "Abstract", "Year", "Fulltext", and "All Search Terms". Below this is a search input field with a magnifying glass icon. To the right of the search field, there's a "Recommendations" section with a "Search examples" sub-section. The "Search examples" section contains several search queries in a table-like format:

Field	Search Example
author	author:"Lee, Eve"
first author	first_author:"Obrzud, Ewelina"
abstract + title	abs:"dark energy"
year	year:2000
year range	year:2000-2005
full text	full:"super Earth"
refereed	property:refereed
astronomy	collection:astronomy
exact search	=body:"intracluster medium"
institution	inst:CfA
author count	author_count:[1 TO 10]
record type	doctype:software

Next lectures

- Friday: How to design and obtain astronomical observations?
- Next Tuesday: part chemical evolution & recent results, and part hands-on
 - bring your laptop if you can