

<b>1</b>	<b>Introduction.....</b>	<b>1</b>
1.1	Motivation.....	2
1.2	Rydberg Atoms and Dipole-Dipole Interactions.....	3
1.3	Atomic Units .....	3
1.4	Dissertation Structure .....	4
<b>2</b>	<b>Experimental Setup.....</b>	<b>8</b>
2.1	Magneto-Optical Trap .....	9
2.1.1	Principle of the Magneto-Optical Trap.....	9
2.1.2	Saturated Absorption Spectroscopy.....	13
2.1.3	Ultra High Vacuum Chamber .....	16
2.1.4	Characterization of the MOT .....	16
2.2	Lasers and Amplifiers .....	17
2.2.1	Nd:YAG Lasers.....	17
2.2.2	Diode Lasers.....	19
2.2.3	Mode-Lock Laser.....	19
2.2.4	Chirped Pulse Amplification .....	21
2.2.5	Regenerative Amplifier .....	23
2.2.6	Linear Amplifier .....	24
2.2.7	Dye Laser and Dye Amplifier .....	25
2.3	THz Pulses.....	27
2.4	Detection and Data Collection .....	30
2.4.1	State-Selective Field Ionization .....	30
2.4.2	Synchronization System .....	31
2.4.3	Measurement Operation.....	33

<b>2.5</b>	<b>Maintenance and Daily Operation .....</b>	<b>34</b>
2.5.1	Daily Examination .....	34
2.5.2	Operation of Regenerative Amplifier .....	35
2.5.3	Operation of the MOT .....	36
<b>3</b>	<b>Models in Simulation .....</b>	<b>41</b>
<b>3.1</b>	<b>Rydberg Atoms .....</b>	<b>42</b>
3.1.1	Modern Picture of Rydberg Atoms.....	43
3.1.2	Properties of Rydberg Atoms .....	45
<b>3.2</b>	<b>Two-Atom Model.....</b>	<b>46</b>
3.2.1	Introduction to the Two-Atom Model.....	46
3.2.2	Nearest Neighbor Distribution .....	47
3.2.3	Förster Resonant Energy Transfer.....	48
<b>3.3</b>	<b>Dipole-Dipole Interaction .....</b>	<b>49</b>
3.3.1	Dipole Moment.....	49
3.3.2	Dipole-Dipole Interaction in the Classical Picture .....	50
3.3.3	Dipole-Dipole Interaction in the Quantum Picture .....	52
3.3.4	Dipole-Dipole Coupled System in an Electric Field.....	55
<b>3.4</b>	<b>Rydberg Decay Model .....</b>	<b>56</b>
3.4.1	Blackbody Induced Transitions.....	57
3.4.2	Radiative Transitions .....	57
<b>4</b>	<b>Absence of Collective Decay in a Cold Rydberg Gas .....</b>	<b>61</b>
<b>4.1</b>	<b>Introduction.....</b>	<b>62</b>
<b>4.2</b>	<b>Experimental Procedure and Results .....</b>	<b>65</b>
<b>4.3</b>	<b>Analysis and Discussion.....</b>	<b>70</b>

4.4	Conclusion .....	79
5	Rydberg Wavepacket Evolution in A Frozen Gas of DD Coupled Atoms.....	85
5.1	Introduction.....	86
5.2	Experimental Procedure.....	88
5.3	Experimental Results.....	90
5.4	Discussion.....	93
5.5	Conclusion .....	97
6	Wavepacket Coherence Transfer via DD interactions .....	102
7	Summary and Future Directions .....	105

Figure 2.1: Simplified one dimensional model for MOT [2]. .....	10
Figure 2.2: Schematic for MOT design. It's a combination of anti-Helmholtz coils and six counter-propagating laser beams. ....	11
Figure 2.3: Hyperfine energy structure of $^{85}\text{Rb}$ . Trap laser drives the transition, $5s_{1/2} F=3$ to $5p_{3/2} F=4$ , and the repump laser drives the transition, $5s_{1/2} F=2$ to $5p_{3/2} F=3$ . ....	13
Figure 2.4: The saturated absorption spectra for the Rb $5s_{1/2}$ to $5p_{1/2}$ hyperfine transitions and their crossover peaks [3]. Small dark arrows are pointing to the real resonances. Blue dotted arrow, pointing to the side of the crossover peak, indicates where the trap laser is actually locked at. Red dotted arrow indicates the position where the repump laser is locked at. ....	15
Figure 2.5: Schematic of Nd:YAG lasing [6]. It is a typical four-level lasing scheme. ....	18
Figure 2.6: Layout of Continuum Surelite Nd:YAG laser. It can output beams of 4 different frequencies [5]. ....	18
Figure 2.7: Basic layout of the mode-lock laser [12]. The green line is the pump light from Millennia Vs laser and the red line is the mode-locked beam in the cavity which is centered at 780-800 nm. The prism pair is used to compensate for group velocity dispersion (GVD). By tapping the 2 <sup>nd</sup> prism, we can produce a temporary unstable beam. Stronger intensity part in this beam will be enhanced, producing the periodic pulsed output. ....	20
Figure 2.8: Schematic of chirped pulse amplification system. Seed light is first temporally stretched using the stretcher. Then the stretched pulse gets amplified. Finally the pulse	

is again compressed to something close to its original duration, but with much greater (106 – 107 times) energy. ....	22
Figure 2.9: Schematic of regenerative amplifier. The switch-in pockels cell controls when a seed pulse is trapped in the resonator and the switch-out pockels cell controls when the pulse is ejected from the cavity. ....	24
Figure 2.10: Schematic of linear amplifier. The beam passes the Nd:YAG pumped gain medium three times and gets amplified. ....	25
Figure 2.11: Schematic for a Hansch dye laser and the 2 <sup>nd</sup> harmonic generation. The angle of the tuning grating determines the output frequency. ....	26
Figure 2.12: Schematic for double cell dye amplifier used in the lab. ....	27
Figure 2.13: A typical THz pulse generated in the lab [8]. The duration is about 5 ps and the peak frequency is 0.2-0.4 THz. This pulse is generated using tilted-pulse-front- pumping optical rectification and the temporal profile of the pulse is characterized using the electro-optic sampling method. The frequency spectrum is then derived from the time-domain profile. ....	28
Figure 2.14: Schematic of TPFP setup [8]. ....	29
Figure 2.15: Schematic of the tipping of electron potential. Solid line is the $1/r$ potential when there is no external field applied to the atom. Dashed line shows the tip of potential when a field is applied to the atom. When such a field is strong enough, electrons are able to escape from the trap. ....	31
Figure 2.16: Schematic of the synchronization system. White circles are inputs and dark circles are outputs. ....	32

- Figure 2.17: A typical ionization signal shown on an oscilloscope. The central peak representing the population of state  $32s + 32p$ . The measurement program uses a user defined gate and integrates the area under the peak within the gate..... 33
- Figure 2.18: Front view of the chamber where the MOT is positioned in. A, B, and C represent three trap beams. .... 38
- Figure 3.1: Classical view of Rydberg orbits of (a) H and (b) Na. In H the electron orbits around the proton. In Na it orbits around the +11 nuclear charge and ten inner shell electrons. In high  $L$  states Na behaves nearly identically to H, but in low  $L$  states the Na electron penetrates and polarizes the inner shell electrons of the Na + core [5]. 43
- Figure 3.2: Schematic of atom pairs in a MOT. For each atom in the MOT, we only consider the effect of its nearest neighbor. One atom and its nearest neighbor is considered to be “a pair of atoms”. .... 47
- Figure 3.3: Schematic for typical FRET. Black cycles represent the initial pair states and gray cycles the final pair states. (a) is  $pp \rightarrow ss'$ , (b)  $ps \rightarrow sp$  and (c)  $ps' \rightarrow s'p$  [13]..... 49
- Figure 3.4: Schematic of the interaction between two classical dipoles. .... 52
- Figure 3.5: Schematic for a DD system in an electric field..... 55
- Figure 3.6: Decay model for atoms starting from state  $40s$  as an example. The red dash lines between two states indicate blackbody stimulated transitions between these two states. The blue dash curves represent the spontaneous decay. .... 58
- Figure 4.1: (a), (c) Probabilities for finding atoms in  $26s + 25p$  (green, fastest decay),  $32s$  (red, intermediate decay), and  $40s$  (blue, slowest decay) as a function of detection time  $\tau$  for Rydberg densities of  $\rho \sim 3 \times 10^9 \text{ cm}^{-3}$  (a) and  $\rho \sim 1.5 \times 10^8 \text{ cm}^{-3}$  (c). Note that the sum of the  $26s$  and  $25p$  populations is shown since their corresponding features

could not be adequately separated in the field-ionization signal. Vertical bars show the experimental data with uncertainties, and the solid curves are calculated as described in the text. Measurements and calculations for the  $40s$  decay extend to  $500\ \mu s$  where the remaining population is negligible. (b), (d) Probabilities for finding atoms in  $26p$  (green, fastest rise and decay),  $32p$  (red, intermediate rise and decay), and  $40p$  (blue, slowest rise and decay) levels as a function of detection time  $\tau$ . The states are populated by blackbody redistribution from the initial  $26s$ ,  $32s$ , and  $40s$  levels, respectively. The data were measured simultaneously with those shown in (a) and (c). Vertical bars show the experimental data with uncertainties, and the solid curves are calculated as described in the text. The measured  $p$ -state probabilities are normalized to the calculations as described in the text. The calculations have no free parameters and consider only the effects of spontaneous emission and blackbody radiation on isolated atoms. .... 68

Figure 4.2: Probabilities for finding atoms in  $26p$  (green, fastest decay),  $32p$  (red, intermediate decay), and  $40p$  (blue, slowest decay) as functions of detection time  $\tau$  for Rydberg densities of  $\rho \sim 3 \times 10^9\ \text{cm}^{-3}$  (a) and  $\rho \sim 1.5 \times 10^9\ \text{cm}^{-3}$  (b). Vertical bars show the experimental data with uncertainties, and the solid curves are calculated as described in the text. Measurements and calculations for the  $40p$  decay extend to  $500\ \mu s$  where the remaining population is negligible. The calculations have no free parameters and consider only the effects of spontaneous emission and blackbody radiation on isolated atoms. .... 70

Figure 4.3: Measured  $32p_{3/2}\ |mj| = 1/2$  (bold line) and  $|mj| = 3/2$  (thin line) excitation probabilities as a function of Rydberg laser frequency in zero applied field. The two data curves are obtained simultaneously in the same laser frequency scan. The small feature on the left (right) of the main  $|mj| = 1/2$  ( $3/2$ ) peak is the result of imperfect discrimination of the  $|mj| = 1/2$  and  $3/2$  components via SSFI. The additional peak on

the right of the main feature in each trace is due to the trap-laser dressing of the $5p_{3/2}$ and $5s$ levels. Its frequency shift from the main peak reflects the Autler-Townes splitting of the $5p_{3/2}$ initial state.....	76
Figure 4.4: Difference (i.e., splitting) in the transition energies for exciting $3p_{3/2}  mj =1/2, 3/2$ from $5p_{3/2}$ as a function of applied electric field. Filled circles are measurements and the solid curve is the result of a numerical Stark map calculation assuming orthogonal “offset” and “residual” electric field components due to the MCP of 2.8 and 1.5 V/cm, respectively. The inset shows a magnified view of the portion of the main figure within the dashed window. ....	77
Figure 5.1: Measured population in the combined $3s+3p$ states as a function of the delay $\Delta t$ between two THz pulses. The left panels show data collected at low Rydberg density, $\rho \sim 3 \times 10^8 \text{ cm}^{-3}$ , for (a) short ( $\Delta t \simeq 0$ ) and (b) long ( $\Delta t \simeq 15\text{ns}$ ) delays, respectively. The right panels show data collected at high Rydberg density, $\rho \sim 2 \times 10^9 \text{ cm}^{-3}$ , for c) short ( $\Delta t \simeq 0$ ) and d) long ( $\Delta t \simeq 15\text{ns}$ ) delays, respectively. The decrease in oscillation amplitude at high density and long delays is apparent. ....	91
Figure 5.2: Fast Fourier transform (FFT) of the delay-dependent populations shown in Figure 5.1. ....	92
Figure 5.3: Measured (filled circles) and simulated (solid curve) decay ratio, $\eta$ , as a function of Rydberg density. ....	93
Figure 5.4: Schematic energy level diagram for the eigenstates of a pair of two level atoms. The diagrams on the left and right sides of the figure depict the situation at large and small interatomic spacing, $R$ , respectively. ....	95



# 1 Introduction

The electronic and nuclear motion degrees of freedom in Rydberg atoms are coupled by long-range dipole-dipole interactions [1]. This coupling is responsible for rich few- and many-body quantum dynamics and its coherent manipulation can enable potential applications in the quantum control of few- and many-body systems, quantum information processing, quantum computing, etc. [2-10]. The research discussed in this dissertation focuses on the influence of dipole-dipole (DD) interactions on electron dynamics within cold Rydberg atoms. This is experimentally challenging due to the long separation of time and distance scales associated with the electronic and nuclear motion, respectively. As summarized below, the result of three sub-projects are described in this dissertation. Details of the common experimental setups, computational approaches, and background knowledge are also provided.

## 1.1 Motivation

Our group’s primary research involves the exploration and manipulation of quantum dynamics in atomic and molecular systems. This dissertation describes a series of sub-projects designed to further this effort. Our results provide new insight into the fundamental problems in atomic physics. More than that, they may contribute to practical applications from designing many-body systems that simulate model condensed matter systems, to quantum information storage and processing, to the development of new radiation sources and detectors.

Recent work done by previous graduates Xiangdong Zhang [11] and Mary Kutteruf [12] provided much of the foundation and motivation for the work presented here. The project “Probing Time-Dependent Electron Interactions in Double Rydberg Wavepackets” done by Xiangdong Zhang helps us understand more about time-dependent electron-electron interactions within individual atoms. The project “Coherence in Rydberg Atoms: Measurement and Control” done by Mary Kutteruf explored the use electric fields to control and measure coherence in electronic wavepackets and between coupled atoms in Rydberg ensembles. My project is an extension of their work but is the first to explicitly examine the impact of dipole-dipole interactions between atoms on electron wavepacket dynamics within them. Another current graduate student Brian Richards is working on utilizing controlled DD interactions to manipulate the position correlation function of cold trapped atoms. Our projects are like puzzle chunks, working together to develop new capabilities and make the scientific picture more complete.

My project exploring the influence of dipole-dipole interactions on Rydberg wavepackets has been divided into three sub-projects. The first one explores the role of DD interactions in suppressing collective decay (“superradiance”) in an ensemble of cold Rydberg atoms (Chapter 4). The second one characterizes the role of DD interactions in

dephasing the macroscopic coherence of an ensemble of Rydberg wavepackets (Chapter 5). The final one examines the transfer of wavepacket coherence between atoms via DD interactions (Chapter 6). These three sub-projects help us understand better how the DD interactions influence electron dynamics in an ensemble, and explore the use of controlled atom-atom coupling to induce coherent wavepacket motion within those atoms. The latter provides an effective demonstration of coherent control beyond unimolecular photo-reactions.

## 1.2 Rydberg Atoms and Dipole-Dipole Interactions

Rydberg atoms are atoms in which an electron is excited to a state with a high principal quantum number. Usually this number is larger than 10. They are good systems for studying atom dynamics. The details about Rydberg atoms and their properties can be found in references such as [1], and are described in Chapter 3.

Dipole-dipole (DD) interactions can influence neighboring Rydberg atoms. Classically the interaction is the result of the electric forces between electrons and the positively charged ion cores to which they are bound. When coupled by DD interactions, Rydberg atoms should not be considered as individuals but rather a system. Details of DD interactions, both from a classical view and a quantum physics perspective, can be found in Chapter 3.

## 1.3 Atomic Units

Atomic units (au or a.u.) are commonly used in atomic physics research. For convenience, we define:

$$\hbar = m_e = e = 4\pi\epsilon_0 = 1 \quad (1.1)$$

where  $\hbar$  is Planck's constant divided by  $2\pi$ ,  $m_e$  is the mass of the electron,  $-e$  is the electron charge and  $\varepsilon_0$  is the permittivity of free space. The conversion factors between a.u. and SI units are shown in Table 1.1.

Quantity	Value in atomic units	Value in SI units
Length	1	$5.2917721092(17) \times 10^{-11}$ m
Energy	1	$4.35974417(75) \times 10^{-18}$ J
Time	1	$2.418884326505(16) \times 10^{-17}$ s
Velocity	1	$2.1876912633(73) \times 10^6$ m·s <sup>-1</sup>
Force	1	$8.2387225(14) \times 10^{-8}$ N
Temperature	1	$3.1577464(55) \times 10^5$ K
Pressure	1	$2.9421912(19) \times 10^{13}$ Pa
Electric field	1	$5.14220652(11) \times 10^{11}$ V·m <sup>-1</sup>
Electric potential	1	$2.721138505(60) \times 10^1$ V
Electric dipole moment	1	$8.47835326(19) \times 10^{-30}$ C·m
Magnetic field	1	$2.35 \times 10^5$ T

Table 1.1: Atomic units to SI units conversion factors.

## 1.4 Dissertation Structure

Subsequent chapters describe the experimental approach, numerical simulations and several distinct projects. Each project contains both an experimental description and details on relevant simulations. Additional introductory material and common aspects of all experiments are found in chapters 2 and 3.

Chapter 2 provides the information about experimental setups. It introduces the apparatus commonly used in the experiments, as well as daily operation procedures. More details about some instruments can be found in their respective manuals and the dissertations from previous students who worked in this lab.

Chapter 3 introduces physics concepts commonly involved in the experiments, and their mathematical expression in simulations. It is not practical to include in this dissertation every single line of simulation code that was used during the research, but by following the models described in Chapter 3, one could reconstruct the simulations in a fairly straightforward way.

Chapter 4 describes a search for collective decay (i.e. “superradiance”) in cold Rydberg gases. We found no evidence for superradiance, despite the results reported by other groups. Our analysis suggests that, due to dipole-dipole dephasing, our null result is the expected one for highly excited Rydberg atoms in the typical magneto-optical-trap geometry.

Chapter 5 describes an exploration of Rydberg wavepacket evolution in dipole-dipole coupled atoms. Our results show, through experiment and simulation, that in dipole-dipole coupled atoms, the wavepackets do not evolve independently, and variations in the coupling strength between pairs of atoms lead to macroscopic dephasing of the electronic wavepacket motion.

Chapter 6 describes a study of coherence transfer between wavepackets on different atoms via resonant dipole-dipole interactions. We observe evidence for the development of wavepacket motion in one set of atoms driven by wavepacket evolution in neighboring atoms. We confirm the coherence transfer by measuring the relative phase between the initial and induced wavepacket oscillations, suggesting the creation of entangled atom pair states with a dynamically evolving Rydberg wavepacket on one, and only one, atom in each pair. Simulations support our interpretation.

Chapter 7 summarizes the work done in this dissertation and briefly discusses the possibilities for future experiments following these results.

# Bibliography

- [1] Thomas F. Gallagher. Rydberg Atoms. Cambridge University Press (1994).
- [2] M. D. Lukin, M. Fleischhauer, R. Cote, L. M. Duan, D. Jaksch, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. 87, 037901 (2001).
- [3] D. Tong, S. M. Farooqi, J. Stanojevic, S. Krishnan, Y. P. Zhang, R. Cote, E. E. Eyler, and P. L. Gould, Phys. Rev. Lett. 93, 063001 (2004).
- [4] F. Robicheaux and J. V. Hernandez, Phys. Rev. A 72, 063403 (2005).
- [5] T. Cubel Liebisch, A. Reinhard, P. R. Berman, and G. Raithel, Phys. Rev. Lett. 95, 253002 (2005).
- [6] T. Vogt, M. Viteau, J. Zhao, A. Chotia, D. Comparat, and P. Pillet, Phys. Rev. Lett. 97, 083003 (2006).
- [7] E. Urban, T. A. Johnson, T. Henage, L. Isenhower, D. D. Yavuz, T. G. Walker, and M. Saffman, Nat. Phys. 5, 110 (2009).
- [8] Alpha Gaëtan, Yevhen Miroshnychenko, Tatjana Wilk, Amodsen Chotia, Matthieu Viteau, Daniel Comparat, Pierre Pillet, Antoine Browaeys, and Philippe Grangier, Nat. Phys. 5, 115 (2009).
- [9] T. Wilk, A. Gaetan, C. Evellin, J. Wolters, Y. Miroshnychenko, P. Grangier, and A. Browaeys, Phys. Rev. Lett. 104, 010502 (2010).
- [10] D. Jaksch, J.I. Cirac, P. Zoller, S.L. Rolston, R. Cote, and M.D. Lukin, Phys. Rev. Lett. 85, 2208 (2000).
- [11] Xiangdong Zhang, Probing Time-Dependent Electron Interactions in Double Rydberg Wavepackets, PhD thesis, University of Virginia (2008).
- [12] Mary Kutteruf, Coherence in Rydberg Atoms: Measurement and Control, PhD thesis, University of Virginia (2010).

- [13] A. Dalgarno, *Rydberg States of Atoms and Molecules*, eds. R. F. Stebbings and F.B. Dunning, Cambridge University Press (1983).
- [14] H. A. Bethe and E.A. Salpeter, *Quantum Mechanics of One and Two Electron Atoms*, Academic Press (1957).

## 2 Experimental Setup

This chapter contains the general experimental setup for the research discussed in this dissertation. It introduces the apparatus and procedure for state excitation, laser cooling, pulse amplification, THz pulse generation, data collection, etc. It also provides procedures for maintenance and daily operation. All experiments are performed on Newport RS 3000 optical tables to reduce mechanical vibrations, in a temperature controlled room to reduce external thermal fluctuations. Other than specifically noted, the repetition rate of all experiments is 15 Hz. Before beginning experiments in the lab, participants must have taken the laboratory safety training.