
EECS 16A
Fall 2018

Designing Information Devices and Systems I

Discussion 12B

1. Least Squares: A Toy Example

Let's start off by solving a little example of least squares.

We're given the following system of equations:

$$\begin{bmatrix} 1 & 4 \\ 3 & 8 \\ 5 & 16 \end{bmatrix} \vec{x} = \begin{bmatrix} 3 \\ 1 \\ 9 \end{bmatrix},$$

where $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$.

- (a) Why can we not solve for \vec{x} exactly?

Answer:

Recall from the earlier linear algebra module that in order for there to be a solution for the matrix system $\mathbf{A}\vec{x} = \vec{b}$, we must have $\vec{b} \in \text{Col}(\mathbf{A})$.

Let us use Gaussian elimination to see if we can find \vec{x} .

$$\left[\begin{array}{cc|c} 1 & 4 & 3 \\ 3 & 8 & 1 \\ 5 & 16 & 9 \end{array} \right] \xrightarrow{R_3 - 2R_1 - R_2 \rightarrow R_3} \left[\begin{array}{cc|c} 1 & 4 & 3 \\ 3 & 8 & 1 \\ 0 & 0 & 2 \end{array} \right]$$

We have reached a point at which there does not exist an \vec{x} that exactly solves the system of equations. Thus, in this case $\vec{b} \notin \text{Col}(\mathbf{A})$. This is because of the last row of \mathbf{A} is $[0 \ 0] \vec{x} = 2$.

- (b) Find $\hat{\vec{x}}$, the *least squares estimate* of \vec{x} , using the formula we derived in lecture.

Reminder: $\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$

Answer:

Recall the equation to find the linear least squares estimate:

$$\hat{\vec{x}} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \vec{b}$$

Plugging in $\mathbf{A} = \begin{bmatrix} 1 & 4 \\ 3 & 8 \\ 5 & 16 \end{bmatrix}$ and $\vec{b} = \begin{bmatrix} 3 \\ 1 \\ 9 \end{bmatrix}$, we get $\hat{\vec{x}} = \begin{bmatrix} -6 \\ 2.41\bar{6} \end{bmatrix}$.

2. Polynomial Fitting

Least squares can only be applied to linear systems. Suppose that we have a vector \vec{x} and a vector \vec{y} , and $\vec{y}[n] = f(\vec{x}[n])$. We would like to approximate f using least squares, where f is not necessarily a linear function.

- (a) Suppose that $y = ax + b$. Set this up as a least squares problem. What are the elements in the matrix \mathbf{A} ?

Answer:

$$\begin{bmatrix} \vec{x} & \vec{1} \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \vec{y}$$

Let's try an example. Say we know that the output, y , is a quartic polynomial in x . This means that we know that y and x are related as follows:

$$y = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$$

We're also given the following observations:

x	y
0.0	24.0
0.5	6.61
1.0	0.0
1.5	-0.95
2.0	0.07
2.5	0.73
3.0	-0.12
3.5	-0.83
4.0	-0.04
4.5	6.42

- (b) What are the unknowns in this question? What are we trying to solve for?

Answer:

The unknowns are a_0 , a_1 , a_2 , a_3 , and a_4 . They are also what we are trying to solve for.

- (c) Can you write an equation corresponding to the first observation (x_0, y_0) , in terms of a_0 , a_1 , a_2 , a_3 , and a_4 ? What does this equation look like? Is it linear?

Answer:

Plugging (x_0, y_0) into the expression for y in terms of x , we get

$$24 = a_0 + a_1 \cdot 0 + a_2 \cdot 0^2 + a_3 \cdot 0^3 + a_4 \cdot 0^4$$

You can see that this equation is linear in a_0 , a_1 , a_2 , a_3 , and a_4 .

- (d) Now, write a system of equations in terms of a_0 , a_1 , a_2 , a_3 , and a_4 using *all of the observations*.

Answer:

Write the next equation using the second observation. You will now get:

$$6.61 = a_0 + a_1 \cdot (0.5) + a_2 \cdot (0.5)^2 + a_3 \cdot (0.5)^3 + a_4 \cdot (0.5)^4$$

And for the third:

$$0.0 = a_0 + a_1 \cdot (1) + a_2 \cdot 1^2 + a_3 \cdot 1^3 + a_4 \cdot 1^4$$

Do you see a pattern? Let's write the entire system of equations in terms of a matrix now.

$$\begin{bmatrix} 1 & 0 & 0^2 & 0^3 & 0^4 \\ 1 & 0.5 & (0.5)^2 & (0.5)^3 & (0.5)^4 \\ 1 & 1 & 1^2 & 1^3 & 1^4 \\ 1 & 1.5 & (1.5)^2 & (1.5)^3 & (1.5)^4 \\ 1 & 2 & 2^2 & 2^3 & 2^4 \\ 1 & 2.5 & (2.5)^2 & (2.5)^3 & (2.5)^4 \\ 1 & 3 & 3^2 & 3^3 & 3^4 \\ 1 & 3.5 & (3.5)^2 & (3.5)^3 & (3.5)^4 \\ 1 & 4 & 4^2 & 4^3 & 4^4 \\ 1 & 4.5 & (4.5)^2 & (4.5)^3 & (4.5)^4 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = \begin{bmatrix} 24 \\ 6.61 \\ 0.0 \\ -0.95 \\ 0.07 \\ 0.73 \\ -0.12 \\ -0.83 \\ -0.04 \\ 6.42 \end{bmatrix}$$

- (e) Finally, solve for a_0 , a_1 , a_2 , a_3 , and a_4 using IPython. You have now found the quartic polynomial that best fits the data!

Answer:

Let \mathbf{D} be the big matrix from the previous part.

$$\vec{a} = (\mathbf{D}^T \mathbf{D})^{-1} \mathbf{D}^T \vec{y} = \begin{bmatrix} 24.00958042 \\ -49.99515152 \\ 35.0039627 \\ -9.99561772 \\ 0.99841492 \end{bmatrix}$$

It turns out that the actual parameters for the polynomial equation were:

$$\vec{a} = \begin{bmatrix} 24 \\ -50 \\ 35 \\ -10 \\ 1 \end{bmatrix}$$

(Remember that our observations were noisy.)

Therefore, we have actually done pretty well with the least squares estimate!