LANSOPRAZOLE

Orodispersible form

Before breakfast

CHEMICAL NAME (RS)-2-({[3-methyl-4-(2,2,2-trifluoroethoxy)pyridin-2-yl]methyl}sulfinyl)-1H-benzimidazole

CLASS Proton pump inhibitor (PPI)

TRADE NAMES Prevacid · Zoton FasTab · Lanzor · Ogast

THERAPEUTIC USES / INDICATIONS

- GORD, oesophagitis.
- Peptic ulcer; NSAID ulcer prophylaxis.
- *H. pylori* eradication.
- Zollinger–Ellison syndrome.

ADMINISTRATION

- **PO** capsules or orodispersible tablets (FasTab — dissolve on tongue or in water for NG/oral syringe), 30 min before breakfast.

DOSAGES

- **GORD / duodenal ulcer:** 30 mg once daily for 4–8 weeks.
- **Maintenance / prophylaxis:** 15 mg once daily.
- **H. pylori:** 30 mg twice daily with antibiotics.
- **Children:** 0.5–1 mg/kg once daily (max 30 mg).

PHARMACOKINETICS

- **Onset:** 1–2 h. **Duration:** >24 h.
- Metabolised by CYP2C19 and 3A4.
- Excreted in bile and urine. **t½:** 1–2 h.

PHARMACODYNAMICS

- Irreversible **H^+/K^+-ATPase** inhibition.

SIDE & ADVERSE EFFECTS

SIDE EFFECTS (COMMON)

- Diarrhoea, headache.
- Nausea, abdominal pain.
- Rash.

ADVERSE EFFECTS (SERIOUS)

- Microscopic colitis (watery diarrhoea).
- Hypomagnesaemia, B12 deficiency.
- *C. difficile*; fractures.

CONTRAINDICATIONS & CAUTIONS

- Hypersensitivity.
- Use with atazanavir.

KEY INTERACTIONS

- **Clopidogrel:** possible ↓ effect.
- **Theophylline:** ↓ levels (minor).
- **Tacrolimus:** ↑ levels.

NURSING MANAGEMENT

- Teach FasTab use.
- Monitor diarrhoea (microscopic colitis).
- Review long-term need.

PATIENT EDUCATION

- Dissolve on tongue before breakfast; do not chew.
- Report persistent watery diarrhoea.
- Lifestyle measures for reflux.

KEY POINTS

- PPI with orodispersible form (NG friendly).
- Microscopic colitis.
- 30 mg treatment / 15 mg maintenance.
- Before breakfast.