DIOSMIN + HESPERIDIN

Take with meals

Pregnancy: avoid in 1st trimester

CHEMICAL NAME Micronised purified flavonoid fraction: diosmin 90% (3′,5,7-trihydroxy-4′-methoxyflavone 7-rutinoside) + hesperidin 10% (as flavonoids)

CLASS Venotonic / vasoprotective agent (micronised purified flavonoid fraction, MPFF)

TRADE NAMES Daflon 500 / 1000 · Detralex · Venoruton (related) · Phlebodia (diosmin)

THERAPEUTIC USES / INDICATIONS

- Chronic venous insufficiency: leg heaviness, pain, oedema, cramps, varicose veins.
- Acute and chronic haemorrhoids (symptom relief).
- Venous leg ulcers (adjunct to compression).

ADMINISTRATION

- **PO** with meals (lunch and dinner).

DOSAGES

- **Chronic venous disease:** 1 g daily (500 mg twice daily or 1,000 mg once daily).
- **Acute haemorrhoids:** 3 g/day for 4 days, then 2 g/day for 3 days.
- **Chronic haemorrhoids:** 1 g daily.

PHARMACOKINETICS

- Micronisation improves absorption; metabolised by gut flora to aglycones (diosmetin).
- Excreted in urine and faeces. **t½:** ~11 h.

PHARMACODYNAMICS

- Prolongs **noradrenaline** vasoconstrictor effect on vein walls → ↑ venous tone.
- Reduces capillary permeability and leucocyte–endothelium interaction (anti-inflammatory).
- Improves lymphatic drainage.

SIDE & ADVERSE EFFECTS

SIDE EFFECTS (COMMON)

- GI upset: nausea, dyspepsia, diarrhoea.
- Headache, dizziness.
- Rash, itching.

ADVERSE EFFECTS (SERIOUS)

- Allergic reactions (rare).
- Colitis (very rare).

CONTRAINDICATIONS & CAUTIONS

- Hypersensitivity.
- Caution in pregnancy (limited data) and breastfeeding.

KEY INTERACTIONS

- No significant interactions reported.

NURSING MANAGEMENT

- Assess leg oedema, pain, skin changes; measure calf circumference.
- Combine with compression therapy and leg elevation.
- Haemorrhoids: assess bleeding; rule out other causes of rectal bleeding.
- Educate on lifestyle measures.

PATIENT EDUCATION

- Take with meals.
- Wear compression stockings, elevate legs, walk regularly, avoid prolonged standing.
- For haemorrhoids: high-fibre diet and fluids; avoid straining.
- Report rectal bleeding that persists.

KEY POINTS

- Venotonic flavonoid (MPFF).
- 1 g/day; higher short course for acute piles.
- Chronic venous insufficiency and haemorrhoids.
- Use with compression and lifestyle advice.