RABEPRAZOLE

Less CYP2C19 dependence

Swallow whole

CHEMICAL NAME (RS)-2-({[4-(3-methoxypropoxy)-3-methylpyridin-2-yl]methyl}sulfinyl)-1H-benzimidazole (sodium)

CLASS Proton pump inhibitor (PPI)

TRADE NAMES Pariet · Aciphex · Rabez · Razo

THERAPEUTIC USES / INDICATIONS

- GORD, erosive oesophagitis.
- Duodenal and gastric ulcer.
- *H. pylori* eradication.
- Zollinger–Ellison syndrome.

ADMINISTRATION

- **PO** enteric-coated tablet swallowed whole, in the morning (before food for best effect).

DOSAGES

- **GORD / ulcer:** 20 mg once daily for 4–8 weeks.
- **Maintenance:** 10–20 mg once daily.
- **H. pylori:** 20 mg twice daily with antibiotics.
- **Zollinger–Ellison:** 60 mg daily (up to 120 mg).

PHARMACOKINETICS

- **Onset:** <1 h (fastest PPI). **Duration:** >24 h.
- Mainly non-enzymatic reduction — less affected by CYP2C19 polymorphism.
- Excreted in urine. **t½:** 1 h.

PHARMACODYNAMICS

- Irreversible **proton pump** inhibition; rapid activation over wider pH range.

SIDE & ADVERSE EFFECTS

SIDE EFFECTS (COMMON)

- Headache, diarrhoea.
- Nausea, abdominal pain.
- Pharyngitis.

ADVERSE EFFECTS (SERIOUS)

- Hypomagnesaemia, B12 deficiency.
- *C. difficile*, fractures.
- Interstitial nephritis.

CONTRAINDICATIONS & CAUTIONS

- Hypersensitivity.
- Use with atazanavir/rilpivirine.

KEY INTERACTIONS

- **Ketoconazole, atazanavir:** ↓ absorption.
- **Methotrexate:** ↑ levels.
- **Clopidogrel:** less interaction than omeprazole.

NURSING MANAGEMENT

- Give in the morning; do not crush.
- Monitor symptoms and long-term need.

PATIENT EDUCATION

- Swallow whole in the morning.
- Report black stools, persistent pain.

KEY POINTS

- PPI with fastest onset.
- 20 mg once daily.
- Less CYP2C19 variability.
- Swallow whole.

SOURCES CHECKED **[514]** Rabeprazole sodium 10 mg gastro-resistant tablets (UK SmPC) **[515]** Rabeprazole Sodium, oral (US label) *Full references p. 432–457*