12.2 Glaucoma*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I did not know I was losing side vision." **Objective** • Raised intraocular pressure • Peripheral field loss	**ACTUAL** Inadequate health knowledge related to chronic eye disease and lifelong treatment as evidenced by unawareness of the condition.	• Can explain why the drops are needed and that they are lifelong.	• Explain that lost vision will not return, though further loss can be prevented. • Teach instillation followed by nasolacrimal occlusion. • Stress the importance of regular pressure checks. • Advise him to mention his glaucoma before any new medicine is started.	• Considered met when he describes the plan accurately.
Subjective • Acute angle closure: "My eye hurts badly and I see halos." **Objective** • Red, painful eye • Vomiting • Very high intraocular pressure	**ACTUAL** Acute pain related to a sudden rise in intraocular pressure as evidenced by severe ocular pain, halos and vomiting.	• Within hours, pain is relieved and the pressure falls.	• Treat this as an emergency and call the ophthalmologist. • Give the prescribed topical agents and acetazolamide straight away. • Give an antiemetic and analgesia. • Lie him supine, as advised.	• Goal reached once the pain settles.
Subjective *No specific complaint* **Objective** • Visual field loss	**RISK** Risk for physical injury as evidenced by risk factors: reduced peripheral vision.	• No injury.	• Teach him to turn his head to scan his surroundings. • Talk through hazards to remove at home.	• Met if he stays free of injury.

*References for this care plan: [32], [35], [37]. Full entries are listed in the References section of the CPMN book.