14.3 Cardiogenic Shock*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I cannot breathe and feel faint." **Objective** • BP 78/50 mmHg after a large MI • Crackles; SpO_2 88% • Cold, mottled skin • Urine 10 mL/hour	**ACTUAL** Ineffective peripheral tissue perfusion related to impaired myocardial contractility as evidenced by hypotension, cold mottled skin and oliguria.	• MAP of 65 mmHg or more and urine at least 0.5 mL/kg/hour within 6 hours.	• Keep ECG, BP (arterial line if ordered) and urine output under continuous watch. • Run inotropes and vasopressors by pump as prescribed. • Prepare for coronary reperfusion or mechanical circulatory support as ordered. • Give fluids cautiously and only if ordered.	• Outcome reached as hemodynamics improve.
Subjective *No specific complaint* **Objective** • Crackles throughout both lungs • SpO_2 88%	**ACTUAL** Impaired gas exchange related to pulmonary congestion as evidenced by crackles and hypoxemia.	• SpO_2 94% or above within 6 hours.	• Sit him up if BP allows. • Give oxygen; prepare for non-invasive or mechanical ventilation. • Give diuretics if ordered.	• Considered met at target SpO_2.
Subjective *No specific complaint* **Objective** • Low cardiac output	**RISK** Risk for impaired water-electrolyte balance as evidenced by risk factors: renal hypoperfusion and diuretic therapy.	• Electrolytes normal and creatinine stable.	• Follow electrolytes and creatinine. • Replace electrolytes as ordered.	• Goal reached if values are normal.

*References for this care plan: [11], [33], [35], [128]. Full entries are listed in the References section of the CPMN book.