15.18 Postpartum Hemorrhage*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel faint and wet." **Objective** • Blood loss over 500 mL • Soft, boggy uterus • HR 120/min, BP 90/55 mmHg	**ACTUAL** Inadequate fluid volume related to excessive blood loss from uterine atony as evidenced by heavy bleeding, a boggy uterus, tachycardia and hypotension.	• Bleeding controlled and vital signs stable within 1 hour.	• Call for help. • Massage the uterus and empty the bladder. • Give uterotonics as prescribed. • Insert two large-bore cannulas; give fluids and blood as ordered. • Measure blood loss and take vital signs every 5–15 minutes.	• Achieved when bleeding is controlled.
Subjective • "Am I going to die?" **Objective** • Frightened • Crying	**ACTUAL** Excessive fear related to heavy bleeding and emergency interventions as evidenced by expressed fear.	• Feels safer as the bleeding comes under control.	• Stay with her. • Explain what you are doing in a few words. • Keep her husband informed.	• Outcome reached when she is calmer.
Subjective *No specific complaint* **Objective** • Still losing blood	**RISK** Risk for shock as evidenced by risk factors: hemorrhage.	• MAP of 65 mmHg or more with adequate urine output.	• Insert a urinary catheter and record output. • Check clotting. • Prepare for surgery if bleeding continues.	• Considered met while she remains stable.

*References for this care plan: [35], [47], [103], [138]. Full entries are listed in the References section of the CPMN book.