15.22 Breastfeeding Support*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My baby does not latch and keeps crying." **Objective** • Shallow latch • Clicking sound during feeds • Unsettled after feeds	**ACTUAL** Ineffective chestfeeding related to poor latch technique as evidenced by a shallow latch and an unsettled infant.	• Effective latch and a settled baby within 48 hours.	• Watch a whole feed. • Teach positioning: baby held close, nose to nipple, mouth wide open. • Encourage feeding on demand, 8–12 times in 24 hours. • Teach the signs of good feeding: wet diapers and audible swallowing.	• Considered met when the baby feeds well.
Subjective • "My nipples are sore." **Objective** • Red nipples • Winces as the baby latches	**ACTUAL** Acute pain related to nipple trauma from poor latch as evidenced by sore nipples.	• Less nipple soreness within 48 hours.	• Correct the latch. • Apply expressed milk to the nipples after feeds. • Vary feeding positions.	• Goal reached as pain lessens.
Subjective *No specific complaint* **Objective** • Poor latch	**RISK** Risk for impaired skin integrity as evidenced by risk factors: repeated nipple trauma.	• Nipple skin stays intact.	• Check the latch at every feed. • Look at the nipples daily. • Let the nipples air-dry after feeds.	• Met if the skin stays intact.

* **References for this care plan:** [35], [38], [47], [134]. Full entries are listed in the References section of the CPMN book.