15.7 Gestational Hypertension and Pre-eclampsia*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My face and hands are swollen." **Objective** • BP 156/102 mmHg • Proteinuria • Swollen face and hands	**ACTUAL** Excessive fluid volume related to reduced renal perfusion and capillary leak as evidenced by edema and proteinuria.	• BP on the prescribed target without worsening edema.	• Measure BP every 4 hours or as ordered. • Test urine for protein and keep an intake and output chart. • Give antihypertensives as prescribed. • Follow liver enzymes, platelets and creatinine. • Check fetal well-being.	• Considered met when BP is controlled.
Subjective • "I am afraid for my baby." **Objective** • Anxious • Keeps asking about the baby	**ACTUAL** Excessive anxiety related to threat to self and fetus as evidenced by worry.	• Less anxious and can state the warning signs.	• Explain the plan of care and monitoring. • Teach the warning signs: severe headache, visual disturbance, upper abdominal pain, fewer fetal movements. • Let a support person stay.	• Goal reached when she is calmer.
Subjective *No specific complaint* **Objective** • Severe pre-eclampsia	**RISK** Risk for physical injury as evidenced by risk factors: possible eclamptic seizure.	• No seizures.	• Give magnesium sulfate as prescribed. • Check reflexes, RR and urine output during magnesium therapy. • Keep calcium gluconate at hand. • Keep the room quiet and dim.	• Met if no seizure occurs.

*References for this care plan: [1], [35], [47], [89]. Full entries are listed in the References section of the CPMN book.