17.19 Pyloric Stenosis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • Mother says, "He vomits forcefully after every feed." **Objective** • Projectile vomiting without bile • Olive-shaped mass felt • Sunken fontanelle; few wet diapers	**ACTUAL** Inadequate fluid volume related to persistent vomiting as evidenced by a sunken fontanelle and reduced wet diapers.	• Rehydrated with normal electrolytes before surgery.	• Nil by mouth; give IV fluids as prescribed. • Follow electrolytes and acid–base status. • Record vomiting, intake, output and weight.	• Outcome reached once he is rehydrated.
Subjective *No specific complaint* **Objective** • Weight loss • Hungry straight after vomiting	**ACTUAL** Inadequate nutritional intake related to vomiting as evidenced by weight loss.	• Tolerates feeds after pyloromyotomy.	• Restart feeds after surgery as ordered. • Burp often. • Weigh daily.	• Considered met when feeds are tolerated.
Subjective *No specific complaint* **Objective** • Losing gastric acid	**RISK** Risk for impaired water-electrolyte balance as evidenced by risk factors: vomiting of gastric contents (metabolic alkalosis, hypokalemia).	• Normal electrolytes and acid–base balance before surgery.	• Follow electrolytes. • Replace as prescribed.	• Goal reached when values are normal.

*References for this care plan: [17], [35], [38]. Full entries are listed in the References section of the CPMN book.