19.11 Incontinence in Older Adults at Home*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I cannot reach the toilet in time at night." **Objective** • Urge incontinence • Gets up twice a night to pass urine	**ACTUAL** Urge urinary incontinence related to overactive bladder as evidenced by leakage before reaching the toilet.	• Fewer episodes within 4 weeks.	• Ask her to keep a bladder diary. • Start bladder training with timed voiding. • Cut down caffeine. • Place a commode beside the bed at night.	• Considered met as episodes become fewer.
Subjective • "I am ashamed." **Objective** • Avoids visitors • Uses many pads	**ACTUAL** Situational inadequate self-esteem related to incontinence as evidenced by shame and avoidance of visitors.	• Voices her feelings and accepts help.	• Discuss discreet continence products. • Talk with her supportively. • Reassure her that treatment can help.	• Goal reached once she accepts help.
Subjective *No specific complaint* **Objective** • Damp skin • Redness of the perineum	**RISK** Risk for impaired skin integrity as evidenced by risk factors: moisture.	• Skin stays intact.	• Teach skin care and the use of barrier cream. • Change pads promptly. • Inspect the skin daily.	• Met if the skin stays intact.

* **References for this care plan:** [35], [92], [126]. Full entries are listed in the References section of the CPMN book.