6.4 Upper Gastrointestinal Bleeding*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I vomited blood and feel faint." **Objective** • Hematemesis • Melena • BP 88/56 mmHg, HR 124/min • Hb 7.5 g/dL	**ACTUAL** Inadequate fluid volume related to acute blood loss as evidenced by hypotension, tachycardia and low hemoglobin.	• Vital signs stable and Hb at target within 24 hours.	• Insert two large-bore cannulas. • Give fluids and blood as prescribed. • Take vital signs every 15 minutes until stable. • Keep him nil by mouth and prepare for endoscopy. • Give the IV PPI as prescribed. • Measure urine output hourly.	• Outcome reached when vital signs are stable.
Subjective • "Am I going to die?" **Objective** • Frightened and trembling • Holding a relative's hand	**ACTUAL** Excessive fear related to visible blood loss and hospital procedures as evidenced by expressed fear.	• Reports less fear within 2 hours.	• Stay with him. • Explain each procedure briefly and calmly. • Remove blood-stained items quickly. • Allow family to stay.	• Considered met when he is calmer.
Subjective *No specific complaint* **Objective** • Still bleeding • HR rising	**RISK** Risk for shock as evidenced by risk factors: hypovolemia.	• MAP above 65 mmHg.	• Take vital signs every 15–30 minutes. • Watch for signs of rebleeding. • Keep cross-matched blood available. • Report changes at once.	• Goal reached if shock does not develop.

*References for this care plan: [35], [46], [61], [128]. Full entries are listed in the References section of the CPMN book.