7.7 Hemodialysis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel dizzy and cramp during dialysis." **Objective** • BP drops to 90/55 mmHg during the session • Muscle cramps	**ACTUAL** Inadequate fluid volume related to rapid fluid removal during dialysis as evidenced by hypotension and cramps.	• Finishes dialysis with a stable BP.	• Weigh before and after dialysis. • Measure BP every 30 minutes during the session. • Report hypotension and reduce ultrafiltration per protocol. • Teach him to limit weight gain between sessions.	• Considered met when the session is stable.
Subjective • "I drink a lot because I am thirsty." **Objective** • Weight gain of 3.5 kg between sessions • Ankle edema	**ACTUAL** Ineffective health self-management related to difficulty adhering to fluid restriction as evidenced by excessive interdialytic weight gain.	• Weight gain between sessions within the prescribed limit.	• Suggest ways to manage thirst: ice chips, sour sweets, small cups. • Cut down sodium. • Review his weight records with him.	• Goal reached when gain is within the limit.
Subjective *No specific complaint* **Objective** • Vascular access • Cannulated repeatedly	**RISK** Risk for infection as evidenced by risk factors: vascular access and uremia-related immunosuppression.	• Access site free of infection.	• Use aseptic technique for every connection. • Inspect the access at each session. • Teach him to keep it clean and report redness.	• Met if no infection develops.

* **References for this care plan:** [32], [35], [37], [43]. Full entries are listed in the References section of the CPMN book.