7.8 Peritoneal Dialysis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "The fluid coming out looks cloudy." **Objective** • Cloudy effluent • Abdominal pain • Fever	**ACTUAL** Acute pain related to peritoneal infection (peritonitis) as evidenced by abdominal pain and cloudy effluent.	• Clear effluent and no pain within 72 hours of treatment.	• Send effluent for cell count and culture. • Give intraperitoneal antibiotics as prescribed. • Record each exchange and how the effluent looks.	• Achieved when effluent is clear.
Subjective • "How do I do the exchange at home?" **Objective** • Recently started peritoneal dialysis • Cannot describe the exchange steps	**ACTUAL** Inadequate health knowledge related to the home dialysis procedure as evidenced by questioning.	• Performs an exchange with aseptic technique.	• Teach hand hygiene, mask use and aseptic connection. • Teach exit-site care. • Teach recording of weight, BP and fluid balance.	• Outcome reached when demonstrated correctly.
Subjective *No specific complaint* **Objective** • Peritoneal catheter • Slight redness at the exit site	**RISK** Risk for infection as evidenced by risk factors: indwelling peritoneal catheter.	• Exit site free of infection.	• Clean the exit site daily. • Teach the signs of exit-site infection and peritonitis. • Reinforce hand hygiene before every exchange.	• Considered met if no infection develops.

* **References for this care plan:** [32], [35], [37], [43]. Full entries are listed in the References section of the CPMN book.