8.5 Hypoglycemia*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I feel shaky, sweaty and hungry." **Objective** • Glucose 52 mg/dL • Trembling and sweating • Poor concentration • HR 110/min	**ACTUAL** Acute confusion related to hypoglycemia as evidenced by glucose 52 mg/dL, tremor, diaphoresis and poor concentration.	• Glucose above 70 mg/dL within 15 minutes.	• If conscious: give 15–20 g of fast-acting glucose, recheck in 15 minutes, repeat if needed. • Follow with a carbohydrate snack or the next meal. • If unconscious: give IV dextrose or IM glucagon as prescribed; nothing by mouth. • Find out what caused the episode.	• Outcome reached when glucose normalizes.
Subjective • "I did not eat lunch but took my insulin." **Objective** • Missed a meal • Cannot say how to prevent hypoglycemia	**ACTUAL** Inadequate health knowledge related to matching food intake and insulin as evidenced by taking insulin without eating.	• Explains how to prevent hypoglycemia.	• Teach how to match meals with insulin. • Advise carrying glucose at all times.	• Considered met when prevention is explained.
Subjective *No specific complaint* **Objective** • Repeated hypoglycemia • Drives to work	**RISK** Risk for physical injury as evidenced by risk factors: hypoglycemic episodes with impaired cognition.	• No injury from hypoglycemia.	• Advise checking glucose before driving. • Teach the family to recognize and treat hypoglycemia, including glucagon if prescribed. • Advise wearing diabetes identification.	• Goal reached if no injury occurs.

* **References for this care plan:** [3], [32], [35], [37]. Full entries are listed in the References section of the CPMN book.