10.13 Anaphylaxis*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "My throat is closing." **Objective** • Stridor and wheeze • Swollen face and lips • Urticaria	**ACTUAL** Ineffective airway clearance related to airway edema and bronchospasm as evidenced by stridor and wheeze.	• Airway open within minutes of treatment.	• Call for help and remove the trigger. • Give IM adrenaline straight away, per protocol. • Give high-flow oxygen. • Prepare for an advanced airway if swelling worsens. • Repeat adrenaline per protocol if there is no improvement.	• Achieved when the airway is patent.
Subjective • "I feel faint." **Objective** • BP 80/40 mmHg • HR 130/min	**ACTUAL** Ineffective peripheral tissue perfusion related to vasodilation and capillary leak as evidenced by hypotension, tachycardia and faintness.	• Systolic BP above 90 mmHg within 30 minutes.	• Lay her flat with the legs raised, or sit her up if lying flat worsens breathing. • Give an IV fluid bolus as prescribed. • Keep BP, HR and SpO_2 on continuous monitoring. • After adrenaline, give the antihistamine and corticosteroid as prescribed.	• Outcome reached once BP is restored.
Subjective *No specific complaint* **Objective** • Known severe allergy • Biphasic reaction possible	**RISK** Risk for allergic reaction as evidenced by risk factors: known severe allergy.	• Avoids future exposure and uses the auto-injector correctly.	• Observe her for the period set by policy after recovery. • Record the allergy clearly in the notes and on the wristband. • Teach use of the adrenaline auto-injector. • Refer to an allergy clinic.	• Considered met when she demonstrates correctly.

* **References for this care plan:** [27], [32], [35], [128]. Full entries are listed in the References section of the CPMN book.