12.3 Retinal Detachment*

Assessment	Nursing Diagnosis	Plan / Expected Outcome	Nursing Interventions	Evaluation
Subjective • "I see flashes and a curtain over my vision." **Objective** • Sudden loss of part of the visual field • Floaters	**ACTUAL** Excessive anxiety related to sudden vision loss as evidenced by fear and questioning.	• Feels less anxious before surgery.	• Explain why treatment is urgent and what the procedure involves. • Keep the environment calm and let her rest. • Allow a relative to stay.	• Achieved if she appears calmer.
Subjective • "How long must I keep my head down?" **Objective** • Intraocular gas bubble after vitrectomy • Specific head position prescribed	**ACTUAL** Inadequate health knowledge related to post-operative positioning as evidenced by questioning.	• Keeps to the prescribed head position.	• Teach the exact position and how long the surgeon wants it held. • Warn against flying or travel to high altitude while gas remains in the eye. • Provide pillows and supports to make the position bearable.	• Outcome reached when the position is maintained.
Subjective *No specific complaint* **Objective** • Fixed positioning • Reduced vision	**RISK** Risk for impaired skin integrity as evidenced by risk factors: prolonged fixed positioning.	• Skin over pressure points stays intact.	• Look at the face and chest where they bear weight. • Pad these areas.	• Considered met if the skin is intact.

*References for this care plan: [19], [32], [35], [37]. Full entries are listed in the References section of the CPMN book.